Mississippi Trustee Implementation Group Draft 2016-2017 Restoration Plan/Environmental Assessment:

Restoration of Wetlands, Coastal, and Nearshore Habitats; Birds; and Nutrient Reduction (Nonpoint Source)

December 2016

EXECUTIVE SUMMARY

On April 20, 2010, the Deepwater Horizon (*DWH*) mobile drilling unit exploded, caught fire, and eventually sank in the Gulf of Mexico, resulting in a massive release of oil and other substances from BP Exploration and Production Inc.'s Macondo well. Initial efforts to cap the well following the explosion were unsuccessful, and for 87 days after the explosion, the well continuously and uncontrollably discharged oil and natural gas into the northern Gulf of Mexico. Approximately 3.19 million barrels (134 million gallons) of oil were released into the ocean (U.S. v. BP et al. 2015), by far the largest offshore oil spill in the history of the United States. Oil spread from the deep ocean to the surface and nearshore environment across the northern Gulf of Mexico. Extensive response actions, including cleanup activities and actions to try to prevent the oil from reaching sensitive resources, were undertaken to try to reduce harm to people and the environment. However, many of these response actions had collateral impacts on the environment. The oil and other substances released from the well in combination with the extensive response actions together make up the *DWH* oil spill.

As an oil pollution incident, the *DWH* oil spill was subject to the provisions of the Oil Pollution Act (OPA) of 1990, which addresses preventing, responding to, and paying for oil pollution incidents in navigable waters, adjoining shorelines, and the exclusive economic zone of the United States. The primary goal of OPA is to make the environment and public whole for injuries to natural resources and services resulting from an incident involving an oil discharge (or substantial threat of an oil discharge). Under the authority of OPA, a council of federal and state *DWH* oil spill Trustees (the Trustees) was established, on behalf of the public, to assess natural resource injuries resulting from the incident and work to make the environment and public whole for those injuries. As required under OPA, the Trustees conducted a natural resource damage assessment (NRDA) to:

- Assess the impacts of the DWH oil spill on natural resources in the Gulf of Mexico and the services those resources provide, and
- Determine the type and amount of restoration needed to compensate the public for these impacts.

Following the assessment, the Trustees determined that the injuries caused by the *DWH* oil spill could not be fully described at the level of a single species, a single habitat type, or a single region. Rather, the injuries affected such a wide array of linked resources over such an enormous area that the effects of the *DWH* oil spill must be described as constituting an ecosystem-level injury.

Given the broad ecological scope of the injuries, restoration planning requires a broad ecosystem perspective to restore the vast array of resources and services injured by the *DWH* oil spill. Thus, the Trustees proposed a comprehensive, integrated ecosystem restoration approach in their programmatic level restoration plan (the Programmatic Damage Assessment and Restoration Plan/Programmatic Environmental Impact Statement, or PDARP/PEIS) to guide and direct the massive restoration effort. The PDARP/PEIS includes a portfolio of Restoration Types that addresses the diverse suite of injuries that occurred at both regional and local scales, and is based on the following five overarching goals:

- Restore and conserve habitat;
- Restore water quality;

- Replenish and protect living coastal and marine resources;
- Provide and enhance recreational opportunities; and
- Provide for monitoring, adaptive management, and administrative oversight to support restoration implementation.

These five goals work both independently and together to restore injured resources and services.

Draft Restoration Plan/Environmental Assessment

This document, the "Mississippi Trustee Implementation Group Draft 2016-2017 Restoration Plan/Environmental Assessment" (Draft RP/EA), was prepared by the Mississippi Trustee Implementation Group (MS TIG) pursuant to OPA and is consistent with the Trustees' findings in the PDARP/PEIS. The MS TIG includes one state trustee agency and four federal trustee agencies: the Mississippi Department of Environmental Quality (MDEQ); the United States Department of Commerce, represented by the National Oceanic and Atmospheric Administration (NOAA); the United States Department of the Interior (DOI), represented by the United States Fish and Wildlife Service (USFWS) and the National Park Service (NPS); the United States Department of Agriculture (USDA); and the Environmental Protection Agency (EPA).

The MS TIG prepared this Draft RP/EA to inform the public about *DWH* NRDA restoration planning efforts and to seek public comment on three preferred restoration alternatives.

In identifying proposed projects for this Draft RP/EA, the MS TIG considered projects submitted by the public via the MDEQ Restoration Project Idea portal¹ and the Trustee Project Submission Portal² as well as those proposed in response to the May 27, 2016 MS TIG Notice of Initiation of Restoration Planning.

In developing a reasonable range of alternatives suitable for addressing the injuries caused by the DWH oil spill, the MS TIG reviewed PDARP Trustee Restoration Goals, MS TIG 2016-2017 Goals and Objectives (which are detailed in 2.4.1.2 of this document), and OPA screening criteria. In addition, the MS TIG considered current and future availability of funds under the DWH oil spill NRDA settlement payment schedule, as well as projects already funded or proposed to be funded by the other DWH restoration funding sources of the National Fish and Wildlife Foundation Gulf Environment Benefit Fund (NFWF GEBF) and the Resources and Ecosystems Sustainability, Tourist, Opportunities, and Revived Economies of the Gulf Coast States (RESTORE) Act. The MS TIG then identified three restoration types - Wetlands, Coastal, and Nearshore Habitats (WCNH), Birds, and Nutrient Reduction (NR) (Nonpoint Source), as appropriate for focus in this plan. Preservation, restoration, regional connectivity and proximity to state and federal conservation lands were key drivers in determining restoration approaches/techniques and in screening of projects to develop a reasonable range of alternatives. Under the Consent Decree discussed in Section 1.1 of this Draft RP/EA, the three largest Natural Resource Damage (NRD) funding categories that will be made available to the MS Restoration Area are to be utilized for the following three Restoration Types: WCNH (\$55,500,000); NR (Nonpoint Source) (\$27,5000,000); and Birds (\$25,000,000). For this

_

¹ http://www.restore.ms/submit-project-idea/

² http://www.gulfspillrestoration.noaa.gov/restoration/give-us-your-ideas/suggest-a-restoration-project/

Draft RP/EA, the MS TIG has chosen to prioritize restoration project alternatives which correspond to these three Restoration Types.

Table ES-1: Settlement of NRD Claims; NRD final allocation to the MS TIG

Major Restoration Categories	Mississippi Restoration Area
Restore and Conserve Habitat	-
Wetlands, Coastal and Nearshore Habitats	55,500,000
Habitat Projects on Federally Managed Lands	5,000,000
Early Restoration (through Phase IV)	80,000,000
Restore Water Quality	-
Nutrient Reduction (Nonpoint Source)	27,5000,000
Replenish and Protect Living Coastal and Marine Resources	-
Sea Turtles	5,000,000
Marine Mammals	10,000,000
Birds	25,000,000
Oysters	20,000,000
Early Restoration Oysters	13,600,000
Provide and Enhance Recreational Opportunities	-
Provide and Enhance Recreational Opportunities	5,000,000
Early Restoration Recreational Opportunities	18,957,000
Monitoring, Adaptive Management, Administrative Oversight	-
Monitoring and Adaptive Management	7,500,000
Administrative Oversight and Comprehensive Planning	22,500,000
Total NRD Funding	\$295,557,000

Section 2 of this Draft RP/EA describes the screening of projects and development of the reasonable range of alternatives for WCNH/Birds and for NR (Nonpoint Source) Restoration Types. Section 3.0 provides the OPA evaluation and NEPA Affected Environment for the reasonable range of alternatives for these Restoration Types. For this Draft RP/EA, the MS TIG proposes moving forward with the following preferred alternatives or projects within the WCNH and Birds Restoration Types: Graveline Bay Land Acquisition and Management and Grand Bay Land Acquisition and Habitat Management, and the following preferred alternative within the NR (Nonpoint Source) Restoration Type: Upper Pascagoula River Water Quality Enhancement.

The three preferred alternatives (projects) proposed in this Draft RP/EA are set forth in Table ES-2 below. The geographic locations of the three proposed alternatives (projects) are depicted below in Figure ES-1. More information about each of these projects, as well as other projects evaluated by the MS TIG, can be found in Section 2 of this Draft RP/EA.

Table ES-2: Proposed Preferred Restoration Alternatives in this Draft RP/EA

Preferred Alternatives/Projects	PDARP/PEIS Restoration Goal and Restoration Type	Proposed Funding
Graveline Bay Land Acquisition and Management	Restore and Conserve Habitat: Wetlands, Coastal and Nearshore Habitats Replenish and Protect Living Coastal and Marine Resources: Birds	\$11,500,000
Grand Bay Land Acquisition and Habitat Management	Restore and Conserve Habitat: Wetlands, Coastal and Nearshore Habitats Replenish and Protect Living Coastal and Marine Resources: Birds	\$6,000,000
Upper Pascagoula River Water Quality Enhancement	Restore Water Quality: NR (Nonpoint Source)	\$4,000,000




Figure ES-1: Locations of Preferred Proposed Alternatives

At this time, the MS TIG proposes moving forward with the three preferred restoration alternatives (projects) in this Draft RP/EA. Pursuant to the National Environmental Policy Act of 1969 (NEPA), an evaluation of environmental consequences is discussed in the PDARP/PEIS and incorporated by reference into this Draft RP/EA, and is also discussed in Section 3.0. Environmental consequences to the physical environment, the biological environment, and the socioeconomic environment are evaluated in this Draft RP/EA (Section 3.3.1, 3.4.1, and 3.9.1). The findings are summarized below.

WCNH/Birds Proposed Alternatives-Environmental Consequence Summary

In addition to land acquisition, proposed habitat restoration measures and management activities for the proposed WCNH/Birds alternatives includes invasive species management via chemical treatment, mechanical treatment, and prescribed fire; access restriction; road repair/removal and culvert placement, and debris removal. Land acquisition and implementation of these restoration measures and management activities would have **short-term**, **minor**, **adverse impacts** to noise, tourism and recreation and public health and safety. There would be **short-term**, **minor to moderate**, **adverse impacts** to hydrology, water quality, wetlands, air quality and greenhouse gases, habitat and wildlife from ground-disturbing activities associated with habitat restoration measures and management activities. Depending on the alternative, the adverse impacts to soils would range from **long-term**, **minor**, due to allowing public access on previously private land, to **short-term**, **minor to moderate** due to habitat management activities. Land acquisition could have a **short-term**, **minor to moderate**

adverse impact on socioeconomic resources due to changes in visitor spending and loss of tax revenues. There would be a **long-term, minor to moderate**, adverse impacts to land and marine management.

There would be **long-term, benefits** to soil, hydrology, floodplains, wetlands, water quality, and habitat and wildlife land and marine management, tourism and recreation, and public health and safety, due to preservation of habitats and floodplains, re-establishment of native plant communities, increased diversity in flora and fauna, implementation of existing resource management plans/initiatives, and the potential for increased visitor use.

NR (Nonpoint Source) Proposed Alternative-Environmental Consequence Summary

Ecological/NR conservation practices and soil and water conservation/NR practices with willing participants, would provide a wide array of benefits to cropland, pasture/grassland, associated agriculture lands and riparian areas. There would be **short-term, minor to moderate**, adverse impacts to soils, water quality, wetlands and habitats and wildlife. The adverse impacts to hydrology would range from **long-term, minor**, due to conservation practices that may require in stream work, to **short-term, minor to moderate** due to non in stream conservation practices. **short-term to long-term, minor to moderate**, adverse impacts to hydrology. Conservation planning and the implementation of conservation practices on privately owned lands would reduce nutrient enrichment and sedimentation and restore water quality in Gulf of Mexico coastal wtaersheds. Conservation practices would provide **long-term benefits** to soil, hydrology, water quality and wetlands, habitat and wildlife, socioeconomic resources, and public health and safety.

Table of Contents

1.0 Introduction	11
1.1 Background and Summary of Settlement	11
1.2 DWH Oil Spill Trustees	12
1.3 Authorities and Regulations	14
1.3.1 OPA and NEPA Compliance	14
1.3.2 PDARP/PEIS Record of Decision	14
1.3.3 Relationship of this Draft RP/EA to the PDARP/PEIS	15
1.3.4 Restoration Planning Context	16
1.4 Draft RP/EA	18
1.5 Purpose and Need	18
1.6 Proposed Action: MS TIG Draft 2016-2017 RP/EA	19
1.7 Public Involvement	20
1.7.1 Public Involvement in the Development of this Draft RP/EA and Next S	Steps21
1.7.2 Administrative Record	21
1.8 Severability of Projects	22
1.9 Decisions to be Made	22
1.10 Document Organization	22
2.0 Restoration Planning Process	23
2.1 Injuries Addressed in this Draft RP/EA	23
2.1.1 Wetlands, Coastal, and Nearshore Habitats	23
2.1.2 Nutrient Reduction (Nonpoint Source)	24
2.1.3 Birds 24	
2.2 DWH Early Restoration Addressing the Injuries to Date	24
2.3 Coordination with Other Gulf Restoration Programs	26
2.4 Screening for Potential Alternatives	26
2.4.1 MS TIG Screening Process	26
2.4.1.1 Consistency with Goals outlined in PDARP/PEIS Restoration Type	pes27
2.4.1.2 MS TIG 2016-2017 Goals and Objectives	29
2.4.1.3 Identification of PDARP/PEIS Restoration Approaches/Technique	es30
2.4.1.4 OPA Screening	32
2.4.1.4.1 WCNH/Birds	32
2.4.1.4.2 NR (Nonpoint Source)	34

2.5	Alterna	atives not Considered for Further Evaluation in this Draft RP/EA	35
2.6	Reason	able Range of Restoration Alternatives Considered	36
2	2.6.1 Res	toration Type: WCNH/Birds	36
	2.6.1.1	Alternative A (Preferred): Graveline Bay Land Acquisition and Managen	nent 38
	2.6.1.2	Alternative B: Grand Bay Land Acquisition (up to 8,000 acres)	39
	2.6.1.3	Alternative C: Grand Bay Habitat Management (up to 17,500 acres)	39
	2.6.1.4	Alternative D (Preferred): Grand Bay Land Acquisition (up to 8,000 acre Habitat Management (up to 17,500 acres)	
	2.6.1.5	Natural Recovery/No Action	40
2	2.6.2 Rest	oration Type NR (Nonpoint Source)	40
	2.6.2.1	Alternative A (Preferred): Upper Pascagoula River Water Quality Enhance	ement40
	2.6.2.2	Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan	ı41
	2.6.2.3	Natural Recovery/No Action	41
2.7	Alterna	ntives Evaluated in this Draft RP/EA	41
3.0		valuation of Alternatives and NEPA Affected Environment and Environment	
	•		
3.1		I/Bird Restoration Type	
		A Evaluation for WCNH/Birds	
		PA Analysis for WCNH/Birds Restoration Type	
3.2		l Recovery/No Action	
3.3		ine Bay Land Acquisition and Management-Background and Project Descri	-
3		rnative A (Preferred): Graveline Bay Land Acquisition and Management Aironment and Environmental Consequences	
	3.3.1.1	Overview of Affected Environment and Environmental Consequences	57
	3.3.1.2	Physical Environment	58
	3.3.1.2	2.1 Geology and Substrates	60
	3.3.1.2	2.2 Hydrology and Water Quality	63
	3.3.1.2	2.3 Air Quality and Greenhouse Gas Emissions	68
	3.3.1.3	Biological Environment	70
	3.3.1.3	3.1 Habitats	73
	3.3.1.3	3.2 Protected Species	77
	3.3.1.3	3.3 Migratory Birds	84
	3.3.1.3	3.2 Wildlife	87
	3.3.1.4	Socioeconomic Resources	88
	3.3.1.4	4.1 Socioeconomic Resources and Environmental Justice	91

3.3.1.4.2 Tourism and Recreational Use	92
3.3.1.4.3 Cultural Resources	93
3.3.1.4.4 Land and Marine Management	94
3.3.1.4.5 Public Health and Safety	96
3.3.2 Site-Specific NEPA Review for WCNH/Birds Proposed Alternative A	(Preferred) 97
3.4 Grand Bay Land Acquisition & Habitat Management-Background & Pro	ject Description97
3.4.1 Grand Bay Land Acquisition and Habitat Management Alternatives B Environment and Environmental Consequences	
3.4.1.1 Overview of Affected Environment and Environmental Consequ	iences 104
3.4.1.2 Physical Environment	104
3.4.1.2.1. Geology and Substrates	105
3.4.1.2.2 Hydrology and Water Quality	109
3.4.1.2.3. Air Quality and Greenhouse Gas Emissions	115
3.4.1.3 Biological Environment	118
3.4.1.3.1. Habitats	119
3.4.1.3.2 Protected Species	124
3.4.1.3.3 Migratory Birds	131
3.4.1.3.4 Wildlife	134
3.4.1.4 Socioeconomic Environment	135
3.4.1.4.1 Socioeconomic Resources and Environmental Justice	137
3.4.1.4.2 Tourism and Recreational Use	139
3.4.1.4.3 Cultural Resources	140
3.4.1.4.4 Land and Marine Management	141
3.4.1.4.5 Public Health and Safety	142
3.4.2. Site-specific NEPA Review for WCNH/Birds Proposed Alternatives I 143	3, C & D-(Preferred)
3.5 Cumulative Impacts for WCNH/Birds Alternatives	144
3.5.1 Identification of Resources Affected	144
3.5.1.1 Cumulative Action Scenario	145
3.5.1.2 Cumulative impact Analysis	146
3.6 Comparison of the Alternatives-WCNH/Birds Restoration Type	148
3.7 NR (Nonpoint Source) Restoration Type	154
3.7.1 OPA Evaluation for NR (Nonpoint Source)	154
3.7.2 NEPA Analysis for NR (Nonpoint Source) Restoration Type	159
3.8 Natural Recovery/No Action Alternative	161

Approa	ach
3.9	1.1 NR (Nonpoint Source) Alternatives A and B: -Affected Environment and Environmental Consequences
:	3.9.1.1 Overview of Affected Environment and Environmental Consequences
	3.9.1.2 Physical Environment
	3.9.1.2.1 Geology and Substrates
	3.9.1.2.2 Hydrology and Water Quality
	3.9.1.3 Biological Environment
	3.9.1.3.1 Habitats and Wildlife
	3.9.1.3.2 Protected Species
	3.9.1.3.3 Migratory Birds
	3.9.1.4 Socioeconomic Resources
	3.9.1.4.1 Socioeconomics and Environmental Justice
	3.9.1.4.2 Cultural Resources 199
3.9	2.2 Site-specific NEPA Review for NR Proposed Alternatives A (Preferred) and B 201
3.10	Cumulative Impacts for NR
3.1	0.1 Identification of Resources Affected
3.1	0.2 Cumulative Action Scenario
3.1	0.3 Cumulative impact analysis
3.11	Comparison of the Alternatives-NR (Nonpoint Source) Restoration Type204
4.0	Compliance with Other Laws and Regulations
4.1	Additional Federal Laws
4.2	Additional State Laws
5.0	Draft Monitoring and Adaptive Management Plan213
5.1	Introduction
5.2	Summary of Restoration Goals, Objectives, and Performance Criteria213
5.3	MAM Plan Administration
6.0	List of Preparers and Reviewers
7.0	Literature Cited

1.0 Introduction

This "Mississippi Trustee Implementation Group Draft 2016-2017 Restoration Plan/Environmental Assessment" (Draft RP/EA) was prepared by the federal and state natural resource trustees for the Mississippi Trustee Implementation Group (MS TIG), which is responsible for restoring the natural resources and services within the Mississippi Restoration Area that were injured by the April 20, 2010, Deepwater Horizon oil spill and associated spill response efforts (*DWH* oil spill). The purpose of restoration, as discussed in this document and detailed more fully in the Programmatic Damage Assessment and Restoration Plan/Programmatic Environmental Impact Statement (PDARP/PEIS), is to make the environment and the public whole for injuries resulting from the incident by implementing restoration actions that return injured natural resources and services to baseline conditions and compensate for interim losses, in accordance with the Oil Pollution Act of 1990 (OPA) and associated Natural Resource Damage Assessment (NRDA) regulations.

The MS TIG includes one state trustee agency and four federal trustee agencies: the Mississippi Department of Environmental Quality (MDEQ); the United States Department of Commerce, represented by the National Oceanic and Atmospheric Administration (NOAA); the United States Department of the Interior (DOI), represented by the United States Fish and Wildlife Service (USFWS) and the National Park Service (NPS); the United States Department of Agriculture (USDA); and the Environmental Protection Agency (EPA) (collectively the MS TIG). The MS TIG has prepared this Draft RP/EA to inform the public about *DWH* NRDA restoration planning efforts and to seek public comment on the proposed restoration alternatives.

1.1 Background and Summary of Settlement

On April 4, 2016, the United States District Court for the Eastern District of Louisiana entered a Consent Decree resolving civil claims by the *DWH* oil spill trustees (Trustees) against BP Exploration and Production Inc. (BP) arising from the *DWH* oil spill. This historic settlement resolves the Trustees' claims against BP for natural resources damages under OPA.

Under the Consent Decree, BP agreed to pay over a 15-year period a total of \$8.1 billion in natural resource damages (which includes \$1 billion that BP previously committed to pay for early restoration projects), and up to an additional \$700 million (some of which is in the form of accrued interest) for adaptive management or to address injuries to natural resources that are presently unknown but may come to light in the future.

Table 1.1-1 below³ outlines the settlement of NRDA claims; this Table provides the final allocation for the MS Restoration Area under NRDA. The total NRDA funding for the Mississippi Restoration Area is \$295,557,000, and the total remaining NRDA allocation for the Mississippi Restoration Area is \$183 million.

More details on the background of the *DWH* oil spill, the impact of the spill on the Gulf of Mexico ecosystem, and additional context for the settlement and allocation of funds can be found in Chapter 2 of the PDARP/PEIS.

³ Table 1.1-1 is a modified version of Table 5.10-1 of the PDARP/PEIS.

Table 1.1-1: Settlement of NRD claims; NRD final allocation

Major Restoration Categories	Mississippi
Wetlands, Coastal, and Nearshore Habitats	\$55,500,000
Habitat Projects on Federally Managed Lands	\$5,000,000
Early Restoration (through Phase IV)	\$80,000,000
Nutrient Reduction (Nonpoint Source)	\$27,500,000
Water Quality (e.g., Stormwater Treatments, Hydrologic Restoration, Reduction of Sedimentation, etc.)	
Fish and Water Column Invertebrates	
Early Restoration Fish and Water Column Invertebrates	
Sturgeon	
Sea Turtles	\$5,000,000
Early Restoration Turtles	
Submerged Aquatic Vegetation	
Marine Mammals	\$10,000,000
Birds	\$25,000,000
Early Restoration Birds	
Mesophotic and Deep Benthic Communities	
Oysters	\$20,000,000
Early Restoration Oysters	\$13,600,000
Provide and Enhance Recreational Opportunities	\$5,000,000
Early Restoration Recreational Opportunities	\$18,957,000
Monitoring and Adaptive Management	\$7,500,000
Administrative Oversight and Comprehensive Planning	\$22,500,000
Adaptive Management NRD Payment for Unknown Conditions	
Total NRD Funding	\$295,557,000

1.2 *DWH* Oil Spill Trustees

As specified in OPA, natural resource trustees are designated to act on behalf of the public to assess and recover damages, develop implementation plans, and implement restoration plans (see Section 7.1 of the Final PDARP/PEIS for further detail). Trustees fulfill these responsibilities by developing restoration plans, providing the public with meaningful opportunities to review and comment on proposed plans (including the information that supports that purpose), implementing and monitoring restoration projects, managing natural resource damage funds, documenting trustee decisions through a public Administrative Record (including those that involve the use of recovered damages), and providing for public involvement and transparency in keeping with the public responsibilities with which they have each been entrusted under OPA.

The DWH Trustees are the government entities authorized under OPA to act as trustees on behalf of

the public to assess the natural resource injuries resulting from the *DWH* oil spill and develop and implement a restoration plan to compensate for those injuries. Collectively, these trustees comprise the *DWH* Trustee Council. The following federal and state agencies are the designated Trustees under OPA for the *DWH* oil spill:

- U.S. Department of the Interior (DOI), as represented by the National Park Service (NPS), U.S. Fish and Wildlife Service (FWS), and Bureau of Land Management (BLM)
- National Oceanic and Atmospheric Administration (NOAA), on behalf of the U.S. Department of Commerce (DOC)
- U.S. Department of Agriculture (USDA)
- U.S. Department of Defense (DOD)⁴
- U.S. Environmental Protection Agency (EPA)
- The State of Alabama's Department of Conservation and Natural Resources (ADCNR) and Geological Survey of Alabama (GSA)
- The State of Florida's Department of Environmental Protection (FDEP) and Fish and Wildlife Conservation Commission (FWC)
- The State of Louisiana's Coastal Protection and Restoration Authority (CPRA) Department of Natural Resources (LDNR); Department of Environmental Quality (LDEQ); Oil Spill Coordinator's Office (LOSCO); and Department of Wildlife and Fisheries (LDWF)
- The State of Mississippi's Department of Environmental Quality (MDEQ)
- The State of Texas' Parks and Wildlife Department (TPWD), General Land Office (TGLO), and Commission on Environmental Quality (TCEQ)

Trustee Implementation Groups (TIGs) are established by the *DWH* Settlement agreement and are composed of Individual Trustee Agency representatives. The TIGs develop plans for, choose, and implement specific restoration actions under the Final PDARP/PEIS. Each TIG makes all restoration decisions for the funding allocated to its Restoration Area, and ensures its actions. The following state and federal agencies are the MS TIG:

- Mississippi Department of Environmental Quality
- U.S. Department of the Interior (DOI), as represented by the National Park Service (NPS), U.S. Fish and Wildlife Service (FWS), and Bureau of Land Management (BLM)
- National Oceanic and Atmospheric Administration (NOAA), on behalf of the U.S. Department of Commerce (DOC)
- U.S. Department of Agriculture (USDA)
- U.S. Environmental Protection Agency (EPA)

⁴ Although a trustee under OPA by virtue of the proximity of its facilities to the *Deepwater Horizon* oil spill, DOD is not a member of the Trustee Council and does not participate in DWH Trustee decision-making.

1.3 Authorities and Regulations

1.3.1 OPA and NEPA Compliance

As an oil pollution incident, the *DWH* oil spill is subject to the provisions of OPA, 33 U.S.C. § 2701 *et seq.* A primary goal of OPA is to make the environment and public whole for injuries to natural resources and services resulting from an incident involving an oil discharge or substantial threat of an oil discharge. Under OPA, each party responsible for a vessel or facility from which oil is discharged, or which poses the substantial threat of a discharge, is liable for, among other things, removal costs and damages for injury to, destruction of, loss, or loss of use of natural resources, including the reasonable cost of assessing the damage.

This process of injury assessment and restoration planning is referred to as NRDA. Under the authority of OPA, a council of federal and state trustees was established to assess natural resource injuries resulting from the incident and to work to make the environment and public whole for those injuries. NRDA is described under Section 1006 of OPA (33 U.S.C. § 2706). Under the OPA NRDA regulations (15 C.F.R. Part 990), the NRDA process consists of three phases: 1) Preassessment; 2) Assessment and Restoration Planning; and 3) Restoration Implementation. The *DWH* Trustees are currently in the Restoration Implementation phase of the NRDA. As part of the initiation of restoration implementation, this Draft RP/EA identifies a reasonable range of restoration alternatives, evaluates those alternatives under various criteria, and proposes a suite of preferred alternatives.

Restoration activities under OPA are intended to return injured natural resources and services to their baseline condition (primary restoration) and to compensate the public for interim losses from the time of the incident until the time resources and services recover to baseline conditions (compensatory restoration). To meet these goals, the restoration activities need to produce benefits that are related to or have a nexus (connection) to natural resource injuries and service losses resulting from the *DWH* oil spill.

Under the OPA regulations, federal trustees must comply with National Environmental Policy Act (NEPA), 42 U.S.C. § 4321 *et seq.*, and its regulations, 40 C.F.R. § 1500 *et seq.*, when planning restoration projects. NEPA requires federal agencies to consider the potential environmental impacts of planned actions. NEPA provides a mandate and framework for federal agencies to determine if their proposed actions have significant environmental effects and related social and economic effects, consider these effects when choosing between alternative approaches, and inform and involve the public in the environmental analysis and decision-making process.

More information about OPA and NEPA, as well as their application to *DWH* oil spill restoration planning, can be found in Chapters 5 and 6 of the PDARP/PEIS.

1.3.2 PDARP/PEIS Record of Decision

Given the potential magnitude and breadth of restoration for injuries resulting from the *DWH* oil spill, the Trustees prepared a PDARP/PEIS under OPA and NEPA to analyze alternative approaches to implementing restoration and to consistently guide restoration decisions. Based on the Trustees' thorough assessment of impacts to the Gulf's natural resources, a comprehensive, integrated ecosystem restoration approach for restoration implementation was proposed. On February 19, 2016,

the *DWH* Trustee Council issued a Final PDARP/PEIS detailing a specific proposed plan to fund and implement restoration projects across the Gulf of Mexico region over the next 15 years. On March 29, 2016, in accordance with OPA and NEPA, the Trustees published a Notice of Availability of a Record of Decision (ROD) for the PDARP/PEIS in the Federal Register (81 Fed. Reg. 17438). Based on the Trustees' injury determination established in the PDARP/PEIS, the ROD set forth the basis for the Trustees' decision to select Alternative A: Comprehensive Integrated Ecosystem Alternative. The Trustees' selection of Alternative A includes the funding allocations established in the PDARP/PEIS.

More information about Alternative A can be found in Sections 5.5 and 5.10 of the PDARP/PEIS.

1.3.3 Relationship of this Draft RP/EA to the PDARP/PEIS

As a programmatic restoration plan, the PDARP/PEIS provides direction and guidance for identifying, evaluating, and selecting future restoration projects to be carried out by the TIGs (PDARP/PEIS Section 5.10.4 and Chapter 7). The Trustees elected to prepare a PEIS to support analysis of the environmental consequences of the selected Restoration Types, to consider the multiple related actions that may occur because of restoration planning efforts, and to allow for a better analysis of cumulative impacts of potential actions. The programmatic approach was taken to assist the TIGs in their development and evaluation and to assist the public in its review of future restoration projects.

For the PDARP/PEIS, the Trustees developed a set of Restoration Types for inclusion in programmatic alternatives, consistent with the desire to seek a diverse set of projects providing benefits to a broad array of potentially injured resources and services they provide. Ultimately, this process resulted in the inclusion of thirteen (13) Restoration Types in the five major Programmatic Trustee Goals including:

- 1) Wetlands, Coastal, and Nearshore Habitats (WCNH)
- 2) Habitat Projects on Federally Managed Lands
- 3) Nutrient Reduction (Nonpoint Source)
- 4) Water Quality (e.g., Stormwater Treatments, Hydrologic Restoration, Reduction of Sedimentation)
- 5) Fish and Water Column Invertebrates
- 6) Sturgeon
- 7) Submerged Aquatic Vegetation
- 8) Oysters
- 9) Sea Turtles
- 10) Marine Mammals
- 11) Birds

12) Mesophotic and Deep Benthic Communities

13) Provide and Enhance Recreational Opportunities

⁵ Available at http://www.gulfspillrestoration.noaa.gov/sites/default/files/wp-content/uploads/PDARP ROD Final-All-Signatures 508.pdf.

For this Draft RP/EA, the MS TIG considered and evaluated proposed alternatives within the following Restoration Types: 1) WCNH; 2) Birds; and 3) Nutrient Reduction (NR) (Nonpoint Source), as described in Section 1.3.4 below. Section 2 of this Draft RP/EA summarizes the injuries addressed and the screening process used to develop a reasonable range of alternatives. The reasonable range of alternatives is consistent with the Trustees' selected programmatic alternative in the PDARP/PEIS, the Consent Decree and OPA. The MS TIG also prepared a NEPA analysis for each of the reasonable range of alternatives (Section 3.0 of this document) which tiers from the PDARP/PEIS programmatic NEPA analysis.

1.3.4 Restoration Planning Context

In 2015, MDEQ began development of the Mississippi Gulf Coast Restoration plan (MGCRP)⁶, which sets forth a coordinated, systematic, and transparent process for sustainable ecological restoration in Mississippi to restore injuries from the *DWH* oil spill.

MDEQ engaged stakeholders throughout the development of the first draft of the MGCRP. Numerous meetings were held with community and non-governmental organizations to share and highlight the individual organization's restoration priorities and objectives. MDEQ also held a series of Community Conversations to ascertain information on individual and organizational values, characteristics, and visions associated with coastal restoration. Utilizing the results of the Community Conversations, MDEQ hosted a series of Resource Summits aimed at a technical audience to provide information on Mississippi's planning tools as well as to refine the priorities identified by the public in earlier engagement efforts. The first draft of the MGCRP was released for public review and input in October 2015. MDEQ hosted a public webinar to present the MGCRP and solicit feedback for improvement.

The MGCRP included development of the Mississippi Comprehensive Ecosystem Restoration Tool (MCERT), a science-based tool that is now in place for identifying and examining ecological resources and threats for improved restoration planning and project sustainability. The MGCRP also includes the Decision Support System (DSS), which is a linear thought process that allows MDEQ to make informed, science-based decisions for enhancing, protecting, or restoring the ecological integrity of coastal Mississippi. MCERT and the DSS are the tools which allow Mississippi to use a comprehensive ecosystem approach to restoration project planning across *DWH* funding streams (NFWF, RESTORE and NRDA). The MGCRP identified three general restoration program areas: Land Resources, Coastal and Marine Resources, and Water Resources. Table 1.3-1 below illustrates the common restoration themes in the MGCRP and the PDARP/PEIS.

⁶ Funded by the National Fish and Wildlife Foundation Gulf Environmental Benefit Fund

Table 1.3-1: The Mississippi Gulf Coast Restoration Plan as Related to PDARP/PEIS Restoration Types

The Mississippi Gulf Coast Restoration Plan; A Path Toward Sustainable Ecosystem Restoration	PDARP/PEIS Restoration Types- May 27th Public Notification			
Programs and Objectives	Wetlands, Coastal and NS Habitats	Birds	Oysters	Nutrient Reduction
Land Resources Program	, ,			
Objective 1: Conserve Priority Habitats	X	X		X
Objective 2: Manage and Restore Priority Habitats	X	X		X
Coastal and Marine Resources Program				
Objective 1: Protect and Restore Marine Habitats			X	
Objective 2: Sustainably manage and enhance coastal and marine resource populations		X	X	
Water Resources Program				
Objective 1: Reduce rural non-point source pollution				X
Objective 2: Reduce urban non-point source pollution				X

On May 27, 2016, the MS TIG published a notice to invite public input regarding natural resource restoration opportunities in Mississippi for the 2016/2017 planning years. The notice indicated a focus on the following range of potential Restoration Types, which may have benefits to living coastal and marine resources:

- 1) restoration of WCNH;
- 2) restoration of water quality through NR (Nonpoint source);
- 3) restoration of Birds; and
- 4) restoration of Oysters.

Because there are several ongoing or completed projects benefitting oysters and secondary productivity in the Mississippi Restoration Area⁷, the MS TIG chose not to prioritize the oyster restoration type in this Draft RP/EA. However, oyster restoration projects will be considered in future MS TIG funding plans.

On October 31, 2016, MDEQ published a <u>Notice of Initiation for Restoration Plan Drafting in Mississippi</u>, indicating an intention to focus on the following restoration types:

- 1) WCNH;
- 2) NR (Nonpoint Source); and
- 3) Birds.

_

⁷ Early Restoration Phase I Mississippi Artificial Reef Habitat and the Mississippi Oyster Cultch Restoration projects; Early Restoration Phase IV Restoring Living Shorelines and Reefs in Mississippi Estuaries project; and the NFWF Oyster Restoration and Management Phase I project

1.4 Draft RP/EA

The MS TIG prepared this Draft RP/EA in accordance with the PDARP/PEIS, the ROD, OPA and NEPA. This Draft RP/EA describes the *DWH* NRDA restoration planning process, identifies a reasonable range of restoration alternatives to address a portion of the injuries to resources and habitats caused by the *DWH* oil spill, and proposes from those alternatives a suite of preferred restoration alternatives proposed by the MS TIG. In accordance with the Trustee Council Standard Operating Procedures and 40 C.F.R. § 1501.5, the MS TIG designated USDA as the lead federal agency responsible for NEPA compliance for this Draft RP/EA. Each of the other federal and state co-Trustees are participating as a cooperating agency pursuant to NEPA (40 CFR § 1508.5). There are no other cooperating federal, state, or local entities or Tribes.

NEPA authorizes a federal agency to adopt another agency's NEPA analysis provided that the analysis meets the standards for an adequate statement under the NEPA regulations (40 CFR § 1506.3). Further, a federal agency participating in the NEPA process as a cooperating agency may adopt the NEPA analysis of a lead agency without recirculating the statement when, after an independent review of the statement, the cooperating agency concludes that its comments and suggestions have been satisfied. NOAA, USDOI, and USEPA are participating in the development of the RP/EA as cooperating federal agencies for purposes of NEPA. Upon completion of the Final RP/EA, each agency intends to independently determine if the EA component of the RP/EA is sufficient for the purposes of informing that agency's decision and hence adopt the EA in accordance with 40 CFR § 1506.3 and its agency-specific NEPA procedures. Adoption of the EA would be completed via signature on the final NEPA decision document.

1.5 Purpose and Need

To meet the purpose of restoring losses to those natural resources and services injured as a result of the *DWH* oil spill, the MS TIG proposes to implement restoration projects evaluated in this Draft RP/EA. This Draft RP/EA is consistent with the PDARP/PEIS, which identifies extensive and complex injuries to natural resources and services across the Gulf of Mexico, as well as a need and plan for comprehensive restoration consistent with OPA. This Draft RP/EA focuses on the restoration of injuries to natural resources and services in the Mississippi Restoration Area, using funds made available in the *DWH* Consent Decree.

Section 5.3 of the PDARP/PEIS describes five Programmatic Trustee Goals for restoration work to benefit injured resources and services. The Programmatic Goals that would be addressed by the reasonable range of alternatives proposed in this Draft RP/EA are 1) Restore and Conserve Habitat, 2) Restore Water Quality, 3) Replenish and Protect Living Coastal and Marine Resources, and 4) Monitoring, Adaptive Management, Administrative Oversight. To help meet these goals, in this Draft RP/EA the MS TIG addresses three Restoration Types: WCNH; NR; and Birds.

Consistent with the Programmatic Trustee Goals, the Trustees also developed specific goals to guide

restoration planning and project selection for each Restoration Type⁸ (PDARP/PEIS Sections 5.5.2 through 5.5.14). The specific goals of each Restoration Type selected by the MS TIG for focus in this Draft RP/EA are also described in Section 2.4 of this Draft RP/EA.

Additional information about the Purpose and Need for *DWH* NRDA restoration can be found in Section 5.3.2 of the PDARP/PEIS at page 5-11.

1.6 Proposed Action: MS TIG Draft 2016-2017 RP/EA

This Draft RP/EA addresses the *DWH* restoration goals discussed above by evaluating six action alternatives and proposing to select three of the restoration alternatives (preferred) for implementation. These proposed alternatives are intended to contribute to primary and compensatory restoration of habitats, species, and services in Mississippi using funds made available in the *DWH* Consent Decree. After evaluating the reasonable range of alternatives, the MS TIG proposes the following preferred restoration alternatives (projects) in this Draft RP/EA (Table 1.6-1). The locations for the preferred restoration alternatives (projects) are shown in Figure 1.6-1.

Table 1.6-1: Proposed Preferred Restoration Alternatives (Projects) in this Draft RP/EA

Project/Program	PDARP/PEIS Restoration Goal and Restoration Type	Proposed Funding
Graveline Bay Land Acquisition and Management	Restore and Conserve Habitat: Wetlands, Coastal and Nearshore Habitats Replenish and Protect Living Coastal and Marine Resources: Birds	\$11,500,000
Grand Bay Land Acquisition and Habitat Management	Restore and Conserve Habitat: Wetlands, Coastal and Nearshore Habitats Replenish and Protect Living Coastal and Marine Resources: Birds	\$6,000,000
Upper Pascagoula River Water Quality Enhancement	Restore Water Quality: NR	\$4,000,000

19

⁸ PDARP/PEIS Section 5.5.2.1 describes the goals for Restoration Type Wetlands, Coastal and Nearshore Habitats, Section 5.5.4.1 describes the goals for Restoration Type Nutrient Reduction (Nonpoint Source), and Section 5.5.12.1 presents goals for the Restoration Type Birds.




Figure 1.6-1: Proposed Preferred Alternatives (Projects) for this Draft RP/EA

In order to identify the reasonable range of alternatives for this Draft RP/EA, the MS TIG reviewed PDARP Trustee Restoration Goals, and developed specific MS TIG 2016-2017 Goals and Objectives. The MS TIG identified three Restoration Types from the PDARP/PEIS - WCNH, Birds, and NR (Nonpoint Source) that the TIG considered as appropriate for focus in this Draft RP/EA. The MS TIG then screened project submittals against OPA appropriateness criteria identified in the PDARP. Further detail on the screening process can be found in Section 2.4 of this Draft RP/EA.

The MS TIG will propose evaluating, developing a reasonable range of alternatives, and implementing additional projects in the Mississippi Restoration Area in subsequent restoration plans. Future projects may potentially include projects or project components included in the reasonable range of alternatives in this Draft RP/EA as well as future restoration plans. In addition, additional NEPA analysis will be performed on future projects.

1.7 Public Involvement

Public input is an integral part of NEPA, OPA, and the *DWH* oil spill restoration planning effort. The purpose of public review is to facilitate public discussion regarding the preferred restoration projects, allow the Trustees to solicit and consider public comment, and ensure that final plans consider

relevant issues. The Trustees conducted an extensive public outreach process as part of the PDARP/PEIS; that process is described more fully in Chapter 8 of the PDARP/PEIS. More discussion on public outreach and involvement can also be found in previous phases of *DWH* NRDA Early Restoration Plans available at http://www.gulfspillrestoration.noaa.gov/restoration/early-restoration.

1.7.1 Public Involvement in the Development of this Draft RP/EA and Next Steps

As discussed above in Section 1.3.4, the MS TIG published a Notice of Initiation for Restoration Planning in response to the *DWH* oil spill on May 27, 2016 (hereafter, May 27 2016 Notice).

In developing this Draft RP/EA, the MS TIG considered projects submitted by the public via the MDEQ Restoration Project Idea portal⁹ and the Trustee Project Submission Portal¹⁰ and those proposed in response to the May 27 2016 Notice. The MS TIG received comments and project proposals in response to the May 27 2016 Notice, all of which were considered in the development of this Draft RP/EA. On October 31, 2016, the MS TIG published a Notice of Initiation for Restoration Plan Drafting in Mississippi.¹¹

The public is encouraged to review and comment on this Draft RP/EA. The Draft RP/EA will be made available for public review and comment for forty-five (45) days following the release of the Draft RP/EA, as specified in the public notice published in the Federal Register, the restore.ms website, and the Trustee Council website. Comments on the Draft RP/EA can be submitted during the comment period by one of following methods:

- Via the internet: http://www.gulfspillrestoration.noaa.gov/restoration-areas/mississippi.
- Via hard copy, write:

Mississippi Department of Environmental Quality Attn: Tabatha Baum P.O. Box 2261 Jackson, MS 39225

Submissions must be postmarked no later than 45 days after the release date of the Draft RP/EA.

After the close of the public comment period, the MS TIG will consider the comments received and revise the Draft RP/EA as needed. A summary of comments received and the MS TIG's responses (where applicable) will be included in the Final RP/EA.

1.7.2 Administrative Record

Pursuant to 15 C.F.R. § 990.45, the Trustees opened a publicly available Administrative Record for the *DWH* oil spill NRDA, including restoration planning activities, concurrently with the publication

⁹ http://www.restore.ms/submit-project-idea/

¹⁰ http://www.gulfspillrestoration.noaa.gov/restoration/give-us-your-ideas/suggest-a-restoration-project/

¹¹ http://www.restore.ms/ms-tig-plan/

of the 2010 Notice of Intent to Conduct Restoration Planning (75 Fed. Reg. 60800). DOI is the lead federal Trustee for maintaining the Administrative Record, which can be found at http://www.doi.gov/deepwaterhorizon/adminrecord. Information about the MS TIG restoration project implementation is being provided to the public through the Administrative Record and other outreach efforts, including http://www.gulfspllrestoration.noaa.gov.

1.8 Severability of Projects

In this Draft RP/EA, the MS TIG proposes three preferred restoration alternatives with proposed funding of \$21,500,000. The proposed preferred alternatives presented in this Draft RP/EA are independent of each other and may be selected independently for implementation in this and/or future restoration plans by the MS TIG.

1.9 Decisions to be Made

This Draft RP/EA is intended to provide the public with information and analysis needed to enable meaningful review and comment on the MS TIG's proposal to proceed with selecting up to three restoration projects (preferred alternatives) using *DWH* NRDA funds. Ultimately, this Draft RP/EA and the corresponding opportunity for the public to review and comment on the document are intended to guide the MS TIG's selection of individual restoration projects. Projects not identified for inclusion in the Draft RP/EA may continue to be considered for inclusion in future restoration plans.

1.10 Document Organization

This Draft RP/EA is divided into the following sections:

- Section 1 (Introduction): Introductory information and context for this document;
- Section 2 (Restoration Planning Process): Background on the NRDA restoration planning process, summary of injuries to resources resulting from the *DWH* oil spill that the MS TIG intends to address in this Draft RP, and screening of a suite of restoration projects to address those injuries, and development of the reasonable range of alternatives;
- Section 3 (OPA Evaluation of Alternatives and NEPA Affected Environment and Environmental Consequences): Evaluation of projects proposed for NRDA restoration, proposal of a suite of preferred restoration alternatives, and discussion of NEPA compliance;
- Section 4 (Compliance with Other Laws and Regulations): Discussion of additional federal and state laws that may apply to the proposed preferred alternatives;
- Section 5 (Monitoring and Adaptive Management): Discussion of monitoring and adaptive management requirements for *DWH* oil spill NRDA restoration projects; and
- Section 6 (List of Preparers and Reviewers): Identification of individuals who substantively contributed to the development of this document.

2.0 Restoration Planning Process

NRDA restoration under OPA is a process that includes evaluating injuries to natural resources and natural resource services to determine the types and extent of restoration needed to address the injuries. This Draft RP/EA is consistent with and tiers from the PDARP/PEIS, a programmatic document developed by the Trustees to provide high-level guidance for identifying, evaluating, and selecting future *DWH* restoration projects. Under OPA, the NRDA regulations (15 C.F.R. § 990.54) provide criteria to be used by Trustees to evaluate projects designed to compensate the public for injuries caused by oil spills. To meet the NRDA regulations, the Trustees must identify a reasonable range of restoration alternatives and then evaluate those proposed alternatives. The MS TIG utilized the MGCRP, numerous other regional restoration and ecosystem management planning documents, as well as the MDEQ Restoration Project Idea portal ¹² and the Trustee Project Submission Portal ¹³ during development of this Draft RP/EA. This section of the Draft RP/EA summarizes the restoration planning process for the MS TIG including planning objectives of the MGCRP, the project screening process developed by the MS TIG, and a discussion of the reasonable range of alternatives.

2.1 Injuries Addressed in this Draft RP/EA

Chapter 4 of the PDARP/PEIS summarizes the injury assessment which established the nature, degree, and extent of injuries from the *DWH* incident to both natural resources and the services they provide. Restoration projects proposed in this Draft RP/EA and in future MS TIG restoration plans are designed to address injuries in the Mississippi Restoration Area resulting from the *DWH* oil spill. As discussed in Section 1, the MS TIG will focus on the following Restoration Types which are described in the PDARP/PEIS: WCNH/Birds and NR.

2.1.1 Wetlands, Coastal, and Nearshore Habitats

The *DWH* oil spill caused significant injuries to Mississippi's nearshore marine ecosystem, including interrelated and biologically diverse habitats such as estuarine coastal wetland complexes, beaches and dunes, barrier islands, submerged aquatic vegetation (SAV), oyster reefs, and shallow unvegetated areas (see PDARP/PEIS Section 4.6.1.1 Ecological Description, p. 4-292). Injuries were detected over a range of species, communities and habitats, affecting a wide variety of ecosystem components (PDARP/PEIS Section 4.6.9). The Trustees allocated the greatest amount of funding to the Restore and Conserve Habitat goal, because of the critical role that coastal and nearshore habitats play in the overall productivity of the Gulf of Mexico. The MS TIG recognizes that one of three general restoration program areas of the MGCRP is Land Resources, which focuses on the need to conserve and manage priority lands, including lands already under protection. Objectives outlined in the program include conserving priority habitats by utilizing land protection tools such as fee title acquisition, conservation easements, and land donations; as well as managing and restoring priority habitats through actions including management plans, invasive species management, the use of prescribed fire (where appropriate), and enhancement of riparian zone buffers.

¹² http://www.restore.ms/submit-project-idea/

¹³ http://www.gulfspillrestoration.noaa.gov/restoration/give-us-your-ideas/suggest-a-restoration-project/

2.1.2 Nutrient Reduction (Nonpoint Source)

Excessive nutrient enrichment, or eutrophication, of Gulf Coast estuaries and their watersheds is a chronic threat that can lead to hypoxia (low oxygen levels), harmful algal blooms, habitat losses, and fish kills (PDARP/PEIS Section 5.5.4). This restoration type ties directly into the Water Resources Program vision described in the MGCRP, which is to restore and enhance the ecological and hydrological integrity of our water resources, including improved water quality and ensuring natural water quantity levels to our coastal bays and estuaries and coastal rivers and streams. One of the defined objectives of this vision is reduction of rural nonpoint source pollution by implementing and improving agricultural, forestry and watershed best management practices. Examples of restoration actions include reducing erosion and thus sedimentation into coastal streams and managing excess nutrient levels to coastal basins.

2.1.3 Birds

As discussed in the PDARP/PEIS (Section 5.5.12), exposure to oil injured a large number of bird species occupying different habitats, from offshore to nearshore, including open water, beaches, island waterbird colonies, bays and coastal marshes (PDARP/PEIS Section 4.7). More than 150 species of birds occur in waters and wetlands of the northern Gulf of Mexico for at least a portion of their lives, and nearly 300 species use either the coast itself or coastal upland habitat directly adjacent to the Gulf. The *DWH* oil spill affected all nearshore habitats along the northern Gulf, including those in the Mississippi Restoration Area. Given the extensive injuries to birds and their various habitats in Mississippi, the MS TIG decided to focus on development of a reasonable range of alternatives for projects that would help restore bird injuries. This Restoration Type is consistent with the MGCRP's Coastal and Marine Living Resources Program, which is intended to restore and stabilize populations of ecologically and commercially and or recreationally important coastal and marine species at sustainable levels. One of the program objectives is sustainable management and enhancement of coastal and marine living resource populations, through restoration actions such as reducing bird nest predation and human disturbance.

2.2 DWH Early Restoration Addressing the Injuries to Date

During *DWH* NRDA Early Restoration, the Trustees selected the following two projects for implementation in the Mississippi Restoration Area that are included in the "Wetlands, Coastal, and Nearshore Habitats" (WCNH) Restoration Type.

Hancock County Marsh Living Shoreline

This project is intended to employ living shoreline techniques, including natural and artificial breakwater material and marsh creation, to reduce shoreline erosion by dampening wave energy while encouraging reestablishment of habitat that once was present in the region. The project will provide for construction of up to 5.9 miles of living breakwater, approximately 46 acres of marsh creation, and approximately 46 acres of subtidal reef restoration in Heron Bay to increase secondary productivity in the area. The project will reduce shoreline erosion, create habitat for secondary

productivity, and protect and create salt marsh habitat. More details on this project can be found in the Phase III Early Restoration Plan. 14

Restoring Living Shorelines and Reefs in Mississippi Estuaries

The project will restore secondary productivity through the placement of intertidal and subtidal reefs and the use of living shoreline techniques including breakwaters. The project will be implemented at locations in Grand Bay, Graveline Bay, Back Bay of Biloxi and vicinity, and St. Louis Bay in Jackson, Harrison, and Hancock Counties, Mississippi. The project builds on recent collaborative projects implemented by the Mississippi Department of Marine Resources (MDMR), NOAA, and The Nature Conservancy. The project will construct over four miles of breakwaters, five acres of intertidal reef habitat, and 267 acres of subtidal reef habitat. Over time, the breakwaters, intertidal and subtidal restoration areas will develop into living reefs that support benthic secondary productivity, including, but not limited to, oysters/bivalve mollusks, annelid worms, shrimp, and crabs. Breakwaters will reduce shoreline erosion as well as marsh loss. More details on this project can be found in the Phase IV Early Restoration Plan. ¹⁵

One Early Restoration project was selected for implementation in the Mississippi Restoration Area that would be included in the "Birds" Restoration Type, as described below.

Enhanced Management of Avian Breeding Habitat Injuries by Response in the Florida Panhandle, Alabama and Mississippi.

The Enhanced Management of Avian Breeding Habitat Injured by Response Activities in the Florida Panhandle, Alabama, and Mississippi project will reduce disturbance to nesting and foraging habitat for beach-nesting birds in the project areas. The project involves three tasks: (1) Placing symbolic fencing (signs and posts connected with rope) around sensitive nesting sites of beach-nesting birds to indicate the site as off-limits to people, pets, and other sources of disturbance; (2) Increasing predator control to reduce disturbance and loss of eggs, chicks, and adult beach-nesting birds at nesting sites; and (3) Increasing surveillance and monitoring of posted nesting sites to assess disturbance to nesting habitat in posted areas. In Mississippi, the project would be implemented on Gulf Island National Seashore (GUIS) - Mississippi District. More details on this project can be found in the Phase II Early Restoration Plan. ¹⁶

No Early Restoration projects have been selected in the Mississippi Restoration Area that would be included in the NR Restoration Type.

More information on the status of all *DWH* NRDA Early Restoration projects and a summary of funds obligated and expended on each project can be found on NOAA's Gulf Spill Restoration Early Restoration Project Atlas. ¹⁷

25

¹⁴ http://www.gulfspillrestoration.noaa.gov/restoration/early-restoration/phase-iii

¹⁵ http://www.gulfspillrestoration.noaa.gov/restoration/early-restoration/phase-iv

¹⁶ http://www.gulfspillrestoration.noaa.gov/sites/default/files/wp-content/uploads/Phase-II-ERP-ER-12-21-12.pdf

¹⁷ http://www.gulfspillrestoration.noaa.gov/restoration/early-restoration/early-restoration-projects-atlas

2.3 Coordination with Other Gulf Restoration Programs

As discussed in the PDARP/PEIS Chapter 1.5.6, the Trustees are committed to coordination with other Gulf of Mexico restoration programs to maximize the overall ecosystem benefit of *DWH* NRDA restoration efforts. This coordination will ensure that funds are allocated strategically to restoration projects across the affected regions of the Gulf of Mexico and within the Mississippi Restoration Area.

The MS TIG will continue ongoing efforts to coordinate project development and leveraging in cooperation with the other *DWH* funding streams – the NFWF GEBF and the RESTORE Act. To that end, the MGCRP described above was funded by NFWF GEBF and was released by MDEQ in 2015. As discussed, this plan sets forth a process for identification of restoration actions in priority habitat and resource areas that result in ecologically sound and sustainable projects. Its purpose was to "Create a plan that would result in a coordinated, systemic, and transparent process for sustainable ecological restoration in Mississippi, that will direct funds associated with the GEBF, and be applicable to informing ecological restoration funding associated with the RESTORE Act."

The Graveline Bay Land Acquisition and Management preferred project alternative would leverage NFWF funding for acquisition and management in the proposed project area.

The Grand Bay Land Acquisition and Habitat Management preferred project alternative would leverage NFWF and RESTORE Act funding already awarded for habitat acquisition and management in the proposed project area.

2.4 Screening for Potential Alternatives

Under the OPA regulations (15 C.F.R. § 990.53), the MS TIG developed a screening process to develop a reasonable range of alternatives to be further evaluated in this plan. The following sections describe the screening process the MS TIG used to identify restoration approaches and a reasonable range of alternatives to include in this Draft RP/EA.

2.4.1 MS TIG Screening Process

The MS TIG reviewed PDARP/PEIS Programmatic Trustee Goals and developed MS TIG 2016-2017 Goals and Objectives for identifying projects to develop a reasonable range of alternatives for restoration in this Draft RP/EA. The MS TIG identified three Restoration Types described in the PDARP/PEIS - WCNH, Birds, and NR that the TIG considered appropriate for this Draft RP/EA.


The project screening process developed by the MS TIG for the purpose of preparing this Draft RP/EA, including ideas submitted by the public via the MDEQ Restoration Project Idea portal ¹⁸ and the Trustee Project Submission Portal ¹⁹, and those submitted in response to the May 27 2016 Notice. The MS TIG queried all projects in both portals identified above and sorted the combined, cumulative

¹⁸ http://www.restore.ms/submit-project-idea/

¹⁹ http://www.gulfspillrestoration.noaa.gov/restoration/give-us-vour-ideas/suggest-a-restoration-project/

project list according to the selected Restoration Types: WCNH, Birds, and NR (Nonpoint Source). Because many portal submissions did not contain sufficient detail or overlapped in scope, the MS TIG further developed restoration project alternatives using components of portal submitted ideas, regional management plans, and resource expertise within the MS TIG (MS TIG Projects²⁰). All projects were evaluated in a similar fashion and against the same criteria.

The MS TIG project screening process is illustrated below. OPA Screening for WCNH/Birds is described in Section 2.4.1.4.1 and OPA screening for NR (Nonpoint Source) is described in Section 2.4.1.4.2. The process is presented in a step-wise manner in Figure 2.4.-1; however, project screening, project development and project selection were iterative processes that were performed collaboratively by the co-Trustees of the MS TIG.



Reasonable Range of Alternatives

Figure 2.4-1: Generalized Process of Identifying the Reasonable Range of Alternatives 2.4.1.1 Consistency with Goals outlined in PDARP/PEIS Restoration Types

The MS TIG considered Restoration Types in developing the reasonable range of alternatives for this Draft RP/EA. The figure below (Figure 2.4-2, taken from the PDARP/PEIS) graphically summarizes the PRDARP/PEIS Restoration Types, the comprehensive restoration plan, the goals and their related restoration type(s) and related restoration approaches. Monitoring, adaptive management, and administrative oversight are planned throughout all restoration types.

²⁰ For the purposes of discussion, MS TIG Projects refers to the pool of projects, proposals, and project ideas that were developed as described and considered in the OPA screening process.




Figure 2.4-2:Trustees' Restoration Goals and Restoration Types (provided as Figure 5.4-1 in the PDARP/PEIS)

Restoration Goals by Restoration Type: Specific restoration goals outlined in the PDARP/PEIS for WCNH, Birds and NR (Nonpoint Source) Restoration Types are described below.

<u>WCNH Restoration Type</u>: Multiple benefits can be derived through restoration of WCNH at a large scale. The specific goals of the WCNH Restoration Type include:

• Restore a variety of interspersed and ecologically connected coastal habitats in each of the five Gulf states to maintain ecosystem diversity, with focus on maximizing ecological

- functions for the range of resources injured by the *DWH* oil spill, such as oysters, estuarine-dependent fish species, birds, marine mammals, and nearshore benthic communities.
- Restore for injuries to habitats in the geographic areas where the injuries occurred, while considering approaches that provide resiliency and sustainability.
- While acknowledging the existing distribution of habitats throughout the Gulf of Mexico, restore habitats in appropriate combinations for any given geographic area. Consider design factors, such as connectivity, size, and distance between projects, to address injuries to the associated living coastal and marine resources and restore the ecological functions provided by those habitats.

<u>Birds Restoration Type</u>: The MS TIG also considered projects that would help restore birds injured by the *DWH* oil spill. Under the Replenish and Protect Living Coastal and Marine Resources Programmatic Goal, the MS TIG will focus on the Birds Restoration Type. Specific restoration goals of the Birds Restoration Type include:

- Restore lost birds by facilitating additional production and/or reduced mortality of injured bird species.
- Restore or protect habitats on which injured birds rely.
- Restore injured birds by species where actions would provide the greatest benefits within geographic ranges that include the Gulf of Mexico.

NR (Nonpoint Source) Restoration Type: The MS TIG recognizes that nutrient pollution adversely impacts water quality and poses a significant threat to localized watersheds across the Gulf Coast. NR measures can benefit the estuaries that are integral habitat for many important species. Under the Restore Water Quality Programmatic Goal, the MS TIG will focus on the NR (Nonpoint Source) Restoration Type, and these specific restoration type goals:

- Reduce nutrient loadings to Gulf Coast estuaries, habitats, and resources that are threatened by chronic eutrophication, hypoxia, or harmful algal blooms (HABs) or that suffer habitat losses associated with water quality degradation.
- Where appropriate, co-locate nutrient load reduction projects with other restoration projects to enhance ecological services provided by other restoration approaches.
- Enhance ecosystem services of existing and restored Gulf Coast habitats.

The MS TIG screened projects for consistency with PDARP/PEIS Restoration Type goals for WCNH, Birds, and NR.

2.4.1.2 MS TIG 2016-2017 Goals and Objectives

After ensuring consistency with the PDARP/PEIS Restoration Type goals for WCNH, Birds, and NR, the MS TIG identified four broad objectives for this Draft RP/EA: regional connectivity; leveraging; project partnering opportunities; and existing regional planning initiatives.

• Regional Connectivity: A key goal in the development of WCNH in the PDARP/PEIS is to restore a variety of interspersed and ecologically connected coastal habitats. In addition, TIGs are encouraged to "Consider design factors, such as connectivity, size, and distance between projects, to address injuries to the associated living coastal and marine resources and restore the ecological functions provided by those habitats." Conservation, management

- and restoration of habitats are also MGCRP Land Program priorities. Preservation, restoration, regional connectivity and proximity to state and federal conservation lands were key factors in determining restoration approaches/techniques and in the screening of projects.
- <u>Leveraging:</u> The MGCRP is Mississippi's plan for *DWH* ecosystem restoration planning in Mississippi. The MS TIG considered opportunities to leverage NFWF GEBF and RESTORE funding in the screening and selection of projects. The MS TIG also considered the extent that NFWF GEBF funds or RESTORE funds have been programmed to accomplish a restoration initiative or projects in the project screening.
- <u>Partnering</u>: The MS TIG considered Trustee expertise from state and federal programs and/or resource management expertise. Opportunities to share resource management expertise and funded programs were considered in the selection of restoration techniques and in the screening of projects.
- <u>Regional Planning Initiatives</u>: Regional plans and planning initiatives were objectives also considered in project screening including the MGCRP, Sand Hill Crane and Grand Bay National Wildlife Refuge planning documents, MDMR management plans, US Army Corps of Engineers (USACE) Mississippi Coastal Improvement Plan (MsCIP), and the MDMR Coastal Preserves (CP) planning initiative²¹.

The MS TIG screened projects for consistency with the MS TIG Goals and Objectives.

2.4.1.3 Identification of PDARP/PEIS Restoration Approaches/Techniques

Based on a review of the PDARP/PEIS Restoration Type goals, MS TIG 2016-2017 Goals and Objectives, and the MGCRP program priorities, the MS TIG identified restoration approaches associated with the following Restoration Types WCNH, Bird, and NR. Table 2.4.-1 demonstrates how the MS TIG's preferred restoration approaches/techniques for WCNH, Birds and NR Restoration Types align with MGCRP program objectives. The rationale for selecting restoration approach/techniques and the decisions made for project screening is outlined below.

_

²¹ http://www.dmr.ms.gov/index.php/mississippi-gems

Table 2.4-1: PDARP/PEIS Restoration Approaches/ MGCRP priorities and MS TIG Goals and Objectives

	Miss Res	issippi Co toration P	astal lan	MS T	TIG Goa	als and Objectives	
PDARP Restoration Types/Approaches/Techniques	MS Land Resources Program Priority	MS Coastal and Marine Resources Priority	MS Water Resources Program Priority	Regional Connectivity	Leveraging	Partnering/Multiple Trustee engagement	Existing Regional Planning Initiatives
Wetland Coastal And Nearshore Habitats							
Acquire lands for conservation	X		X	х	X	X	X
Develop and implement management actions in conservation areas and/or restoration projects	х		X	x	x	х	X
Nutrient Reduction (Nonpoint Source)-(Also includes Protect and Cons	serve Marine (Coastal, Es	tuarine, and	Riparia	n habita	ts)	
Nutrient Reduction (Nonpoint Source)-(Also includes Protect and Cons Reduce Nutrient Loads to Coastal Watersheds	serve Marine (Coastal, Es	tuarine, and	Riparia	n habita	ts)	X
	serve Marine (Coastal, Es		Riparia			x x
Reduce Nutrient Loads to Coastal Watersheds	serve Marine (Coastal, Es	x	Riparia	X	x	
Reduce Nutrient Loads to Coastal Watersheds Agricultural conservation practices	serve Marine (Coastal, Es	x x	Riparia	x x	x x	x
Reduce Nutrient Loads to Coastal Watersheds Agricultural conservation practices Forestry management practices			x x x	Riparia	X X X	x x	x
Reduce Nutrient Loads to Coastal Watersheds Agricultural conservation practices Forestry management practices Implement erosion and sediment control (ESC) practices			x x x	Riparia	X X X	x x	x
Reduce Nutrient Loads to Coastal Watersheds Agricultural conservation practices Forestry management practices Implement erosion and sediment control (ESC) practices Birds (Also includes Protect and Conserve Marine Coastal, Estuarine, a			x x x	Riparia	X X X	x x x	x

Rationale for Protect and Conserve Marine, Coastal, Estuarine and Riparian Habitats as a restoration approach: WCNH is a broad Restoration Type which could include restoration techniques such as beneficial use, land acquisition, habitat enhancement/restoration on public and private lands, backfilling canals and numerous other techniques. The restoration goals of the WCNH Restoration Type include the consideration of connectivity, size and proximity to other land conservation projects. OPA screening was focused on land acquisition for the purpose of protection, conservation and restoration/management of coastal marine and riparian habitats so as to achieve multiple benefits,

including birds, which fall under the Bird Restoration Type ²². The Bird Restoration Type also includes Protect and Conserve Marine, Coastal, Estuarine, and Riparian habitats, enhancing habitat through vegetation management and nesting and foraging area stewardship. For these reasons, the MS TIG completed a screening process where the WCNH Restoration Type and the Bird Restoration Type were combined and evaluated together (See Section 2.4.1).

Rationale for screening based on agricultural and forestry management practices restoration techniques: The NR (Nonpoint Source) Restoration Type also includes Protect and Conserve Marine, Coastal, Estuarine, and Riparian Habitats as a restoration approach, as well as agricultural conservation practices and forestry management practices. For the purposes of the Draft RP/EA, screening focuses on agricultural and forestry management practices as restoration techniques.

The MS TIG selected the following restoration approaches and techniques:

WCNH

Approach: Protect and Conserve Marine, Coastal, Estuarine, and Riparian Habitats *Techniques*: Acquire lands for conservation

Develop and implement management actions in conservation areas and/or restoration projects

Birds

Approach: Restore and Conserve Bird Nesting and Foraging Habitat *Techniques*: Enhance habitat through vegetation management Nesting and foraging area stewardship

• NR (Nonpoint Source)

Approach: Reduce Nutrient Loads to Coastal Watersheds

Techniques: Agricultural conservation practices

Forestry management practices

2.4.1.4 OPA Screening

2.4.1.4.1 WCNH/Birds

For WCNH/Birds, MS TIG projects were screened against OPA appropriateness criteria described in the PDARP/PEIS, consistency with MS TIG 2016-2017 Goals and Objectives and some additional considerations needed to identify the reasonable range of alternatives.

PDARP/PEIS OPA Appropriateness Criteria: The MS TIG considered two approaches under the WCNH and Birds Restoration Types. The approaches and their OPA appropriateness criteria are discussed below:

Protect and Conserve Marine Coastal, Estuarine and Riparian Habitats restoration approach (Appendix 5 D; D.1.7.2; page 5-239). This restoration approach can help return injured natural resources and services to baseline conditions by minimizing or eliminating the potential for future loss or degradation of protected lands and or enhancing the ecosystem services provided by protected lands over time when compared to the future of those protected areas without the conservation action.

²² Birds also includes Land Acquisition-Protect and Conserve Marine, Coastal, Estuarine, and Riparian habitats.

It can also help to compensate for interim service losses to 1) coastal and riparian buffer uplands; 2) coastal wetland, oyster, submerged aquatic vegetation (SAV) or beach/barrier island habitats; and 3) nearshore and offshore living coastal and marine resources (fish, shellfish, birds, sea turtles, and marine mammals) that were adversely affected by the *DWH* oil spill. These techniques have been well demonstrated and this approach is highly likely to succeed. Additionally, collateral injury to other natural resources is expected to be minimal or avoided entirely. The MS TIG does not anticipate that the approach will negatively affect public health or safety and consider the approach likely to benefit other natural resources.

The Restore and Conserve Bird Nesting and Foraging Habitat restoration approach (Appendix 5 D; D.6.1.2; page 5-307) can restore injured natural resources and services to baseline conditions by supporting increased health and reproduction of birds. This approach may also help compensate for interim services losses to birds adversely affected by the *DWH* oil spill through restoring, rehabilitating, and/or replacing habitats that provide services to injured bird species. These are established techniques to provide services to birds. Collateral injury to other natural resources is expected to be minimal and short-term, however, project selection and design considered potential impacts on existing habitat. The project approach is not expected to negatively affect public health or safety. Additionally, the MS TIG considers it likely that it will also benefit additional natural resources.

MS TIG projects were screened against combined OPA appropriateness criteria. Consistent with the goals and restoration approaches discussed in Section 2.4.1.3 above, the MS TIG considered projects that would help compensate for interim services losses to injured birds, and selected projects for further consideration that provided bird nest protection/bird production, habitat acquisition, and habitat restoration. Projects that were not consistent with the goals and restoration approaches discussed above and were eliminated from further consideration in this plan included monitoring, wildlife rehabilitation, recreational loss, stormwater management, nutrient reduction, and prairie restoration projects, as well as those already funded by another source or multiple redundant entries of the same project.

Consistency with MS TIG 2016 - 2017 Goals and Objectives: The MS TIG considered projects that would provide regional connectivity, leveraging opportunities, multiple trustee engagement, and were consistent with the MGCRP and with the PDARP/PEIS. Projects considered further included large acquisitions, habitat restoration, and projects that could be leveraged with funds outside of the NRDA process, such as RESTORE or NFWF funds. Projects that were eliminated included projects that provided only limited regional connectivity.

Additional considerations allowed under OPA: The MS TIG also screened projects for compliance with the following additional considerations allowed under OPA, which were developed by the MS TIG for the 2016-2017 funding cycle.

- Project is consistent with regional planning efforts or ongoing restoration efforts including National Wildlife Refuge (NWR) management plans, the MS CP program and others;
- Project has willing seller(s); and
- Project has management opportunity on adjacent lands.

The MS TIG further considered projects that were included in regional planning efforts and projects with willing sellers, and involved large-scale acquisition and habitat restoration. Projects that were

not further considered included those for which sufficient detail was lacking and where project components were included in other programs/projects (CP Tracts).

Selection of Projects for Development of the Reasonable Range of Alternatives for WCNH/Birds: The MS TIG selected two projects for the development of the reasonable range of alternatives, in addition to the No Action Alternative: 1) Graveline Bay Land Acquisition and Management proposed alternative; and 2) Grand Bay Land Acquisition and Habitat Management proposed alternatives. The development of the reasonable range of alternatives is discussed in Section 2.6.

2.4.1.4.2 NR (Nonpoint Source)

For NR (Nonpoint Source), MS TIG projects were screened against OPA appropriateness criteria described in the PDARP/PEIS, consistency with MS TIG 2016 - 2017 Goals and Objectives and additional considerations used to identify a final suite of project ideas. These project ideas were then employed to develop the reasonable range of alternatives described in Section 2.6.

PDARP/PEIS OPA Appropriateness Evaluation: Appendix 5D of the PDARP/PEIS describes four individual restoration approaches that could be used to implement the NR (Nonpoint Source) Restoration Type. Outlined below is the OPA appropriateness evaluation for the restoration approach that was considered by the MS TIG under the NR (Nonpoint Source) Restoration Type.

The Reduce Nutrient Loads to Coastal Watersheds restoration approach (D.2.1.2 page 5-242): This approach enhances ecosystem services provided by restored habitats and resources and may return injured natural resources and services to baseline conditions by reducing nutrient loads to coastal watersheds, improving water quality, reducing the extent of eutrophication and occurrence of hypoxia and/or HABs, reducing turbidity, and increasing light penetration. This approach can help compensate for interim services losses to estuarine-dependent water column resources, oysters, SAV, and recreational uses adversely affected by the *DWH* oil spill. It also compensates for lost ecosystem services by reducing nutrient runoff, which will improve water quality and mitigate chronic ecosystem threats and impaired recreational use to provide ecosystem benefits to injured resources and habitats. This project approach has demonstrated effectiveness as shown in numerous studies by the USDA's Conservation Effects Assessment Program (CEAP) and water quality restoration "Success Stories" for the EPA Section 319 Nonpoint Source Management Program grants. The risk of collateral injury to other natural resources is expected to be minimal. Collateral injury could occur during project construction, but these effects can be minimized during the design process. The MS TIG does not anticipate that the project will negatively affect public health or safety and considers it likely to benefit additional natural resources.

MS TIG projects were screened against the OPA appropriateness criteria. The MS TIG considered nutrient reduction projects and those that would compensate for interim or lost services, and retained nutrient reduction, water quality, and sediment reduction projects for further consideration. Projects that were eliminated from consideration included multiple redundant entries of the same project, monitoring projects, infrastructure projects, and water quality projects such as dredging, debris removal, and drainage improvements.

Consistency with MS TIG 2016 - 2017 Goals and Objectives: The MS TIG screened projects for consistency with the goals developed for the Draft RP/EA, outlined above. The MS TIG considered

agricultural conservation practices and forestry management practices that would provide nutrient and sediment reduction, projects that would leverage other funding opportunities, those with the potential for multiple trustee engagement, and those that were consistent with the MGCRP and with the PDARP/PEIS. As a result, nutrient and sediment reduction projects were considered further. Projects that were eliminated included multiple redundant entries of the same project or projects that provided similar benefits, or projects where nutrient and sediment reduction was only a minor component.

Additional considerations allowed under OPA: The MS TIG also screened projects for compliance with the following additional considerations which were developed by the MS TIG for the Draft RP/EA.

- Conservation practices on agricultural lands in cooperation with landowners, and that could be leveraged by existing Mississippi TIG Trustee programs, such as the USDA-NRCS Farm Bill and other programs;
- Projects that would reduce nutrient and sediment load contribution in the Pascagoula River watershed, which contains Gulf sturgeon Critical Habitat; and
- Selection of the appropriate sub-watershed in which conservation practices applied to land uses would maximize water quality benefits in the Mississippi Sound, particularly sediment removal.

The MS TIG considered projects that would result in nutrient and sediment reduction, that would involve USDA program participation, and for which the MS TIG had demonstrated experience in the geographic area. The MS TIG kept projects that included the implementation of agricultural conservation practices that would reduce nutrient runoff from the landscape; reduce nutrient loads to streams and downstream receiving waters; and could provide benefits to marine resources and benefits to coastal watersheds. Projects that were eliminated included best management practices (BMPs) that do not provide a benefit to agricultural and forested lands.

Selection of Projects for Development of the Reasonable Range of Alternatives for NR: The MS TIG selected two projects for the development of the reasonable range of alternatives, in addition to the No Action Alternative; 1) Upper Pascagoula River Water Quality Enhancement proposed alternative and 2) Pascagoula River Basin Riparian Buffer Maintenance Plan proposed alternative. The development of the reasonable range of alternatives is discussed in Section 2.6.

2.5 Alternatives not Considered for Further Evaluation in this Draft RP/EA

The MS TIG OPA screening described in this Section resulted in the reasonable range of alternatives. Following the screening steps outlined above, there were a number of land acquisition and management project submittals which included acquisition and management of larger acreage that would have provided benefits to WCNH and Birds. These projects collectively include acquisition to expand the Mississippi CP sites and related management activities (discussed below). These individual projects were considered through the screening process and it was determined that they met the screening criteria, and with further development could be selected in a future restoration plan. However, these projects, although valuable in terms of WCNH/Bird benefits, are not further considered for evaluation in this plan. A review of the collective Mississippi C P project is

summarized below. See Section 2.4 for a discussion of screening of projects and reasons for eliminating various projects that were not considered for development as the reasonable range of alternatives in this Draft RP/EA.

Review of Mississippi CP Project: This review combines project submittals proposing protection of ecologically significant parcels from willing sellers in the three coastal counties. The parcels would be located within or adjacent to CP boundaries, and would then be preserved and managed by the MDMR Mississippi CP system. Currently, the Mississippi CP system manages over 36,000 acres of coastal lands. Project proposals include up to 183,000 acres of acquisition that were not selected for evaluation in this Draft RP/EA, the larger projects occurring in the following areas:

- Biloxi River Marsh CP Acquisition
- Escatawpa River Marsh CP Acquisition
- Pascagoula River Marsh CP Acquisition
- Wolf River Restoration Project
- Bellefontaine Marsh Preserve Land Protection
- Old Fort Bayou Land Protection
- Tchoutacabouffa Land Protection
- Delisle Bayou Land Protection
- Ansley Area Land Protection Land Acquisition- Jourdan River CP

Although this large-scale land acquisition program would protect bird habitat, including slash pine forest, estuarine and intertidal wetlands, and beaches, the project components are geographically disparate, and beyond the financial scope of the current funds available to the MS TIG. The MS TIG coordinated with MDMR to consider their acquisition priorities as well as MS TIG goals and objectives including connectivity and use of existing management plans. The MS TIG developed the large land acquisition and management projects including Graveline Bay CP and the Grand Bay Savanna CP from the list because of the existing planning at Grand Bay NWR and Grand Bay National Estuarine Research Reserve (NERR), and the Grand Bay Savanna CP, as well as the proximity/connectivity benefits that Graveline Bay CP provides considering these and other large conservation areas nearby (Sandhill Crane NWR). Additionally, activities within Grand Bay NWR, Grand Bay NERR, and Graveline Bay CP have been considered in previous restoration plans, NEPA analyses, and state planning efforts. Components of proposed projects not selected for development of the reasonable range of alternatives, and thus for analysis in this Draft RP/EA, may be considered in future MS TIG restoration plans.

2.6 Reasonable Range of Restoration Alternatives Considered

The development of the reasonable range of alternatives for the OPA selected projects for WCNH/Birds and NR Restoration Types is discussed here.

2.6.1 Restoration Type: WCNH/Birds

WCNH/Bird screening resulted in identification of two projects for development as the reasonable range of alternatives: Graveline Bay Land Acquisition and Management and Grand Bay Land

Acquisition and Habitat Management. There is one proposed alternative for the proposed Graveline Bay project area, three alternative means of accomplishing the goals in the proposed Grand Bay project area, and the No Action alternative. The WCNH/Birds alternatives for the Draft RP/EA are listed here:

Alternative A (Preferred): Graveline Bay Land Acquisition and Management

Alternative B: Grand Bay Land Acquisition (up to 8,000 acres)

Alternative C: Grand Bay Habitat Management (up to 17,500 acres)

Alternative D (Preferred): Grand Bay Land Acquisition (up to 8,000 acres) and Habitat

Management (up to 17,500 acres); Alternatives B and C combined

Natural Recovery/No Action (No Action)

Management Plans and Planning Initiatives Considered in the Development of the Reasonable Range of Alternatives

The MS TIG considered existing management plans and planning initiatives in the identification of projects and in the development of the reasonable range of alternatives.

Graveline Bay CPs: The MDMR CP Program includes tracts throughout the coastal counties in Mississippi. The Graveline Bay CP was one of the initial acquisitions by the State of Mississippi dedicated as a CP. Existing monitoring activities by the state and cooperative partners include marsh bird monitoring, routine salinity monitoring and shellfish surveys. Much of the property is considered tidal wetlands owned by the State. The State will manage the area as a CP for conservation purposes to protect ecological integrity of tidal marsh and adjacent uplands. Threats to the ecological integrity of the Graveline Bay CP include the potential for future development of habitat adjacent to the marsh, septic tank contamination from adjacent development, and limited flushing action of bay. Graveline Bay CP priorities include acquisition of properties within and adjacent to CP boundaries and habitat management of the same.

<u>Grand Bay NWR, NERR and Grand Bay Savanna CP</u>: There are currently several management documents used by natural resource agencies managing habitats within the project boundary. These documents would be used as guidance to select and prescribe the appropriate restoration activities and management measures on a parcel by parcel basis. A summary of each of these documents is provided below:

Grand Bay National Estuarine Research Reserve Final Environmental Impact Statement/Reserve Management Plan²³: This EISwas finalized in 1998 by the MDMR. The purpose of this plan was to designate the area as part of the NERR. For designation, a reserve management plan was produced and in 2013 was updated. The Grand Bay NERR Management Plan 2013-2018 frames out stewardship, resource protection, public use/access, research and monitoring, education and coastal training plans.

 $[\]frac{23}{http://grandbaynerr.org/wp-content/uploads/2010/08/Grand-Bay-NERR-Final-Environmental-Impact-Statement-Reserve-Management-Plan.pdf}$

<u>Grand Bay National Wildlife Refuge Comprehensive Conservation Plan²⁴:</u> This plan was finalized in 2008 by USFWS. The purpose of the plan was to guide management actions and direction over a 15-year period. Specifically, the Comprehensive Conservation Plan (CCP) was written to:

- Provide a clear statement of the refuge's management direction;
- Provide refuge neighbors, visitors, and government officials with an understanding of the USFWS's management actions on and around the refuge;
- Ensure that the USFWS's management actions, including land protection and recreation/education programs, are consistent with the mandates of the NWR System; and
- Provide a basis for development of the refuge's budget requests for operations, maintenance, and capital improvement needs.

<u>Land Protection Plan and Final Environmental Assessment for the Expansion of Grand Bay National Wildlife Refuge²⁵</u>: This plan was finalized in 2012 by USFWS. This plan identified the proposed acquisition boundary for the proposed expansion of NWR. It delineated approximately 8,428 acres from four areas adjacent to the refuge for acquisition, restoration, enhancement, and management. The purpose of the proposed refuge expansion was to conserve valuable riverine habitat, to protect threatened and endangered species, to restore and protect key habitats (i.e., coastal savanna and longleaf pine), and to manage populations of migratory birds and other interjurisdictional trust species.

2.6.1.1 Alternative A (Preferred): Graveline Bay Land Acquisition and Management

The proposed Graveline Bay Land Acquisition and Management alternative includes acquiring and managing parcels within the existing Graveline Bay CP and nearby publicly owned lands. Development is a threat to habitats adjacent to the preserve. To the north, residential developments are adjacent to developable uplands that currently buffer the Graveline marsh. Municipal land use plans reflect the project area for the proposed alternative is zoned as new growth, which would allow residential and commercial development. Without protection, the MS TIG anticipates that future residential development would continue in these areas. The proposed alternative would be implemented at locations in Graveline Bay in Jackson County, Mississippi. The planning process for Alternative A has been a collaboration between MDMR and the MS TIG. Proposed Alternative A includes the acquisition of land from willing sellers, preservation and habitat enhancement of up to 1,410 acres. Habitat to be acquired includes estuarine marsh, shoreline (beach) and other coastal riparian habitats which provide foraging, loafing and nesting for bird species that were injured in the DWH oil spill. Restoration measures and benefits would include acquisition to reduce the threat of development, direct enhancement of habitat, decreased habitat fragmentation and increased habitat connectivity to other large conservation parcels in the area. Protection of shoreline habitat from vehicle traffic would also enhance shorebird nesting success. Additional details for proposed Alternative A (including restoration measures) are provided in Section 3 of this document.

 $\frac{25}{\text{https://restorethegulf.gov/sites/default/files/documents/images/Grand\%20Bay\%20Final\%20Land\%20Protection\%20Plan\%20and\%20Environmental\%20Assessment.pdf}$

²⁴ https://catalog.data.gov/dataset/grand-bay-national-wildlife-refuge-comprehensive-conservation-plan

2.6.1.2 Alternative B: Grand Bay Land Acquisition (up to 8,000 acres)

The goal of Alternative B is to acquire up to 8,000 acres of estuarine marsh, coastal pine flatwoods, coastal pine savanna, maritime forest, bottomland hardwoods, and coastal marsh within the boundaries of Grand Bay NWR/NERR and Grand Bay Savanna CP. Doing so would help restore injuries to WCNH injured by the *DWH* oil spill as well as habitats on which injured bird species rely. Public ownership of these habitats would help protect them in perpetuity. Acquiring these habitats would also facilitate more efficient and effective restoration and management of lands and waters within these boundaries by leading to larger blocks of contiguous habitat which can be managed and protected. The MS TIG proposes to allocate up to \$4.2M from its WCNH resource category and up to \$1.8M from its Birds resource category to fund this activity. Acquisition would continue with available funding for up to 15 years. Additional details for Alternative B are provided in Chapter 3 of this document.

2.6.1.3 Alternative C: Grand Bay Habitat Management (up to 17,500 acres)

The goal of Alternative C is to implement management activities, which are further described in Chapter 3, on up to 17,500 acres of current publicly owned land over the course of 15 years. Target habitats would include coastal marsh, beach, freshwater marsh, savannas and flatwoods, and forested freshwater scrub-shrub and bird habitats in Grand Bay NWR, Grand Bay NERR and Grand Bay Savanna CP. The MS TIG would propose to allocate up to \$4.2M from its WCNH resource category and up to \$1.8 M from its Birds resource category to fund this activity.

2.6.1.4 Alternative D (Preferred): Grand Bay Land Acquisition (up to 8,000 acres) and Habitat Management (up to 17,500 acres)

Alternative D proposes to implement both habitat acquisition and restoration (a combination of alternatives B and C) to help restore injuries to WCNH and birds in the Mississippi Restoration Area. The primary objective of coastal land acquisition and restoration is to protect important contiguous lands and waters in an effort to maximize efficiencies and effectiveness in restoring and managing those habitats for the benefit of coastal resources. Implementing these activities within the proposed alternative boundary is consistent with and supports the mission and goals of the Grand Bay NWR, Grand Bay NERR and Grand Bay Savanna CP management plans and initiatives. For Grand Bay NWR acquisition and restoration measures and management activities have been developed in plans which incorporated a public vetting process and analyses under NEPA^{26,27}. While land acquisition

 $\frac{https://www.fws.gov/southeast/planning/PDFdocsLandAcquisition/Grand\%20Bay\%20Final\%20LPP\%20EA/GrandBay4_Final_LPP\%20Formatted.pdf$

.

²⁶ Grand Bay National Wildlife Refuge comprehensive conservation plan, available at https://www.fws.gov/southeast/planning/CCP/GrandBayFinalPg.html.

²⁷ Land protection plan and final environmental assessment for the expansion of Grand Bay National Wildlife Refuge, available at

alone can be a valuable habitat protection tool, habitat management is often necessary for landscape-level conservation to allow connectivity with adjacent habitats. Private inholdings and associated land use and structures within the project boundary for proposed Alternative D create challenges for landscape-level habitat management in the area. The MS TIG therefore believes a combined strategy of land acquisition and habitat management represents the most comprehensive approach to help restore injuries to WCNH at this site, as well as maximizing the potential to provide services to injured bird species within target habitats affected by the proposal. Prioritizing public ownership of acquisitions ensures permanent protection of the MS TIG's investment. Collaborating with managers and staff at Grand Bay NWR, Grand Bay NERR, and Grand Bay Savanna CP would constitute a valuable partnership in reaching MS TIG goals. Acquisition and management would be implemented with available funding for up to 15 years. The estimated cost for this alternative is \$6 M. The MS TIG proposes to allocate up to \$4.2M from its WCNH restoration type and up to \$1.8 M from it Birds Restoration Type for this activity. Additional details for proposed Alternative D are provided in Section 3 of this document.

2.6.1.5 Natural Recovery/No Action

Under this alternative, no additional restoration to restore injuries to WCNH/Birds in the Mississippi Restoration Area using NRDA funding would be done by the MS TIG at this time. The Natural Recovery/No Action Alternative for the WCHN/Birds Restoration Type is described in detail in Section 3.2

2.6.2 Restoration Type NR (Nonpoint Source)

The development of the reasonable range of alternatives for the NR Restoration Types is discussed here. Screening of NR projects resulted in identification of two projects for development as the reasonable range of alternatives:

Alternative A (Preferred): Upper Pascagoula River Water Quality Enhancement Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan Natural Recovery/No Action (No Action)

2.6.2.1 Alternative A (Preferred): Upper Pascagoula River Water Quality Enhancement

The health of the Gulf of Mexico depends upon the health of its estuaries, and the health of those coastal waters is influenced by land use upstream along tributary rivers. Runoff from cropland, pasture/grassland, associated agriculture lands and forestland contributes nutrients and sediment that adversely impact the health of coastal waters of the Gulf. While agricultural and forested lands are not the sole contributors of nutrients to coastal waters, there are tremendous opportunities to address this resource concern at its source. The primary goal of the Upper Pascagoula River Water Quality Enhancement alternative is water quality improvement through the development and implementation of conservation plans and practices to reduce nutrient and sediment runoff into coastal waters. The Chunky-Okatibbee watersheds were selected for implementation of the proposed Alternative A based on sediment load contributions to coastal waters. Alternative A would be implemented by the United States Department of Agriculture (USDA), Natural Resource Conservation Service (NRCS) in Mississippi (USDA-NRCS). The USDA-NRCS would provide outreach and technical assistance to voluntary participants (landowners), especially on the most vulnerable acres in the watersheds, to

develop conservation plans and would use all conservation practices as shown in Appendix B and typically planned and funded by USDA-NRCS programs. The USDA would develop conservation plans within a 20,000-acre area with a priority on opportunities in critical areas for the greatest reduction in nutrient losses that are also within one mile of tributaries. Given the success of USDA-NRCS Farm Bill programs, their strong acceptance by private landowners, and the existence of an effective program execution, there is a significant opportunity to implement conservation practices, especially in critical acres, on private lands that will reduce the levels of nutrients and sediments entering the Gulf of Mexico. Upper Pascagoula River Water Quality Enhancement would be a 5-year program. Conservation practices, especially those systems that avoid, control and trap nutrient and sediment losses, would be implemented on cropland, pasture/grassland, forestland, and associated agriculture land within the Okatibbee-Chunky watersheds. The estimated cost for this preferred alternative is \$4.0 M. The MS TIG proposes to allocate \$4.0 M from its NR (Nonpoint Source) Restoration Type for this activity, and the USDA will invest \$1.0 M of Farm Bill funding in the same watershed. Additional details for Proposed Alternative A are provided in Section 3 of this document.

2.6.2.2 Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan

The Pascagoula River Basin Riparian Buffer Maintenance Plan (Alternative B) would provide outreach and technical assistance to voluntary participants (landowners) to develop conservation plans in riparian areas and would use all conservation practices as shown in Appendix B and typically planned and funded by USDA-NRCS programs. Riparian buffers act to partially protect streams from the impact of adjacent land uses. Buffers increase water quality in associated streams as sediment and runoff is intercepted. Riparian buffers also serve to provide habitat and reduce bank erosion by providing bank stabilization. With planning and monitoring, riparian buffers will help control channel instability, head-cutting, mass slumping, and wetland degradation. Like Alternative A, USDA-NRCS would develop conservation plans within a 20,000-acre area with priority on opportunities that are within one mile of tributaries. Ecological/NR conservation practices would be implemented in riparian areas within forestland and associated agriculture lands and forests on farmsteads in the Chunky- Okatibbee watersheds in Mississippi. This alternative would be a 5-year program. The estimated cost for this preferred alternative is \$4 M which would be allocated from the NR Restoration Type. USDA-NRCS will invest \$1.0 M of Farm Bill funding in the same watershed. Additional details for proposed Alternative B are provided in Section 3 of this document.

2.6.2.3 Natural Recovery/No Action

Under this alternative, no additional restoration to enhance water quality by NR (Nonpoint Source) in the Mississippi Restoration Area using NRDA funding would be done by the MS TIG at this time. The Natural Recovery/No Action Alternative for the NR Restoration Type is described in detail in Section 3.8.

2.7 Alternatives Evaluated in this Draft RP/EA

The following alternatives are evaluated in Section 3 under both OPA and NEPA. The map below (Figure 2.7-1) shows the locations of the proposed project areas for proposed alternatives that are evaluated in this RP/EA. A summary is also provided below.




Figure 2.7-1: Project Areas for the Proposed Reasonable Range of Alternatives for this Draft RP/EA

Wetlands, Coastal, and Nearshore Habitats/Birds

The reasonable range of alternatives for the WCNH/Birds Restoration Types includes five proposed alternatives including the Natural Recovery/No Action. There is one alternative for Graveline Bay Land Acquisition and Management, three alternative means of accomplishing Grand Bay Land Acquisition and Habitat Management, and the Natural Recovery/No Action Alternative. The WCNH/Birds alternatives for the Draft RP/EA described in the above sections are listed here:

- Alternative A (Preferred): Graveline Bay Land Acquisition and Management-Preferred
- Alternative B: Grand Bay Land Acquisition (up to 8,000 acres)
- Alternative C: Grand Bay Habitat Management (up to 17,500 acres)
- Alternative D (Preferred): Grand Bay Land Acquisition (Up to 8,000 acres) and Habitat Management (Up to 17,500 acres)
- Natural Recovery/No Action Alternative

NR-Nonpoint Source

The reasonable range of alternatives for NR (Nonpoint Source) Restoration Type includes three proposed alternatives including the Natural Recovery/No Action. The proposed alternatives for the NR Restoration Type for the Draft RP/EA are described in the above sections are listed here:

- Alternative A (Preferred): Upper Pascagoula River Water Quality Enhancement
- Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan

• Natural Recovery/No Action Alternative

3.0 OPA Evaluation of Alternatives and NEPA Affected Environment and Environmental Consequences

This section analyzes the proposed alternatives evaluated under OPA for the Restoration Types and proposed for selection in this Draft RP/EA. Section 3.1.1 provides an OPA evaluation for the WCNH/Birds Restoration Type. Section 3.8.1 provides an OPA evaluation for the Nutrient Reduction (Nonpoint Source) Restoration Type.

Similarly, Section 3.3.1 and 3.4.1 provides the NEPA affected environment and environmental consequences for proposed WCNH/Bird Restoration Type alternatives, and Section 3.9.1 provides the NEPA affected environment and environmental consequences for proposed NR (Nonpoint Source) Restoration Type alternatives.

The MS TIG elected to prepare a programmatic analysis of the environmental consequences of the range of alternatives developed for the selected Restoration Types, (1) to consider the multiple related actions that would occur because of this restoration planning effort, and (2) to allow for a better analysis of cumulative impacts. Prior to implementation of restoration measures, management activities, and conservation practices, site-specific environmental evaluations as described herein would be conducted. So long as the adverse impacts of particular site-specific restoration measures and management activities are at or below the levels described in this RP/EA, no additional environmental assessments or environmental impact statements would be required before implementation. Should site-specific environmental evaluation indicate the potential for significant adverse effects or effects beyond those disclosed in this RP/EA, an EA or EIS would be prepared, or the site-specific project would be modified so that the level of impacts were at or below the levels described in this RP/EA or other site-specific project(s) explored.

3.1 WCNH/Bird Restoration Type

Section 3.1.1 provides the OPA evaluation for the No Action Alternative and WCNH/Bird Alternatives A-D. Land acquisition and related habitat management would be dependent on willing sellers, successful acquisition, and planning of restoration activities and management measures. Section 3.1.2 also describes the programmatic nature of the NEPA analysis for WCNH/Bird alternatives A-D as well as MS TIG approach for NEPA review after restoration measures and management activities have been identified for specific parcels.

3.1.1 OPA Evaluation for WCNH/Birds

The proposed project alternatives are consistent with the Restore and Conserve Habitat Programmatic Goal, the WCNH/Birds Restoration Type, the Replenish and Protect Living Coastal and Marine Resources Programmatic Goal, and the Birds Restoration Type in the PDARP/PEIS. Table 3.1-1 provides an OPA evaluation of each proposed alternative and the No Action alternative using the standard OPA evaluation criteria described in OPA Section 990.54.

These OPA evaluation criteria are listed below:

- The cost to carry out the alternative (The Cost).
- The extent to which each alternative is expected to meet the Trustees' goals and objectives in returning the injured natural resources and services to baseline and/or compensating for interim losses (Restoration Goals and Objectives).
- The likelihood of success of each alternative (Likelihood of Success).
- The extent to which each alternative will prevent future injury as a result of the incident and avoid collateral injury as a result of implementing the alternative (Avoidance of Further Injury/Collateral Injury).
- The extent to which each alternative benefits more than one natural resource and/or service (Multiple Resource Benefits).
- The effect of each alternative on public health and safety (Public Health and Safety).

Table: 3.1-1: OPA Evaluation Criteria

Alternatives	OPA Evaluation Criteria
	Cost
Natural Recovery/No Action Alternative	There is no financial cost associated with the No Action Alternative .
Alternative A: Graveline Bay Land Acquisition and Management	Alternative A: The cost of \$11.5 M for land acquisition, management and monitoring is reasonable for the proposed Alternative A. The Implementing Trustees, through individual and partnering agency experience, have implemented similar projects and anticipate that implementation of Alternative A would result in benefits to WCNH and birds and would provide connectivity benefits. Parcel acquisition costs would be based on appraised value and any management and monitoring contracts would be subject to either MS or Federal acquisition regulations to ensure open competition and competitive pricing.
Alternative B: Grand Bay Land Acquisition; Up to 8,000 acres; Alternative C: Grand Bay Habitat Management (up to 17,500 acres); Alternative D: (Alts. B+C)	Alternatives B-D: The cost of \$6 M for Alternative B, land acquisition (up to 8,000 acres), Alternative C, habitat management (up to 17,500 acres) or Alternative D (B+C) where ultimate funding for habitat management (C) would depend, in part, on funding used for acquisition (B) is reasonable for the proposed alternatives, and is based on the costs of similar acquisition and habitat management projects conducted in the area. The MS TIG anticipates that funding would result in benefits to WCNH and birds, and would provide connectivity benefits. For Alternative B, the cost would allow for the acquisition of more acreage without the benefits of habitat management. Parcel acquisition costs would be based on appraised value. For Alternative C, more acres of habitat could be managed without the benefit of preserving additional habitat through acquisition. Any management and monitoring contracts would be subject to either MS or Federal acquisition regulations to ensure open competition and competitive pricing. Although Alternative D would provide for less acreage in acquisition and habitat management, it would provide more flexibility for strategic and opportunistic acquisition, while focusing appropriate habitat management measures on parcels to maximize the cost per unit of benefit. Further, similar to Alternatives B and C, land acquisition would be based on appraised values and management and monitoring would be subject to either MS or Federal acquisition regulations to ensure open competition and competitive pricing.

Alternatives	OPA Evaluation Criteria
	Restoration Goals and Objectives
Natural Recovery/No Action Alternative	The No Action Alternative is not consistent with the MS TIG's goal to pursue restoration projects that would provide restoration benefits to WCNH and birds.
Alternative A: Graveline Bay Land Acquisition and Management	Alternative A has a clear nexus to the WCNH/Bird injuries described in the PDARP/PEIS because it would result in the acquisition and restoration of interrelated and biologically diverse habitats, which were injured as a result of the spill. Restoration measures would include land acquisition and preservation of WCNH including estuarine marsh, beach, beech-magnolia forest, coastal plain small stream forest, fire suppressed pine savanna, and open water including tidal creeks and bayous habitats. Land acquisition and preservation would provide habitat connectivity by expanding state ownership of parcels near and adjacent to Graveline Bay Marsh Preserve, where the threat of development is high. Land acquisition and preservation would also serve to restore and conserve bird nesting and foraging habitat for bird species that were injured in the DWH oil spill. Restriction of vehicle traffic from sensitive shoreline areas would improve shorebird nesting success, and the bay, marsh, adjoining upland forest, and undeveloped beach front near the mouth of Graveline Bayou are an important landing area for neotropical migrant birds. Direct habitat management measures would include chemical treatment, mechanical treatment, prescribed fires, debris removal and road removal/culvert replacement to enhance these habitats for use by many species and to restore them to more natural condition. Further, Alternative A is consistent with existing MS TIG goals and objectives that focus on the ues of existing management plans and initiatives, leveraging DWH funds, and habitat connectivity. This alternative meets these goals by providing habitat connectivity with the Grand Bay NWR/NERR/Savanna CP, the Sandhill Crane NWR and other wildlife corridors adjacent the project area for the proposed alternative. Leveraging would include NFWF funding for acquisition and management in the project area for Alternative A.

Alternatives OPA Evaluation Criteria Alternative B: *Grand Bay Land Acquisition; Up to 8,000* **Alternatives B-D** have a clear nexus to the WCNH/Bird injuries described in the PDARP/PEIS because they would result in the Alternative C: Grand Bay Habitat Management (up to acquisition and/or restoration, or both, of interrelated and biologically diverse habitats, which were injured as a result of 17,500 acres); **Alternative D:** (Alts. B+C)the spill. Alternative B restoration measures would include land acquisition and preservation of WCNH including coastal marsh, beach, freshwater marsh, pine savanna flatwoods, forested freshwater scrub-shrub, and open water including tidal creeks and bayous. Acquisition and preservation would reduce the threat of development and would provide habitat connectivity to other large conservation parcels in the area. Land acquisition and preservation would also provide services to bird species that were injured in the *DWH* oil spill. Further, this alternative is consistent with existing MS TIG goals and objectives and would result in the acquisition and preservation of land that would expand habitat protection in the project area for Alternative B. Leveraging would include RESTORE funding and NFWF funding for acquisition in the project area for Alternative B. Alternative C would directly benefit WCNH and Birds by restoring habitats using techniques that have been well established by Grand Bay NWR and NERR resource managers would be implemented to restore the structure and function of target habitats within the project area for Alternative C, thereby restoring ecosystem services to WCNH and to birds, fish and other wildlife injured by the *DWH* oil spill. Direct habitat management measures would include chemical treatment, mechanical clearing, controlled burns, debris removal and road removal/culvert replacement. Futher, alternative C is consistent with existing MS TIG goals and objectives and would support enhancement of up to 17,500 acres of WCNH, including habitat used by bird species injured by the *DWH* oil spill. Alternative C would leverage NFWF funding already awarded for habitat management in the Alternative C project area. Alternative D (B+C) has a clear nexus to the WCNH / Bird injuries described in the PDARP/PEIS because it would result in the acquisition and restoration of interrelated and biologically diverse habitats, which were injured as a result of the DWH oil spill. This alternative would provide collective habitat connectivity with the Sandhill Crane NWR and Graveline Bay CPs as well as several other wildlife corridors in the area. Elements of Alternative D are discussed in the Grand Bay NERR Management Plan, the Grand Bay NWR Land Protection Plan and the Comprehensive Conservation Plan. Alternative D would leverage NFWF and RESTORE Act funding already awarded for habitat acquisition and management in the Alternative D project area. By combining Alternatives B and Alternative C, Alternative D would provide maximum benefits to WCNH and birds through the strategic targeted combination of land acquisition and habitat management practices, selected by the resource managers with site specific expertise and experience.

Alternatives	OPA Evaluation Criteria
	Likelihood of Success
Natural Recovery/No Action Alternative	The No Action Alternative would not contribute to restoring, replacing, or enhancing injured natural resources and would not provide for compensation of interim natural resource losses that occurred as result of the <i>DWH</i> oil spill.
Alternative A: Graveline Bay Land Acquisition and Management; Alternative B: Grand Bay Land Acquisition; Up to 8,000 acres; Alternative C: Grand Bay Habitat Management (up to 17,500 acres); Alternative D: (Alts. B+C)	Alternatives A-D: The Implementing Trustees, through individual and partnering agency experience have successfully implemented projects similar to the proposed project alternatives (land acquisition/preservation and habitat management) in the Graveline Bay area, the NWR, and other similar habitats in the CP system, and proposed restoration activities take advantage of similar ongoing work in these and other nearby areas. This documented experience and successful completion of land acquisition and habitat management projects demonstrates that the proposed project alternatives would have a high likelihood of success. The MS TIG would ensure compliance with all applicable federal laws, regulations and executive orders prior to project implementation, and would conduct all necessary agency consultations for NEPA compliance. The proposed alternatives would meet all OPA and NEPA requirements as discussed in Sections 3.0 and 4.0 of this Draft RP/EA. The proposed restoration projects and practices do not generally harm species or their habitats.
	Avoidance of Further Injury/Collateral Injury
Natural Recovery/No Action Alternative	The No Action Alternative would not cause further injury, but would also not provide benefit to offset interim losses.
Alternative A: Graveline Bay Land Acquisition and Management; Alternative B: Grand Bay Land Acquisition; Up to 8,000 acres; Alternative C: Grand Bay Habitat Management (up to 17,500 acres); Alternative D: (Alts. B+C)	Alternatives A-D: There would be minor to moderate impacts from implementing various restoration measures; however, restoration measures would result in long-term benefits to WCNH and the birds that utilize the habitats. Acquisition and management of large parcels of land would result in benefits to WCNH and birds injured in the <i>DWH</i> oil spill that rely on these habitats and therefore would likely prevent ongoing and future injuries to the same types of habitats and resources affected by the <i>DWH</i> oil spill. The risk of collateral injury would be minimized by the use of Best practices (as described in Sections 3.3.1 and 3.4.1) that would be considered in developing parcel specific management actions and during the implementation of habitat management activities.
	Multiple Resource Benefits
Alternative A: Graveline Bay Land Acquisition and Management; Alternative B: Grand Bay Land Acquisition; Up to 8,000	The No Action Alternative could provide for multiple resource benefits; however, recovery rates of multiple resources would be less than if the MS TIG pursued active restoration activities included in the Proposed Actions. Alternatives A-D: Acquisition and preservation of lands in the Graveline Bay CP and the Grand Bay project area for the proposed alternatives would provide multiple resource benefits.
acres; Alternative C: Grand Bay Habitat Management (up to 17,500 acres); Alternative D: (Alts. B+C)	The proposed alternatives would include the acquisition of land adjacent to other large conservation parcels in the area owned and managed by the state, serving to increase habitat connectivity by reducing the threat of development. The combination of acquisition and management of these parcels at the landscape scale will provide benefits including the direct enhancement of habitat, to both WCNH as well as service to bird species injured by the <i>DWH</i> oil spill. Additionally, the habitats protected, restored and enhanced under these

Alternatives	OPA Evaluation Criteria
	alternatives provide food, shelter, breeding, and nursery habitat for many ecologically and economically important animals, including fish, shrimp, birds, and terrestrial mammals. Direct habitat management measures include prescribed fire, mechanical treatment, chemical treatment, access restriction, debris removal, and road removal/culvert replacement.
	Public Health and Safety
Natural Recovery/No Action Alternative	Under the No Action Alternative any potential public health and safety issues or concerns that exist under current and future natural resource management activities would likely remain the same.
Alternative A: Graveline Bay Land Acquisition and Management; Alternative B: Grand Bay Land Acquisition; Up to 8,000 acres; Alternative C: Grand Bay Habitat Management (up to 17,500 acres); Alternative D: (Alts. B+C)	Alternatives A-D: Effects on public health and safety would include minor short-term impacts resulting from prescribed fires. However, there would be long-term benefits to public health and safety from acquisition, preservation and management of parcels in the floodplain that could be developed if they were not acquired through the proposed alternatives. Restored hydrology resulting from road removal/culvert replacement provides a flood risk/public safety benefit by enhancing floodplain functions. The proposed alternative would have a beneficial effect to the surrounding communities. It would promote healthy lifestyles by allowing recreational use on previously private parcels of land.

Project Alternatives A, B, C, and D would meet the evaluation criteria established by OPA because:

- Cost estimates are reasonable, based on the experience of the MS TIG and project partners on similar acquisition and habitat management projects completed in the area;
- The project alternatives have a clear nexus to the WCNH/Bird injuries described in the PDARP/PEIS and the MS TIG's restoration goals and objectives (use of existing management plans and initiatives, leveraging *DWH* funds and providing habitat connectivity) would be met;
- The MS TIG Trustees and project partners have substantial experience successfully implementing similar projects to the proposed project alternatives (land acquisition/preservation and habitat management) in the Graveline Bay area, the NWR, and other similar habitats in the CP system, and proposed restoration activities take advantage of similar ongoing work in these and other nearby areas. This documented experience and successful completion of land acquisition and habitat management projects demonstrates that the proposed alternatives would have a high likelihood of success;
- Acquisition and management of large parcels of land would result in benefits to WCNH and birds injured in the *DWH* oil spill that rely on these habitats and therefore would likely prevent ongoing and future injuries to the same types of habitats and resources affected by the *DWH* oil spill. Future and collateral injury would be avoided by employing best practices in project implementation;
- Each alternative is likely to benefit more than one resource; and
- There would be a long-term benefit to public health and safety from preserving parcels in the floodplain that otherwise might be developed.

Project Alternatives A, B, C and D are also consistent with the MGCRP and other regional planning initiatives and approved management plans being implemented within the Grand Bay NWR, NERR and CP and Graveline CP project areas. Acquiring and/or restoring biologically diverse habitats

demonstrates a nexus between injury and the restoration goals. Future management planning and implementation of acquired properties would not require additional OPA evaluation.

3.1.2 NEPA Analytical Approach for WCNH/Birds Restoration Type

This section provides the NEPA analytical approach for the WCNH/Birds Restoration Type(s) including:

- 1. a description of the general NEPA analytical approach for the WCNH/Birds project alternatives;
- 2. the MS TIG plan for site-specific NEPA review for the selected alternative; and
- 3. the organization of the affected environment and environmental consequences for WCNH/Birds Restoration Type.

1) The NEPA Analytical Approach for the Development of WCNH/Birds Project Alternatives: Proposed WCNH/Birds Alternatives A-D, include acquisition and management of habitat that would benefit the target Restoration Types. If the preferred alternative(s) are ultimately selected in the final RP/EA, the implementing agency would begin willing landowner identification, title surveys, appraisals, etc. and acquisitions²⁸. Acquisition of parcels will only be made at appraised value. Additionally, if the preferred alternative(s) are selected, habitat inventories, restoration planning and restoration measures and management activities would be developed for newly acquired land and current publicly owned parcels consistent with existing management plans. The size and location of these acquisitions would depend on successful negotiations to acquire targeted parcels and therefore the extent of the potential adverse and beneficial impacts are evaluated in this Draft RP/EA as a range of potential impacts. Further, restoration measures and management activities would be implemented on a site-specific basis and would vary for each depending on the current condition of the habitat on that site.

The environmental consequences analysis in this Draft RP/EA would be corroborated by a site-specific review because the exact parcels and associated restoration measures and management activities that would be most appropriate on those parcels are not known at this time. The environmental consequences in the Draft RP/EA are based on the extent of the anticipated restoration measures and management activities contemplated on parcels for proposed alternative project areas. This analysis provides a maximum impact to each of the resource categories based on the MS TIG's knowledge of the proposed alternative project area. This Draft RP/EA also presents a process that the MS TIG would follow to complete the requirements of NEPA and other environmental statutes as site-specific restoration measures and management activities are planned. The process is described in more detail below.

2) The MS TIG Approach to Site-Specific NEPA Review for the Selected Alternative: In the future, the Implementing Trustee would perform additional environmental reviews once parcels and site-specific restoration measures and management activities are developed for a site. The following is a description of the proposed approach to NEPA evaluation for future site-specific restoration measures and management activities for the WCNH/Bird alternatives in this Draft RP/EA.

-

²⁸ The act of acquiring individual parcels would not require parcel-specific NEPA evaluation

- Future NEPA evaluations would be conducted by the Implementing Trustee or their designee by completing an Environmental Evaluation that would document whether impacts are at or below the maximum impacts described in the Final RP/EA. An example of an Environmental Evaluation Worksheet is attached as Appendix A.
- If impacts from the site-specific restoration measures and management activities are at or below the maximum impacts described in the Final RP/EA, then the Implementing Trustee/designee would route the finalized Environmental Evaluation Worksheet through the MS TIG for inclusion in the project's Administrative Record.
- If impacts from the site-specific restoration measures and management activities are above maximum impacts described in the Final RP/EA (e.g. exceed), then the Implementing Trustee/designee²⁹ would notify the MS TIG and conduct additional environmental planning in the form of an environmental assessment on behalf of the MS TIG for TIG review and approval. As an alternative, the Implementing Trustee/designee could re-design the restoration measures and management activities to ensure that they are below the maximum impacts described in the Final RP/EA.
- 3) Organization of the Affected Environment and Environmental Consequences for WCNH/Birds Restoration Type: Guidelines for NEPA impact determinations for the Final PDARP/PEIS are described in Section 6.3.2 of the PDARP/PEIS and are incorporated here by reference. The intensity definitions are used in this RP/EA for identifying adverse impacts of the proposed restoration approaches. The analysis uses the intensity definitions in evaluating whether the proposed restoration approaches may result in minor, moderate, or major adverse impacts. WCNH/Birds Alternatives A, B, C and D include land acquisition, habitat management or the combination of both. The NEPA affected environment and environmental consequences for the WCNH/Birds Restoration Type is structured as follows:
 - Section 3.2 Natural Recovery/No Action Alternative for WCNH/Bird Restoration Type
 - Section 3.3 Alternative A (Preferred): Graveline Bay Land Acquisition and Management
 - Section 3.4 Alternative B (Grand Bay Land Acquisition), C (Grand Bay Habitat management), and Alternative D (Preferred): Grand Bay Land Acquisition and Habitat Management
 - Section 3.5 Cumulative Impacts for WCNH/Birds Alternatives
 - Section 3.6 Comparison of Alternatives

3.2

Natural Recovery/No Action

In addition to the proposed alternatives listed above for the WCNH/Birds Restoration Type, the MS TIG evaluated the Natural Recovery/No Action Alternative (No Action alternative). NEPA [§ 1502.14(d)] requires consideration of a No Action Alternative as a basis for comparison of potential environmental consequences of the action alternatives. The Natural Recovery/No Action Alternative evaluation under NEPA parallels a natural recovery alternative under OPA. OPA regulations also require that "trustees must consider a natural recovery alternative in which no human intervention would be taken to directly restore injured natural resources and services to baseline" (40 CFR §

²⁹ If the Implementing Trustee is MDEQ, then the designee for the purposes of conducting the follow up environmental assessment must be one of the MS TIG's federal trustees.

990.53(b)(2)). The OPA alternatives analysis (which included the No Action alternative) was presented above in Table 3.1-1.

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur, which could result in one of four outcomes for injured resources: 1) gradual recovery, 2) partial recovery, 3) no recovery, or 4) further deterioration. Although injured resources could presumably recover to or near baseline conditions under this scenario, recovery would take much longer compared to a scenario in which restoration actions were undertaken.

Under the No Action alternative, the NRDA Early Restoration projects already approved (Hancock County Marsh Living Shoreline; Restoring Living Shorelines and Reefs in Mississippi Estuaries; and Enhanced Management of Avian Breeding Habitat Injuries by Response in the Florida Panhandle, Alabama and Mississippi) would be the only NRDA restoration implemented for the WCNH/Birds Restoration Type in the MS Restoration Area at this time.

This alternative would have no beneficial impacts to WCNH/Birds because this alternative would largely result in a continuation of the conditions described in the PDARP/PEIS Chapters 3, Ecosystem Setting and Chapter 4, Injury to Natural Resources, and there would be no associated benefits to WCNH/Birds. Under the No Action alternative, some WCNH recovery could result from other *DWH* funded projects which propose acquisition and habitat management in the Grand Bay and Graveline Bay proposed project areas (RESTORE and NFWF GEBF), but not from the federal action being evaluated in this Draft RP/EA. Even if funding and implementation of other *DWH* projects does occur in the project areas, the full suite of WCNH/Birds restoration benefits would not be realized due to diminished funding and the lost opportunity for leveraged funding. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

When analyzed in combination with other past, present and reasonably foreseeable future actions, the No Action alternative would provide no beneficial impacts, because existing conditions would not change. This alternative is not expected to contribute to short-term or long term, cumulative adverse impacts to physical resources, biological resources, or socioeconomics, with the following exception. For the proposed Alterative A (Preferred) Graveline Bay Land Acquisition and Management, without NRDA funding for acquisition, it is likely that these properties would be developed. There is a threat of development of privately-held land adjacent to the proposed alternative, including areas which are designated as new growth areas by the cities of Ocean Springs and Gautier. Acquisition and preservation of land in perpetuity would prevent land development in floodplains and loss of habitat. For Alternative D (Preferred) Grand Bay Land Acquisition (up to 8,000 acres) and Habitat Management (Up to 17,500 acres), the No Action alternative would result in a lack of contiguous parcels acquired for large-scale prescribed fire management. Therefore, by preventing the acquisition and habitat management of parcels in the Alternative A and Alternative D areas, the No Action alternative would have an adverse long-term minor to moderate impact to geology and substrates, and habitats. The No Action could result in long-term, minor to moderate adverse impacts to floodplain as well as public health and safety related to floodplain filling for the proposed alternative A.

3.3 Graveline Bay Land Acquisition and Management-Background and Project Description

The Graveline Bay Land Acquisition and Management proposed alternative includes acquiring parcels near publicly owned lands in the Graveline Bay CP in Jackson County, Mississippi. Habitat management measures that are currently used on the adjacent public lands are also planned including prescribed fire. The proposed alternative would be implemented at proposed locations in Graveline Bay (Figure 3.3-1). The project planning process has been a collaboration between the MDMR and the MS TIG. Potential acquisitions from willing sellers in the proposed alternative area include approximately 1,410 acres of habitat that could be acquired. Estuarine marsh, shoreline (beach), and other coastal riparian habitats are in the proposed alternative project area, some of which are expected to provide foraging, loafing and nesting for bird species injured by the *DWH* oil spill. The estimated cost to implement this proposed alternative is \$11.5 M. The proposed alternative would be located on parcels adjacent to and near Graveline Bay in Jackson County, Mississippi. The parcels are located in Sections 4, 5, 9, 10, 15, and 16 of Township 8 South, Range 7 West (Figure 3.3-1).




Figure 3.3-1: Graveline Bay Land Acquisition and Management –Parcels and Habitats

Graveline Bay and Bayou is over 2,100 acres and represents one of the few relatively undisturbed estuarine bays and small tidal creeks in Mississippi. Graveline Bay coastal wetland and nearshore habitats include estuarine marsh, beach, beech-magnolia forest, coastal plain, small stream forest, fire-suppressed pine savanna, and open water including tidal creeks and bayous (Table 3.3-1). Wetlands, coastal, and nearshore habitat in this area is utilized for foraging, nesting and loafing by bird species injured by the *DWH* Oil Spill. Acquisition and management of parcels in the proposed alternative project area would provide benefits to wading birds and other species. Beach habitat enhancements would benefit nesting shorebirds injured by the *DWH* Oil Spill. The coastal bay and estuarine marsh system of this area consists largely of black needle rush (*Juncus roemerianus*) dominated marsh along its entire length. Smooth cordgrass (*Spartina alterniflora*) occurs largely as narrow (1-3 m) bands along the creeks and bayous. The area supports salt marsh, brackish marsh, and several oyster beds.

Table 3.3-1: Proposed Graveline Bay Land Acquisition and Management - Habitats and Ownership³¹

Habitat	Publicly Owned (acres)	Privately Owned (acres)	Total Acreage of Habitat
Estuarine Marsh	582	636	1,218
Beach	1	5	6
Beech Magnolia Forest	0	115	115
Fire-suppressed pine savanna	36	460	496
Coastal Plain Small Stream Forest	0	66	66
Open water, Tidal Creeks and bayous	156	128	284
Total	775	1,410	2,185

Development is a threat to habitats adjacent to the CP. Residential developments exist to the north of the proposed alternative project area. Municipal land use plans would allow the forested habitats within the proposed alternative project area to be similarly developed. Without protection, the MS TIG anticipates that future residential development would continue in these areas.

The proposed alternative has several objectives including: acquisition of properties that have a high threat of development; preserving a buffer to keep adjacent marsh habitat intact; and reducing habitat fragmentation and realizing connectivity benefit that would result from habitat management adjacent to existing state-owned CP land, PDARP/PEIS restoration approaches for this alternative include:

- Protect and conserve marine, coastal, estuarine and riparian habitats
- Restore and conserve bird nesting and foraging habitat

-

³⁰ http://www.dmr.ms.gov/mississippi-gems/215-graveline-bay

³¹ Acreage is based on parcels that are targeted for purchase, some of which are within the CP boundary, some are adjacent.

The proposed alternative includes (1) the acquisition and preservation of up to 1,410 acres within and adjacent to the CP, and (2) habitat management of both currently owned CP lands and those which would be acquired as part of the alternative. Acquisition and management would be within the 2,185 acres of total habitat within and adjacent to CP boundaries shown on Table 3.3-2.

Restoration Measures-Methodology and Timing

This proposed alternative would include management of habitats within the proposed alternative project area which includes the CP and newly acquired parcels in and adjacent to the CP. The Implementing Trustee would begin landowner identification, title surveys, appraisals, etc. and acquisitions after final RP/EA approval. Additional data collection on target habitats needed to facilitate restoration and management (e.g., habitat inventories, identification of appropriate restoration measures and management activities, etc.) would also be conducted following approval of the project. Restoration measures and management activities would be implemented on a site-specific basis and may vary across the project area depending on the current condition of habitats. Habitat restoration measures and management activities could include vehicular access restriction on Graveline beach; chemical treatment; mechanical treatment; prescribed fire; debris removal; and road repair/removal and culvert placement, described below. Proposed restoration measures and management activities are summarized in Table 3.3-2 and described below.

Table 3.3-2: Restoration Measures and Management Activities by Habitat

Habitat	Acres of Habitat	Acquisition/Preservation	Access Restriction	Invasive Species Management- Chemical Treatment	Invasive Species Management- Mechanical Treatment	Prescribed Fire	Debris Removal	Road repair/removal and culvert placement
Estuarine Marsh	1,218	X						
Beach	6	X	X	X			X	
Beech Magnolia Forest	115	X		X			X	
Fire-suppressed pine savanna	496	X		X	X	X	X	X
Coastal Plain Small Stream Forest	66	X		X	X		X	

Acquisition and Preservation: Protection of habitats is consistent with the MS TIG goal to increase connectivity of coastal habitats. Lands would be purchased in fee from willing sellers at appraised value. Acquisition and preservation includes the purchase of land and preservation in perpetuity, facilitating protection of habitat through prevention of large scale development. Acquisition of parcels would only be made at appraisal value. Acquisition and preservation would apply to up to 636 acres of estuarine marsh, 5 acres of beach, 115 acres of beech magnolia forest, 460 acres of fire-suppressed pine savanna, and 66 acres of coastal plain small stream forest. This would be a 10-year project.

Access Restriction: Access restriction following acquisition of parcels containing beach habitat would provide protection of shorebird habitat. Barriers would be placed to restrict all vehicle traffic to sensitive shoreline areas. Restricted access would reduce direct impacts from vegetation and sand disturbance, as well as reduce litter, noise pollution, and environmental effects resulting from target shooting and vehicle traffic. Pedestrian access would be allowed.

Invasive Species Management: Invasive species management will focus on prevention, control and eradication of known exotic invasive plant species in the project area for the proposed alternatives. Example species include, but are not limited to, Chinese privet (*Ligustrum sinense*), Chinese tallow (*Sapium sebiferum*), common reed (*Phragmites australis*), Cogon grass (*Imperata cylindrica*), Japanese climbing fern (*Lygodium japonicum*), Japanese honeysuckle (*Lonicera japonica*) and others. A number of techniques are commonly utilized on the NWR and NERR, and at the nearby Sandhill Crane NWR, to accomplish invasive species management are incorporated by reference here (USFWS, 2007, USFWS, 2008, GBNERR, 2016). For example, prescribed fire is used for both reduction of fuel loads and invasive species management in fire suppressed pine savanna to promote grassy-herbaceous ground cover. For the purposes of discussion and to facilitate a programmatic impact analysis, invasive species management techniques will be divided into three categories, which are described below: 1. Chemical Treatment; 2. Mechanical Treatment, and 3. Prescribed Fire. Resource managers could use an integrated approach including a variety of techniques for site specific restoration and management measures depending on existing habitat conditions.

- 1) Chemical Treatment: Chemical treatments could include basal-bark application, cut stump treatments, foliar spray applications, and stem injection of herbicides to target eradication or control of invasive plant species. These applications are completed seasonally and typically occur in small target areas. Activities could require the vehicular transport of personnel into areas, use of approved herbicides, use of established safety and containment procedures, and the targeted application of herbicide in small areas. Personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities. Treatments are typically done in areas that range from several acres up to 50 acres for a large-scale treatment by trained personnel. On Graveline Bay CP, chemical treatment would be limited to small areas within the 6-acre beach for treatment of common reed; the 115-acre beech magnolia forest; in 496 acres for fire-suppressed pine savanna; and selectively within the 66 acres of coastal small stream forest (avoiding streams and aquatic vegetation) for treatment of Chinese tallow, privet, Cogon grass, and other exotic invasive plant species.
- 2) Mechanical Treatment: Mechanical treatment is often used in combination with prescribed fire to restore and maintain openness, recycle nutrients, and reduce woody vegetation. Use of these techniques results in an increase in savanna species including sun-loving graminoids (grass-like plants) and forbs (flowering plants). Mechanical treatment could include removal of trees using commercial tree contracts, chain saws, bulldozing, and use of a bulldozer or gyrotrac with roller chopper to remove shrubs and small trees or drum chopping to push over and crush small, pre-commercial pines and shrubs. In wet areas, soft track or wide track equipment would be used to distribute the equipment weight and minimize ground disturbance. Alternatively, crews access areas on foot and may remove material with chainsaws or by hand. Replanting could also be part of invasive management operations. These techniques can be for large areas and are used successfully; several thousand acres of undesirable vegetation has been cleared by mechanical treatment in the Sandhill Crane NWR

- (USFWS, 2007). Mowing, tilling and disking are also used to prevent the spread of Cogon grass. For the proposed alternative, mechanical treatment would be used within 496 acres of fire-suppressed pine savanna and in 66 acres coastal plain small stream forests. Operations could occur over several seasons depending on the timing of acquisitions and other restoration priorities.
- 3) Prescribed Fire: Native habitats within the southeastern United States, including those within the project boundary, evolved in the midst of reoccurring, natural fires (USFWS, 2007, USFWS, 2008, GBNERR, 2016). These habitats therefore depend on a reoccurring fire schedule. Historically, natural fire occurred on a three to five-year interval. Fires were of low intensity, fueled by grasses and pine litter. Habitat management agencies in the project area successfully use prescribed fires to restore and maintain high quality, natural habitats. Prescribed fires reduce woody vegetation and tree encroachment in pine savanna habitat and can be effective in helping prevent the spread of certain exotic invasive species (e.g., Cogon grass and Chinese tallow), when used in combination with other methods (e.g., chemical and mechanical treatment). This project proposes to implement a schedule of prescribed fires on publicly owned property within the project boundary to accomplish habitat restoration and management goals. Wire grass, for example, is a fire-dependent savanna species. Only after being burned during the growing season will this grass produce seeds. Their complex system of underground roots and shoots helps them survive the fire. By increasing species such as this, the project is also expected to provide services to wildlife that use them, such as many declining populations of grassland bird species that rely on savanna habitat.³² Prescribed fire and associated management within the project boundary would simulate these historic. natural fires.

Site preparation for a prescribed fire often involves compression of vegetation using equipment like roller choppers, gyrotracks, and excavators and/or other mechanical treatments included above to create habitat conditions which facilitate desired burns. Clearing, plowing and disking may be used to prepare fire breaks, zones devoid of fuel that border burn units and help manage burn boundaries. Fire could be applied using handheld drip torches to initiate prescribed fire. Aerial ignition from helicopters could also be used. Prescribed burns would follow standardized planning protocols and methodologies, such as considering environmental factors (certain weather, fuel and moisture conditions that would make the fire manageable³³) and burning on a 2-3 year rotation during the growing season (Spring and Summer months, when possible). Prescribed fires could range in size depending on habitats and logistics. Average prescribed burns at Grand Bay NWR are 79 acres, however, 20% of the Grand Bay burns may reach 100 acres or more. Prescribed fires average 59 acres at Mississippi Sandhill Crane NWR, however, 13% of those may reach 100 acres or more (USFWS 2005). For Alternatives C and D, prescribe fire would applied on up to 6,276 acres of savanna and flatwoods. For the proposed alternative, prescribe fire would applied to up to 496 acres of fire-suppressed pine savanna.

³² https://www.fws.gov/refuge/Grand Bay/what we do/resource management.html

³³ https://www.fws.gov/mississippisandhillcrane/fire.html

Debris Removal: Debris removal would include the use of equipment such as trucks, ATVs, bobcats, chainsaws and other equipment to remove debris such as dead vegetation, garbage, and other refuse. Debris would be disposed of properly at a landfill or other approved site. This would apply to up to 6 acres of beach, 115 acres of beech magnolia forest, 496 acres of fire-suppressed pine savanna, and 66 acres of coastal plain small stream forest.

Road Removal/Repair and Culvert Replacement: These measures include roadbed and culvert removal/replacement, filling and rerouting of drainage ditches, geotextile placement, ditch bank stabilization and other services needed to remove the roadbed. In addition, minor repair of the roadbed could also be required depending on site conditions. Roadbed material would be disposed of properly at a landfill or other approved site. Road repair/removal would apply to up to 4 acres of fire-suppressed pine savanna. Best practices would be implemented including erosion control measures, re-contouring and revegetation of the roadbed after hard surface is removed.

Best Practices: The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS to avoid and minimize impacts to resources during the implementation of restoration measures and management activities described above. Best practices listed in the PDARP/PEIS are intended to evolve as an adaptive management component of implementing the PDARP/PEIS; as such, the appendix to the PDARP/PEIS is a living document. As new best practices are established, existing best practices are refined, or new techniques and information are informed by implementation, these measures will be added to or updated in the relevant websites identified in the appendix of the PDARP. In this capacity, new projects will have available the current range of best practices to support project design and implementation. In addition to PDARP/PEIS best practices, the MS TIG could develop best practices for site-specific restoration measures and management activities in different locations due to differences in relevant site conditions.

3.3.1 Alternative A (Preferred): Graveline Bay Land Acquisition and Management Affected Environment and Environmental Consequences

This section discusses Alternative A (Preferred), the Graveline Bay Land Acquisition and Management proposed alternative. For WCNH/Birds, Alternative A is one of two preferred alternatives that is proposed for implementation.

Graveline Bay Land Acquisition and Management (Proposed Action)

The proposed action includes acquisition of up to 1,410 acres of habitat in the vicinity of the Graveline Bay CP and restoration and management activities on up to 2,185 acres of the proposed alternative project area (existing CP and newly acquired parcels in the vicinity of the CP). Management activities that are anticipated include access restriction, chemical treatment, mechanical treatment, prescribed fire, debris removal, road repair/removal and culvert replacement.

3.3.1.1 Overview of Affected Environment and Environmental Consequences

This analysis incorporates by reference the relevant portions of Section 3.5.1(Nearshore Ecosystem) of the PDARP/PEIS. The PDARP/PEIS provides programmatic evaluation of the environmental consequences of the restoration approaches "Protect and conserve marine, coastal, estuarine and riparian habitats" and "Restore and conserve bird nesting and foraging habitat", which are considered

in this Draft RP/EA. PDARP/PEIS evaluations from sections 6.4.1 and 6.4.10 are incorporated by reference here. Tiering from that analysis, this section presents the Affected Environment of Graveline Bay and environmental consequences of the proposed actions in context of the project-specific affected environment.

The programmatic analysis looked at a series of resources as part of the biological, physical, and socioeconomic environment. As appropriate in a tiered analysis, the evaluation of each alternative focuses on the specific resources with a potential to be affected by the proposed project. To avoid redundant or unnecessary information, resources that are not expected to be affected are evaluated summarily in the respective sections. These resources include noise, marine and estuarine fauna, infrastructure, fisheries and aquaculture, marine transportation, and aesthetics and visual resources which will be discussed in Sections 3.3.1.2, 3.3.1.3, and 3.3.1.4.

3.3.1.2 Physical Environment

<u>Introduction to Affected Environment (Physical Environment)</u>: Geology and Substrates, Hydrology and Water Quality, and Air Quality and Greenhouse Gas Emissions are discussed in this Section. PDARP/PEIS sections 6.4.15.1 is incorporated by reference here. The affected environment for the proposed alternative physical environment is described in respective sections below.

<u>Programmatic Review of Environmental Consequences (Physical Environment)</u>: Sections 6.4.1.5.1 and 6.4.10.1.1 of the PDARP/PEIS describe the impacts to Physical Resources for the relevant restoration approaches and are incorporated by reference and briefly described here.

PDARP/PEIS consequences related to geology and substrates and water resources: Specific restoration activities identified as part of land management plans could result in short-term, minor to moderate adverse effects on geology, substrates, and water resources. Fire management may have short-term adverse impacts on soils, substrates, and air quality. Land acquisition could permit public access for recreational use which could result in short-term, minor to moderate adverse effects through increased soil compaction, rutting, or erosion caused by human presence and activity within the conservation area. Increased public use could result in short-term, minor effects on surface water through increased sedimentation. Fee title land acquisition could reduce disturbance of geology and substrates by protecting lands from development pressure. This would be a long-term beneficial effect that will extend beyond the life of the project. Where protected lands overlap ground water recharge zones, surface water, or brackish-water resources, water sources and water quality could be further protected from future degradation by helping to reduce runoff. Similarly, where protected land overlaps wetlands or shorelines, the protection of natural hydrologic processes could indirectly help limit development and associated effects on water quality, including via saltwater intrusion. These would be long-term beneficial effects.

Environmental consequences for the proposed alternative are within the general range of impacts as described in the PDARP/PEIS with some variances related to specific actions. Table 3.3-3 summarizes the environmental consequences of the proposed alternative on the physical environment. Detailed analyses are provided below the summary table.

Table 3.3-3: Environmental Consequences to the Physical Environment Due to the Proposed WCNH/Birds Alternative A (Preferred)

WCNH/Bird	ls Alternati	<u>ve A (Preferi</u>	red)				
	Acquisition/Preservation	Access restriction	Chemical treatment	Mechanical treatment	Prescribed fire	Debris removal	Road repair/removal and culvert placement
		Geol	logy and Subst	trates			
Adverse Impact Duration	long-term	short-term	short-term	short-term	short-term	short-term	short-term
raverse impact Baration	long term	SHOTE COIN	SHOTE COIN	minor	SHOTE CETTI	SHOTE CHIII	Short term
Adverse Impact Intensity	minor	minor	minor	tomoderate	moderate	minor	moderate
Beneficial Impact							
Duration							long-term
		Hydrole	ogy and Water	· Quality			
Hydrology							
			-1	-1	-1		-1
Adverse Impact Duration			short-term	short-term minor to	short-term minor to		short-term
Adverse Impact Intensity			minor	moderate	moderate		moderate
Beneficial Impact			minor	moderate	moderate		moderate
Duration	long-term						long-term
Water Quality	Ü						<u> </u>
Adverse Impact Duration			short-term	short-term	short-term		short-term
Adverse impact Buration			SHOTE-term	minor to	minor to		SHOTE-term
Adverse Impact Intensity			minor	moderate	moderate		moderate
Beneficial Impact							
Intensity	long-term	short-term					long-term
Floodplains							
Adverse Impact Duration							
Adverse Impact Intensity							
Beneficial Impact							
Duration	long-term						long-term
Wetlands							
Adverse Impact Duration			short-term	short-term	short-term	short-term	short-term
A 1 T T				minor to	minor to	•	minor to
Adverse Impact Intensity Beneficial Impact			minor	moderate	moderate	minor	moderate
Duration	long-term				long-term	long-term	long-term
Duration	10115 101111		_		Tong willi	10116 101111	long term
Air Quality and							
Greenhouse Gas							
Emissions							
Adverse Impact Duration			short-term	short-term	short-term	short-term	short-term
A dament To the transfer of th					minor to		
Adverse Impact Intensity Beneficial Impact			minor	minor	moderate	minor	minor
Duration							
Duration							

As appropriate in a tiered analysis, the evaluation of the proposed alternative focuses on the specific resources with a potential to be affected. Noise impacts for the proposed alternative would be negligible to minor. To avoid redundant or unnecessary information, noise is evaluated here.

Noise: There would be short-term, minor, adverse noise impacts from equipment and operations associated with mechanical treatment, establishment of fire breaks, prescribed fire operations, and road repair/removal and culvert replacement. Restoration activities would occur sporadically and seasonally and would be short in duration. Noise receptors in the area of the work would be buffered by forested areas between the receptor and the site of noise-producing activity. Acquisition and preservation of developable areas would provide a long-term benefit by reducing ambient noise pollution when compared to a build out scenario if property were developed. In addition, the following best practices would be implemented, to the extent practicable, for the selected alternative: minimize construction noise to the maximum extent practicable when working near protected species and their habitats.

For the physical environment, the following resources are further analyzed in this section:

Geology and Substrates Water Quality and Hydrology Air Quality and Greenhouse Gas Emissions

3.3.1.2.1 Geology and Substrates

Affected Environment

Section 3.3.3 of the PDARP/PEIS discusses the geomorphological zones of the northern Gulf of Mexico. The proposed alternative is located within the Gulf Coastal Plain and the Mississippi Alluvial Plain physiographic regions. Seismic activity in the area of the proposed alternative is low. Since the late 1800s, about ten earthquakes large enough to be detected have occurred in the Gulf of Mexico. These earthquakes were mostly small-magnitude events (magnitudes of 3 to 4 on the Richter scale).

Landforms and substrates are generally comprised of Holocene sediments. These sediments are composed of sand, silt, and clay with comparatively high organic matter content. The coastal estuaries of Mississippi are composed of mostly sandy fine-grained sediment, silt and clays (Schmid 2015). The habitats can be divided into two classes - intertidal and subtidal. Intertidal zones (typical tidal range of 0.5 ft.) are generally composed of mud flats and small areas of natural sand beach. In general, the nearshore subtidal habitat is composed mostly of unconsolidated bottom types including sand, muddy sand, and mud bottom.

Data from the Mississippi State Geological Survey generally indicates that surface soils in the area of the proposed alternative consist of Holocene age coastal deposits of loam, sand, gravel, and clay. The USDA-NRCS Web Soil Survey identifies 14 soil-mapping units within the footprint of the proposed alternative. These soil map units located within the proposed alternative footprint area are listed on Table 3.3-4 (NRCS 2016). Of these soils, the Guyton silt loam and Handsboro association soil are listed as hydric and minor inclusions of the Atmore loam, 1 to 3 percent slopes; Benndale fine sandy loam, 0 to 2 percent slopes; Benndale fine sandy loam, 2 to 5 percent slopes; Smithton loam, 0 to 1 percent slopes, occasionally flooded; Ocilla loamy sand, 0 to 2 percent, occasionally flooded; Axis mucky sandy clay loam, frequently flooded; Handsboro mucky silt loam, frequently flooded; Bayou

sandy loam, 0 to 1 percent slopes; and Harleston fine sandy loam, 0 to 2 percent slopes are listed as hydric (NRCS 2016a). Soils characteristics are listed in Table 3.3-4.

Table 3.3-4: Soils Characteristics in the project area for WCNH/Birds Alternative A (Preferred)

Soil Type	Texture	Drainage Class
Atmore loam, 1 to 3 percent	Loam (upper)	Poorly Drained
Slopes	Loam (lower)	·
Benndale fine sandy loam, 0 to	Fine Sandy Loam (upper)	Well Drained
2 percent slopes	Loam (lower)	
Benndale fine sandy loam, 2 to	Fine Sandy Loam (upper)	Well Drained
5 percent slopes	Loam (lower)	
Smithton loam, 0 to 1 percent	Loam (upper)	Poorly Drained
slopes, occasionally flooded	Sandy Loam (lower)	·
Escambia very fine sandy loam,	Very Fine Sandy Loam	Somewhat Poorly Drained
0 to 2 percent slopes	(upper)	
	Loam (lower)	
Ocilla loamy sand, 0 to 2	Loamy Sand (upper)	Somewhat Poorly Drained
percent, occasionally flooded	Sandy Clay Loam (lower)	
Prentiss silt loam, 0 to 2 percent	Silt Loam (upper)	Moderately well Drained
Slopes	Loam (lower)	·
Wadley loamy sand, 0 to 5	Loamy Sand (upper)	Somewhat Excessively Drained
percent slopes	Sandy Clay Loam (lower)	
Axis mucky sandy clay loam,	Mucky Sand Clay Loam	Very Poorly Drained
frequently flooded	(upper)	
	Sandy Loam (lower)	
Handsboro mucky silt loam,	Mucky Silt Loam (upper)	Very Poorly Drained
frequently flooded	Muck (lower)	
Bayou sandy loam, 0 to 1	Sandy Loam (upper)	Poorly Drained
percent slopes	Sandy Loam (lower)	
Harleston fine sandy loam, 0 to	Fine Sandy Loam (upper)	Moderately well Drained
2 percent slopes	Sandy Loam (lower)	·
Harleston fine sandy loam, 2 to	Fine Sandy Loam (upper)	Moderately well Drained
5 percent slopes	Sandy Loam (lower)	
Latonia loamy sand, 0 to 2	Loamy Sand (upper)	Well Drained
percent slopes	Sandy Loam (lower)	

Environmental Consequences for WCNH/Birds Proposed Alternative A (Preferred)

Table 3.3-3 lists environmental consequences to geology and substrates of the activities associated with the proposed alternatives. There would be no adverse effect to geologic resources in the proposed alternative project area from acquisition/preservation, access restriction, chemical treatment, mechanical treatment, prescribed fire, debris removal or road repair/removal and culvert replacement. A review of impacts to substrates (soils) is provided here.

<u>Acquisition/Preservation</u>: Acquisition and preservation would open new areas to recreational activities including hiking, fishing, bird-watching, and camping. Access via motorized vehicles would be limited. The increased public use could result in a long-term, minor, adverse impact to soils due to potential compaction, but these would be limited to relatively small areas.

<u>Access restriction</u>: For beach habitat, barriers would be placed to restrict ATV and vehicle traffic to sensitive shoreline areas. During the placement of barriers and signage, small areas of soils would be disturbed and compacted by personnel and equipment. This would be a short-term, minor, adverse impact to soils.

<u>Chemical treatment</u>: Treatment activities could require the use of ATVs, pickups or other small equipment that could result in soil disturbance, rutting and compaction. The use of equipment would

result in a short-term minor adverse impact to soils. Removal of nuisance species and replanting could result in short-term, minor, adverse impacts to soils.

Mechanical treatment: Activities include but would not limited to use of brush-hog, mowing, disking, and use of chainsaws. In addition, use of gyro-tracks and in some cases bobcats or bulldozers to lay down or remove vegetation could be used. Turning over soils, soil compaction, disturbance and/or rutting from equipment use could result in short-term, minor to moderate, adverse impacts, depending on the size of the operation, soil wetness and season of the operation. To minimize these effects, care would be taken in the selection of equipment used and timing of operations, particularly in wetter soil conditions.

<u>Prescribed fire</u>: Preparations for prescribed fires could include installation of fire breaks, and use of light to heavy equipment to fell or lay down woody underbrush. Fire breaks would be constructed around the boundary of the burn unit by clearing and or disking. Soils would be turned and to exponse mineral underlayers. Soil could be disturbed and compacted during the burn operations due to equipment use. Vegetation laydown/removal operations using light to heavy equipment could result in soil disturbance or rutting. In wet areas, soft track or wide track equipment would be used to distribute the equipment weight and minimize impact. Alternatively, crews may remove material with chainsaws. There could be short-term, moderate adverse impacts from mineral soil exposure, rutting, and soil disturbance during the site preparation and burn operations.

<u>Debris removal</u>: The use of equipment such as trucks, ATVs, bobcats, and other equipment to remove debris such as dead vegetation, garbage, and other refuse could cause compaction of the soil which would result in short-term, minor impacts.

<u>Road repair/removal and culvert placement</u>: Removal of road beds of up to 4 acres would require the use of excavation equipment, dump trucks, and other large equipment. Soils adjacent to the road bed may become compacted. Removing the roadbed would allow soils to return to a more naturally functioning state. Disturbed soils and road surfaces graded and prepared for revegetation. There would be short-term, moderate, adverse impacts to soils during road bed removal. Roadbed areas would be recontoured by disking ad prepared for planting of native vegetation. The operations could provide long-term benefits to soils by restoring more historic hydrologic patterns to soils.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to geology and substrates (soils):

- Allow revegetation of fire breaks or actively revegetate with native species or annual grasses, if prolonged period of greening up is anticipated.
- Develop and implement spill prevention and response plan, including conducting daily inspections during chemical treatment, mechanical treatment and prescribed fire operations to ensure there are no leaks of antifreeze, hydraulic fluid, pesticides or other substances.
- To the extent practicable, for equipment use in wet areas, soft tracked or wide tracked equipment should be used to distribute the equipment weight and minimize impacts to soils. Alternatively, crews may remove vegetative material with chainsaws.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. If these properties were developed, adverse impacts to soils would be expected. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.2.2 Hydrology and Water Quality

Affected Environment

Section 3.3.2 of the PDARP/PEIS addresses river flows on the Northern Gulf geography and water quality. Section 6.14.2 discusses future sea level rise, storm surge and storm intensity projections and is incorporated by reference here. For the proposed alternative, the affected resources consist of shallow water within bays, bayous, and wetlands within Graveline Bay. Mississippi's water quality standards specify the appropriate levels for which various water quality parameters or indicators support a water body's designated use(s). Each use assessed for a water body is determined to be either "Attaining" or "Not Attaining" in accordance with the applicable water quality standards and U.S. Environmental Protection Agency (EPA) guidelines for assessments pursuant to Clean Water Act Section 305(b). A water body's use is said to be impaired when—based on current and reliable site-specific data of sufficient quantity, quality, and frequency of collection—it is not attaining its designated use(s). Where data and information of appropriate quality and quantity indicate nonattainment of a designated use or uses for an assessed water body, the water body will be placed on the Mississippi 2014 Section 303(d) List of Impaired Water Bodies (MDEQ 2014). The proposed alternative is located in the Mississippi Coastal Streams watershed. This watershed has a drainage area of approximately 1,550 square miles (MDEQ 2014) and includes portions of Lamar, Hancock, Pearl River, Stone, Harrison, and Jackson counties. Major tributaries within the Mississippi Coastal Streams watershed include Bayou Casotte, Wolf River, Rotten Bayou, DeLisle Bayou, Bayou La Croix, Bayou Bacon/Jourdan River, Turkey Creek/Bernard Bayou, Biloxi River, and Tuxachanie Creek.

Major rivers carry high sediment loads into the Mississippi Sound. Inland fresh water drainage from these and other smaller rivers create an estuarine environment. Variable salinity levels can affect the productivity and survival of organisms living in the area, as well as economic and recreational activities. Pollution from agriculture, cities, improperly treated sewage, roadways, accidental oil spills, industrial discharges, and other sources also affect the health of the habitats. Graveline Bay is influenced by freshwater flow from several small tributaries. The waters in this area are classified by the State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters (MDEQ 2012) as "shellfish harvesting", "recreation", and "fish and wildlife" (within Graveline Bay proper), and "recreation" and "fish and wildlife" for all other areas in the proposed alternative location. Commercial harvest of oysters is currently restricted in Graveline Bayou and Graveline Bay. None of the waterbodies that drain directly into Graveline Bay are listed as impaired on the State of Mississippi 303(d) list (MDEQ 2014).

Floodplains

The proposed alternative is in FEMA Federal Insurance Rate Maps 28059C0406G, 28059C0314G, and 28059C0405G. A large portion of the area is mapped as Zone VE. Zone VE is defined as Coastal flood zone with velocity hazard. This includes beach areas, open water and most estuarine marsh. Some estuarine marsh, streams, and riparian areas are mapped as Zone AE. Zone AE is defined as "Base Flood Elevations Determined". Upland areas are mostly Zone X. Zone X are defined as "Areas of 0.2% annual change flood; areas of 1% annual chance flood with average depths of less than 1 foot or with drainage areas less than 1 square mile; and areas protected by levees from 1% annual chance flood".

Wetlands

In general, estuarine areas within the proposed alternative are composed of low, mid, and high marsh zones. In the low marsh areas, regularly flooded by tidal activity, the area consists of mesohaline habitat. The intermediate (mid) marsh zone is irregularly flooded by tidal activity and is typically dominated by black needlerush (*Juncus roemerianus*), which can be intermixed with salt grass (*Distichlis spicata*) in oligohaline areas. In higher elevation areas, it is not uncommon to observe numerous species intermixed including salt grass, black needlerush, and salt meadow cordgrass (*Spartina patens*). Fire- suppressed pine savanna and coastal plain small stream forest habitat may be jurisdictional wetlands having prolonged durations of surface water hydrology in a depressional landscape context. Plant communities are discussed in Section 3.3.1.3 (Biological Environment).

Environmental Consequences for WCNH/Birds Proposed Alternative A (Preferred)

Environmental consequences affecting hydrology, water quality, floodplains, and wetlands are discussed below. Table 3.3-4 lists the environmental consequences of each proposed alternative activity to hydrology and water quality.

Hydrology and Water Quality

Acquisition/Preservation: Acquisition and preservation would open new areas to recreational activities including hiking, fishing, bird-watching, and camping. Access via motorized vehicles would be limited. Preservation of lands would have indirect, long-term benefits by preventing development and disturbances, which can reduce surface water runoff and result in long-term water quality benefits to the proposed alternative project area.

<u>Access restriction</u>: Access restriction on Graveline beach would provide short-term benefits to water quality resulting from a decrease in disturbance/equipment use on the beach.

Chemical treatment: Chemical treatment activities would include the use of herbicides. There could be unavoidable spills near the intended application area. However, best practices would be used to prevent any harmful chemicals from entering the environment. Implementation of best practices that the MS TIG would consider, described in the best practices summary below includes development and implementation of a spill prevention and response plan, including conduction daily inspections during chemical treatment to ensure there are no leaks of pesticides or other substances. Personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities. As such this activity, would have short-term, minor, adverse impacts, if any, on water quality. There could be short-term, minor impacts to hydrology as a result of minor rutting/soil disturbance and temporary changes in hydrologic patterns from vehicular transport of personnel to treatment areas.

Mechanical treatment/Prescribed fire: Mechanical treatment would apply to up to 496 acres of fire-suppressed pine savanna and up to 66 acres of coastal plain small stream forest. Prescribed fire would apply on up to 496 acres of fire-suppressed pine savanna. Since large equipment may be needed, soil disturbance, rutting, compaction and any resulting erosion could have a short-term, minor to moderate, adverse impact to water quality. There could be small, temporary changes to stormwater flows and runoff retention patterns due to rutting by equipment and vegetation removal resulting in a short-term, minor to moderate adverse impact to hydrology. There would be short-term, minor to moderate, adverse impacts resulting from mechanical treatment of woody underbrush and construction of fire breaks. There could be small, temporary changes to stormwater flows and runoff retention patterns due to rutting by equipment and vegetation removal. Soft tracked or wide tracked equipment would be used in wet areas to the extent practicable. Alternatively, crews may access the area on foot and remove vegetative material with chainsaws or by hand.

<u>Debris removal</u>: Debris removal could result in limited compaction and soil movement due to the use of equipment, and physical removal of debris. Impacts to water quality would be negligible. There would be no debris removal operation in water or in estuarine marsh.

Road repair/removal and culvert placement: Removal of road beds of up to four acres would require the use of excavation equipment, dump trucks, and other large equipment. Soils adjacent to the road bed could be disturbed or compacted from operations. Erosion control measures would be implemented during construction operations. Roadbed areas would be recontoured and prepared for revegetation. There could be short-term, moderate, adverse impacts to water quality during road bed removal as a result of construction-related sediment movement, and sedimentation of surrounding areas until vegetation is established on the disturbed area. Design of road repair/removal and culvert placement would include, to the extent practicable, efforts to restore historic hydrologic patterns. Road repair/removal and culvert placement could result in long-term, beneficial impacts to local hydrology and stormwater runoff patterns. The activity would result in long-term, water quality and hydrology benefits by restoring the natural hydrologic connection of the area surrounding the road.

Floodplains

Acquisition and preservation of land in perpetuity would prevent land development in floodplains. There would be a long-term benefit to floodplains. Chemical treatment, mechanical treatment and prescribed fire operations would not result in a detectable change to natural and beneficial floodplain values. Road removal/repair would restore natural hydrologic connectivity to areas adjacent to the roadways and would exchange compacted road surface with ground that would eventually be vegetated, providing a long-term benefit to floodplains.

Wetlands

<u>Acquisition and Preservation</u>: There would be a long-term benefit to wetlands from acquisition and preservation. Wet fire-suppressed pine savanna areas that are acquired would not be filled for development.

<u>Access Restriction</u>: Access restriction would occur on the Graveline beach. Barriers would not be placed in wetland areas. There would be no effect to wetlands as a result of this activity.

<u>Chemical treatment</u>: Chemical treatment activities would require the use of herbicides and equipment during applications. Personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities. Only chemicals

approved for use in wetlands would be used. Equipment traffic in wetlands would be avoided to the extent practicable. Best practices would be used during the application of herbicides. Accidental spillage could result in minor, short-term adverse impacts to wetland habitat. However, best practices would be used to prevent any harmful chemicals from entering the environment and for clean up if a spill occurred.

Mechanical Treatment: Mechanical treatment in wetland areas would be done in a manner that would minimize impacts to wetlands to the extent practicable. If mechanical treatment is conducted in wetlands, soft track or wide track equipment would be used to distribute the equipment weight and minimize ground impacts. Alternatively, crews may remove material with chainsaws. If required, a USACE permit would be obtained; likely a Nationwide 27 (Aquatic Habitat Restoration, Establishment, and Enhancement Activities) as well as MDMR Coastal Wetlands Permit (if required). Nationwide 27 allows for mechanized land clearing to remove non-native invasive, exotic or nuisance vegetation and other related activities. If there is clearing within wetlands or stream boundaries, damage to vegetation, soil compaction and any resulting erosion could have a short-term, minor to moderate impacts to wetlands. USACE permit and/or MDMR Coastal Wetlands permit conditions (if required) would be adhered to in all operations.

<u>Prescribed fire:</u> Prescribed fire would apply to up to 496 acres of fire-suppressed pine savanna, a portion of which, are likely wetlands. Intermittent fires were historically a critical perturbation for this habitat. There would be short term minor to moderate impacts to wetlands resulting from mechanical treatment of woody underbrush and construction of fire breaks if the fire breaks are in wetlands or streams. Permit requirements and minimization measures are discussed above in mechanical treatment. There would be long-term beneficial effects to wet fire-suppressed pine savannas including a re-establishment of wetland communities, and increased diversity in flora and faunal populations that colonized the prescribed burn unit.

<u>Debris removal</u>: Debris removal from wetlands would be completed in a manner that would not substantially disturb or redistribute soils including avoidance of equipment in saturated areas and hand removal by field crews. Debris removal could have short-term, minor, adverse impacts to wetlands. Debris removal would have a long-term beneficial impact to wetlands.

Road repair/removal and culvert placement: Removal of road beds of up to 4 acres would require the use of excavation equipment, dump trucks, and other large equipment. Removing the roadbed would allow wetlands to return to a more naturally functioning state. There could be short-term, minor to moderate impacts to surrounding wetlands and streams during road bed removal as a result of increased erosion and sedimentation until vegetation is established. There would be a long-term, wetland benefit from culvert placement if design of the project enhances natural historic hydrologic patterns. Section 404 permits would be obtained as required. All activity would be conducted in compliance with applicable permit conditions. Erosion control and spill prevention measures would be implemented during construction activities.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to water quality and hydrology:

- In the execution of land acquisition and the design of habitat management measures the MS TIG would consider resiliency measures to facilitate habitat migration due to sea level rise (CEQ, 2016).
- Develop and implement an erosion control plan to minimize erosion during and after construction and where possible use vegetative buffers (100 feet or greater), revegetate with native species or annual grasses, and to the extent practicable, conduct work during dry seasons.
- For chemical treatment, personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities.
 Personnel will apply herbicide in accordance with the direction and guidance provided on the appropriate U.S. Environmental Protection Agency (EPA) labels and state statutes during land-based activities.
- Soft track or wide track equipment would be used in wet areas to the extent practicable. Alternatively, crews may remove vegetative material with chainsaws.
- Avoid and minimize, to the maximum extent practicable, placement of dredged or fill material in wetlands and other aquatic resources. Design construction equipment corridors to avoid and minimize impacts to wetlands and other aquatic resources to the maximum extent practicable. If required, a USACE permit and/or MDMR Coastal Wetlands Permit would be obtained; likely a Nationwide 27 (Aquatic Habitat Restoration, Establishment, and Enhancement Activities) as well as MDMR Coastal Wetlands Permit (if required). USACE permit and/or MDMR Coastal Wetlands permit conditions (if required) would be adhered to in all operations.
- Designate a vehicle staging area removed from any natural surface water resource or wetland to perform fueling, maintenance, and storage of construction vehicles and equipment.
 Inspect vehicles and equipment daily prior to leaving the storage area to ensure that no petroleum or oil products are leaking.
- Develop and implement a spill prevention and response plan, including conducting daily inspections of all construction and related equipment to ensure there are no leaks of antifreeze, hydraulic fluid, or other substances and cleaning and sealing all equipment that would be used in the water to rid it of chemical residue.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. If these properties were developed, there would likely be adverse effects on hydrology, water quality, floodplains, and wetlands. Adverse hydrologic affects could include increased runoff rates due to impervious surfaces related to development. Increases in sediment entering waterways could result in adverse effects to water quality. Floodplain and wetland function could be adversely affected by development of parcels proposed for acquisition, preservation and management under proposed Alternative A.The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.2.3 Air Quality and Greenhouse Gas Emissions

Affected Environment

The following section is a discussion of air quality for the proposed alternative project area. EPA has set national ambient air quality standards (NAAQS) for six principal air pollutants (also called criteria pollutants): Ground-Level Ozone (O3), Particulate Matter (PM), Nitrogen Dioxide (NO2), Sulfur Dioxide (SO2), Carbon Monoxide (CO), and Lead (Pb). MDEQ is the state agency responsible for development and maintenance of state specific air emission standards for Mississippi, and monitors all of these pollutants. In Jackson County, the following parameters are monitored: Ozone, Particulate Matter, Nitrogen Oxides, and Sulfur Dioxide. According to MDEQ 2015 Air Quality Data Summary³⁴ the entire state of Mississippi, including Jackson County, is meeting all of the NAAQS.

Environmental Consequences for WCNH/Birds Proposed Alternative A (Preferred)

The environmental consequences for this section is divided into two discussions: 1- environmental consequences from equipment operation/best practices and; 2- environmental consequences resulting from prescribed fire/best practices.

<u>1-Environmental Consequences Resulting from Equipment Operation/Best Practices:</u> The following proposed alternative implementation activities would produce emissions during equipment operation: chemical treatment, mechanical treatment, and road repair/removal and culvert placement. Because these restoration activities would occur seasonally, and would be limited in scope and distribution, the adverse impacts on air quality or to emissions of greenhouse gases would be short-term and minor.

Best Practices

Unavoidable short-term, minor adverse impacts from equipment emissions would be offset through the following best practices to the extent practicable:

- Shut down idling construction equipment, if feasible.
- Encourage the use of the proper size of equipment for the job to maximize energy efficiency.
- Use of ultra-low sulfur diesel fuel in off-road construction equipment with engine horsepower (HP) rating of 60 HP and above.

2- Environmental Consequences Resulting from Prescribed Fire/Best Practices: The use of prescribed fire is included in this project as a restoration activity to provide major long-term benefits for native species habitats, water and soil quality, and nutrient cycling³⁵. However, short-term minor to moderate adverse impacts to air quality and greenhouse gases may occur during the prescribed fire. Smoke emissions are primarily composed of water vapor and carbon dioxide but also contains carbon monoxide, nitrogen oxide, hydrocarbons, particulate matter, and trace minerals. According to the

 $[\]frac{34}{http://www.deq.state.ms.us/mdeq.nsf/pdf/Air} 2015AirQualityDataSummary/\$File/2015\%20Air\%20Quality\%20Data\%20Summary.pdf?OpenElement (MDEQ 2015)$

³⁵ https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/16/stelprdb1046311.pdf (NRCS 2015)

National Coalition of Prescribed Fire Councils Guide to Smoke Management (September 2007 version)³⁶, the primary concerns of smoke as an air pollutant are as follows:

- Carbon Dioxide: The emission factor for carbon dioxide for prescribed burning is 2,000-3,500 pounds/ton (pounds of emissions/ton of organic matter burned)..
- Carbon monoxide: The emission factor for carbon monoxide for prescribed burning is 20-500 pounds/ton. It is classified as a criteria pollutant by EPA. Because of rapid dilution and its instability, carbon monoxide emissions from prescribed burning are not a concern to the general public.
- Water vapor: The emission factor for water vapor for prescribed burning is 50-1500 pounds/ton. The only possible concern about water vapor is visibility reduction in the vicinity of the fire.
- Particulate matter: The emission factor for particulate matter for prescribed fire is 20-180 pounds/ton. Particulates are a criteria pollutant and can impact health and visibility.
 Particulates are presently the major pollutant of concern from prescribed burning. They represent a health risk by inhalation and also reduce visibility.
- Hydrocarbons: The emission factor for hydrocarbons for wildland fire is 10-40 pounds/ton. While hydrocarbons are not a criteria pollutant, they may impact health and visibility and in some cases, may contribute to excessive ozone concentrations.
- Nitrogen oxides: The emission factor for nitrogen oxides for wildland fire is 1-9 pounds/ton. Nitrogen oxides are a criteria pollutant and can impact health and visibility. The low emission factor reduces concern of ambient air quality standards on a local level; however, nitrogen oxides can affect ozone formation.
- Secondary emissions: Secondary emissions are pollutants which are formed in the atmosphere by photochemical transformation of primary emissions. They include oxidants such as ozone which is a criteria pollutant. Specific emission factors from prescribed burning are unknown but are believed to be relatively small.
- Air Toxics: There is an emerging concern about the potential emission of air toxics (acetaldehyde, acrolein; 1,3 butadiene; formaldehyde; and polycyclic organic matter (POM). POM includes eight major categories of compounds including polycyclic aromatic hydrocarbons (PAHs) which include number

Adverse impacts to air quality by controlled burns would be minimized by the frequency and timing of the events; typically, they would be conducted every 1-3 years on managed burn areas per the management plan. Unavoidable short-term minor to moderate adverse impacts from prescribed fire would be offset through the development of a Prescribed Burn Plan, which would include some or all the following Best Smoke Management Practices (BSMPs) and would be part of the management plan. These BSMPs (October 2011) were developed by USDA Forest Service/NRCS³⁷ to mitigate the impacts of smoke to public health (See Section 3.3.1.4.5), public safety and nuisance, and visibility. These six BSMPs have applicability depending on the type of burn, fuel to be burned, and level of effort needed to address air quality concerns. BSMPs are utilized by the individual fire manager and

³⁶http://www.garxfire.com/pdf%20files/The_National_Coalition_of_Prescribed_Fire_Councils_Guide_to_Smoke_Management.pdf (The national Coalition of Prescribed Fire Councils 2007)

³⁷ https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/16/stelprdb1046311.pdf (NRCS 2015)

may be an expectation of a state-wide smoke management program and any applicable conservation plans which are in place for the proposed alternative area (Table 3.3-5).

Table 3.3-5: Summary of Basic Smoke Management Practices

Basic Smoke Management Practice	Benefit achieved with the BSMP	When the BSMP is Applied
Evaluate Smoke Dispersion Conditions	Minimize smoke impacts	Before, During, After
Monitor Effects on Air Quality	Be aware of where the smoke is going and degree it impacts air quality	Before, During, After
Record-Keeping/Maintain a Burn/Smoke Journal	Retain information about the weather, burn and smoke. If air quality problems occur, documentation helps analyze and address air regulatory issues	Before, During, After
Communication- Public Notification	Notify neighbors and those potentially impacted by smoke, especially sensitive receptors	Before, During
Consider Emission Reduction Techniques	Reducing emissions can reduce downwind impacts	Before, During
Share the Airshed Coordination of Area Burning	Coordinate multiple burns in the area to manage exposure of the public to smoke	Before, During, After

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. If these properties were developed, there would likely be adverse impacts to air quality due the potential of development, the additional traffic and other air pollution related to development, and removal of vegetation that benefits air quality. Under the No Action alternative, prescribed fire would not take place as an additional management activity, resulting in no additional short-term, minor to moderate impacts to air quality from burning. This short term impact however would be offset by the potential for development with its resultant potential for long-term impacts. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.3 Biological Environment

<u>Introduction to Affected Environment (Biological Environment)</u>: Biological environment resources discussed in the section include habitats, wildlife, and protected species. PDARP/PEIS sections 6.4.1.5.2 and 6.4.10.1.2 are incorporated by reference here. The affected environment for the proposed alternative biological environment is described in respective sections below.

<u>Programmatic Review of Environmental Consequences (Biological Environment)</u>: Sections 6.4.1.5.2 and 6.4.10.1.2 of the PDARP describe the impacts to Biological Resources for the restoration approaches relevant restoration approaches and are incorporated by reference and briefly described here.

<u>PDARP/PEIS</u> consequences related to land management plans: Specific restoration activities identified as part of land management plans could result in short-term, minor to moderate adverse effects on conservation areas. Consequences reviewed in the PDARP/PEIS are incorporated here and summarized.

<u>PDARP/PEIS</u> consequences related to Invasive species: Activities that may occur on conserved lands may result in introduction of invasive species. Use of best practices would help prevent the introduction of invasive species. Implementation of land management plans, located within or near restoration activities, could result in disturbed, removed, or altered habitats, which could cause minor to moderate, short- and long-term adverse effects on species that use those habitats for forage or nesting purposes.

<u>PDARP/PEIS</u> consequences related to public access: Land acquisition could permit public access for recreational use. This public use, depending on management stipulations, could result in long-term, minor to moderate adverse effects on area species through increased human presence and activity on acquired habitats.

<u>PDARP/PEIS</u> consequences related to habitat: Conservation of habitat through fee title acquisition and improved management could have a long-term benefit to any habitat on the property acquired or protected. Conservation would also allow for upland migration of beach, wetland, or other habitats as the sea level rises and could limit development encroachment.

<u>PDARP/PEIS</u> consequences related to habitat and resource benefits: Conservation of habitat through fee title acquisition could have a long-term benefit to fish, birds, and terrestrial wildlife through the protection of coastal, riparian, or terrestrial habitat. These habitats can be important for food supply and various life stages of some species. Benefits of the proposed restoration approach include conservation of bird nesting and foraging habitat that would increase bird health and reproduction by preventing habitat loss through land conversion.

<u>PDARP/PEIS</u> consequences related to access restriction: Restrictions on seasonal or overall human use that could result from changes in land management would reduce habitat degradation. Improvements in habitat associated with this approach may draw additional visitors to the area, resulting in potential indirect adverse impacts from human presence. Human disturbance can lead to failure of nests, increased egg and chick predation, or even total colony abandonment.

<u>PDARP consequences related to vegetation management</u>: Managing vegetation is a common restoration technique to enhance habitat for specific bird species. Reducing vegetation on beaches, for example, can provide nesting and foraging habitat for birds such as such as snowy plover, least tern, black skimmer, and American oystercatcher. Conversely, adding vegetation can provide habitat for other bird species such as wading birds and brown pelicans. Common vegetation management methods include mechanical treatments, application of pesticides or herbicides, and biological control to manage plant species.

Environmental consequences for the proposed alternative are within the general range impacts as described in the PDARP/PEIS with some variances related to specific actions. Table 3.3-6 summarizes the environmental consequences to the biological environment that would result from the proposed alternative. These impacts to these resources is discussed below.

Table 3.3-6: Environmental Consequences to the Biological Environment due to the Proposed WCHN/Bird Alternative A (Preferred)³⁸

WCHN/Bird Alternative A (Preferred) ³⁸								
	Acquisition/Preservation	Access restriction	Chemical treatment	Mechanical treatment	Prescribed fire	Debris removal	Road removal/repair and culvert placement	
Habitats					•			
Beach								
Adverse Impact Duration		short-term	short-term					
Adverse Impact Intensity		minor	minor					
Beneficial Impact Duration	long-term	long-term	short-term			short-term		
Beech-Magnolia Forest	Ü	Ü						
Adverse Impact Duration			short term			short term		
Adverse Impact Intensity			minor			minor		
Beneficial Impact Duration	long-term		long-term			short-term		
Fire Suppressed Pine	8		2418 44111					
Savanna								
Adverse Impact Duration			short-term	short-term	short-term	short-term	short-term	
- carrier confusion				minor to	minor to			
Adverse Impact Intensity			minor	moderate	moderate	minor	moderate	
Beneficial Impact Duration	long-term		long-term	long-term	long-term	short-term	long-term	
Coastal Plain Small Stream			long term	rong term	Tong term	SHOTE COTTE	long term	
	l							
Adverse Impact Duration			short-term	short-term	short-term	short-term		
				minor to	minor to	_		
Adverse Impact Intensity			minor	moderate	moderate	minor		
Beneficial Impact Duration	long-term		long-term	long-term	long-term	short-term		
Estuarine Marsh								
Adverse Impact Duration								
Adverse Impact Intensity								
Beneficial Impact Duration	long-term							
Invasive Species								
Adverse Impact Duration								
Adverse Impact Intensity								
Beneficial Impact Duration			long-term	long-term	long-term			
Wildlife Species (including l	birds)							
Adverse Impact Duration			short-term	short-term	short-term	short-term	short-term	
-				minor to	minor to			
Adverse Impact Intensity			minor	moderate	moderate	minor	minor	
Beneficial Impact Duration	long-term	long-term	long-term	long-term	long-term	short-term	long-term	

As appropriate in a tiered analysis, the evaluation of the proposed alternative focuses on the specific resources with a potential to be affected. Marine and estuarine fauna are not expected to be affected by the proposed alternative as there is no in-water work. To avoid redundant or unnecessary information, marine and estuarine fauna are evaluated summarily here.

_

³⁸ Protected species are not included in this table and are addressed in Section (3.3.1.3.2).

Marine and Estuarine Fauna (Submerged Aquatic Vegetation, Nearshore Benthic Invertebrates, Marine Mammals, Essential Fish Habitat): There would be no in-water work. Estuarine marsh would be acquired and preserved, but there are no management activities planned in this habitat in the proposed alternative project area. Acquisition and preservation of habitat would prevent development and preclude habitat removal or stresses that could result from shoreline development.

For the biological environment, the following resources are further analyzed in this section:

- Habitats
- Protected Species
- Migratory Birds
- Wildlife

3.3.1.3.1 Habitats

The section includes habitats found in the proposed alternative area and the environmental impacts from restoration activities that would be implemented in those habitats.

Affected Environment

Section 3.5 of the PDARP/PEIS provides a discussion of habitats of the northern Gulf of Mexico; Section 3.7.4 covers invasive species. This section covers habitats in the proposed alternative project area. The Mississippi Sound extends along the southern coasts of Mississippi and Alabama. The Mississippi Sound is separated from the Gulf of Mexico by several narrow barrier islands and sand bars (including Cat Island, Ship Island, Horn Island, and Petit Bois Island), which provide dynamic and diverse habitats especially for over 300 species of migratory or permanent resident bird species (USACE 2009). Along the Mississippi Sound, there are numerous coastal bays including St. Louis Bay, Biloxi Bay, Back Bay of Biloxi, Pascagoula Bay, Graveline Bay and Grand Bay. The Mississippi Sound is shallow with water depths generally not exceeding 20 ft. Water is exchanged with the Gulf of Mexico through the openings between the barrier islands. This partially protected nature and the influx of riverine freshwater create a salinity gradient within the Sound (Priddy et al. 1955). This delicate mix of fresh and salt water provides a suitable habitat for oysters, shrimp, and other fisheries. Christmas and Waller (1973) reported 138 fish species in 98 genera and 52 families taken from areas across the Mississippi Sound. Vittor and Associates (1982) identified over 437 taxa of macrofauna from the sound with densities varying from approximately 1,200 to 38,900 individuals per square yard.

• Graveline Bay and waterways represent one of only a few relatively undisturbed estuarine bays and small tidal creeks in Mississippi. It is located between Grand Bay to the east and Biloxi Bay to the west. The area supports salt marsh, brackish marsh, and several degraded oyster beds (which are intended to be restored under a *DWH* Early Restoration Project). This shallow, coastal bay/marsh estuarine system receives only local freshwater runoff and consists largely of black needle rush dominated marsh along its entire length. Smooth cordgrass occurs largely as narrow (1 to 3 m) bands along the waterways. Subtidal ecological communities/habitats include muddy sand embayment, small tidal creeks and mollusk reefs. Intertidal ecological communities/habitats include sand beach, mesohaline marsh, and oligohaline marsh. Much of the marsh area is already part of the MDMR CP Program.

• Within the proposed alternative area, coastal wetland and nearshore habitats include estuarine marsh, beach, beech-magnolia forest, coastal plain small stream forest, fire suppressed pine savanna, and open water including tidal creeks and bayous (Figure 3.3-2).

Estuarine Marsh: Approximately 1,218 acres of estuarine marsh exists within the proposed alternative area, 636 acres are in private ownership. Estuarine marsh consists largely of black needle rush dominated marsh along its entire length. Smooth cordgrass (*Spartina alterniflora*) occurs largely as narrow (1-3 m) bands along the creeks and bayous.

Beach: Approximately 6 acres of natural beaches of the Graveline area are located directly adjacent to the Mississippi Sound. Sandy material is distributed and deposited via longshore currents. The beach habitat also exhibits soft, easily erodible marsh terraces directly in front of the beach deposits. Currently, the beach is primarily unvegetated with common reed as a dominant on the northern interface between the beach and marsh. The beach is used as nesting habitat by the diamondback terrapin (*Malaclemys terrapin*) on a regular basis as well as several solitary nesting shorebird species. There has been one recorded atypical use of this site by a nesting loggerhead turtle (*Caretta caretta*).




Figure 3.3-2: Habitats in the Proposed Land Acquisition and Management Project Area

Beech-Magnolia Forest: Approximately 115 acres of beech magnolia habitat exists within the proposed alternative area including the public owned parcels in the CP and the acquisition parcels in and adjacent to the CP. The Beech-Magnolia forest community occurs in transitional areas from upland longleaf pine high relief areas to stream bottoms. American beech (*Fagus grandifolia*) and magnolias (*Magnolia grandiflora*) are the dominant trees of the canopy, but the forested community can be very diverse with several species of hardwoods (e.g. oaks) and pines also occurring. This community represents the climax community of this ecoregion (MMNS 2015).

Coastal Plain Small Stream Forest: Coastal plain small stream swamp forests are alluvial swamps along small drainages. In the proposed alternative area, there is a total of 66 acres of this habitat.

Their floodplains are often protected by a dense mat of interwoven tree roots, traversed by braided streams. Sweetbay (*Magnolia virginiana*) and water tupelo (*Nyssa aquatic*) are the most common trees. Red maple (*Acer rubrum*), and water oak (*Quercus nigra*) are also common (MMNS 2015). The understory of these habitats remains open with regular fires, but quickly becomes overgrown by rapidly growing shrubs such as swamp titi, buckwheat tree, and large gallberry in the absence of fire.

Fire Suppressed Pine Savanna: Approximately 496 acres of fire suppressed pine savanna exists in the proposed alternative project area. In Mississippi, the historical longleaf pine forest extended from the wetlands of the coast to the mixed pine- hardwood forests of central Mississippi and from the border of Alabama to the Loess Hills. Natural fires maintained forests and savannas of massive, well-spaced longleaf pine trees. Combustible leaf litter and grassy understory carried natural wildfires through the longleaf region. Sampling of virgin forests over a century ago indicated that tree densities averaged about 100 per acre, or 400 square feet per tree. In the absence of frequent burns, other pines, hardwood trees and shrubs rapidly move into these longleaf pine savannas. In addition, many of the areas were planted in faster growing species such as slash pine (*Pinus elliotii*). In just a few years, the midcanopy and shrub layers of this community can become thick and impenetrable, eliminating natural regeneration of the shade-intolerant longleaf seedlings. If left unaltered, this community succeeds to an oak-hickory-pine community on drier sites and to beech-magnolia in mesic areas (MMNS 2015). If managed, wetter pine savannas can have a diverse community of carnivorous plants including pitcher plants (*Sarracenia alata*), sundews (*Drosera spp.*) and in ponding areas, bladderwort (*Utricularia* sp.)

Open Water: Approximately 485 acres of open water exists in the proposed alternative project area. Graveline Bay is tidally influenced, with wide ranging salinities levels. Graveline Bay supports subtidal and intertidal oysters and is a popular fishing area.

Invasive Species EO 13112 applies to all federal agencies whose actions may affect the status of invasive species, requires agencies to identify such actions, and to the extent practicable and permitted by law, requires agencies to 1) take actions specified in the Order to address the problem consistent with their authorities and budgetary resources and 2) not authorize, fund, or carry out actions that they believe are likely to cause or promote the introduction or spread of invasive species in the United States or elsewhere unless pursuant to guidelines that it has prescribed, the agency has determined and made public its determination that the benefits of such actions clearly outweigh the potential harm caused by invasive species; and that all feasible and prudent measures to minimize risk of harm will be taken in conjunction with the actions. The proposed alternative habitat management is primarily invasive species management with restoration actions and measures including chemical treatment, mechanical treatment and prescribed fire. Best practices that would be used to control or eliminate invasive species are discussed in the environmental consequences section below.

Environmental Consequences for WCNH/Birds Proposed Alternative A (Preferred)
A summary of proposed restoration activity and adverse and beneficial impacts are listed in Table 3.3-6 and discussed in this section.

Estuarine Marsh: Acquisition and preservation of habitat would prevent development and preclude habitat removal or stresses that could result from site development. There would be a long-term benefit to acquiring estuarine marsh.

Beach: Acquisition/preservation, access restriction, chemical treatment for common reed and debris removal would be conducted on this habitat. Acquisition and preservation of habitat would prevent development. There would be short-term, minor, adverse impacts resulting from the installation of barriers on the beach. Access restriction would allow the beach to recover from current use impacts and would protect shorebird habitat, providing long-term benefits. For chemical treatment of common reed management, there could be minor impacts to adjacent vegetation from the misapplication in the intended area and incidental spillage of chemicals. Personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities. Chemical treatment could result in short-term, minor impacts to habitat. Chemical treatment would have a short-term beneficial impact by preventing the spread of common reed. Debris removal would have a short-term beneficial effect on beach habitat.

Beech-Magnolia Forest: Acquisition/preservation, chemical treatment, and debris removal would be conducted on this habitat. Acquisition and preservation of habitat would prevent development, habitat loss and fragmentation. This would result in a long-term benefit to the habitat. Equipment use during chemical treatment and debris removal could result in short-term minor impacts to habitat. The restoration activities would have a beneficial impact by preventing the spread of invasive species and restoring native species composition. This would result in a long-term benefit to the habitat. Debris removal would have a short-term benefit to Beech-magnolia forest.

Fire-Suppressed Pine Savanna: Acquisition/preservation, chemical treatment, mechanical treatment, prescribed fire, debris removal, and road removal/repair and culvert placement would be conducted on this habitat. Acquisition and preservation of habitat would prevent development, habitat loss and habitat fragmentation. Acquisition and preservation provide a long-term benefit to the habitat.

Chemical treatment could result in short-term, minor impacts from equipment use and incidental spillage of herbicide, both localized to small areas. Eradication and control of invasive species using chemical treatment would result in a long-term benefit to this habitat.

Mechanical treatment/Prescribed fire: Activities include but would not limited to use of brush-hog, and use of chainsaws. In addition, use of gyro-tracks and in some cases bobcats or bulldozers to lay down or remove vegetation could be used as a stand-alone treatment or in combination/preparation for prescribed fire. The preferred prescribed fire regime would be completed, ideally, on a two-year rotation, with 50% of the prescribed burns occurring during the growing season. Weather conditions, seasonal wetness, availability of trained staff, invasive species present and other factors are considerations in maintaining the fire frequency; 1-3 years. These activities would largely be applied in areas that were colonized by woody invasive and understory shrubs such as gallberry (*Ilex glabra*), privet, saw palmetto, Chinese tallow, and other species. Impacts to soils and wetlands were discussed in previous section. These could result in short-term, minor to moderate, adverse impacts, to existing habitats depending on the size of the operation. There would be long-term benefits to fire suppressed savannas from mechanical clearing alone or in combination with prescribed fire by creating conditions that would result in the re-establishment of diverse plant communities.

Debris Removal: There could be short-term, minor, adverse impacts from equipment related to debris removal in fire-suppressed pine savannas. There would be a short-term beneficial affect from debris removal.

Road Removal/Repair and Culvert Placement: Equipment used for road removal/repair and culvert placement would result in short-term, moderate impacts to habitat. The equipment would cause disturbance to vegetation and soils adjacent to existing roads, which would temporarily impact habitats. The impacts would be adjacent to up to 4 acres of roadway constituting a moderate impact. The restoration activities would have a long-term, beneficial impact which include restoring historic hydrologic conditions beneficial to fire-suppressed pine savannas.

Coastal Plain Small Stream Forest: Acquisition/preservation, debris removal, and road removal/repair and culvert placement would be conducted on this habitat. The adverse and beneficial impacts described in fire-suppressed pine savanna for these activities would apply here.

Open Water: There would be no work in open water. Therefore, no adverse or beneficial impacts would result.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to reduce the spread of invasive species:

- Prior to bringing any equipment (including personal gear, machinery, vehicles, or vessels) to the work site, inspect each item for mud or soil, seeds, and vegetation. If present, clean the equipment, vehicles, or personal gear until they are free from mud, soil, seeds, and vegetation.
- Inspect the equipment, vehicles, and personal gear each time they are being prepared to go to a site or prior to transferring between sites to avoid spreading exotic, nuisance species.

No Action

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. If these properties were developed, there would likely be adverse impacts to habitats including habitat removal and/or fragmentation. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.3.2 Protected Species

Affected Environment

Section 3.6 of the PDARP/PEIS discusses biota of the northern Gulf of Mexico. This section covers endangered species in the proposed alternative area. The U.S. Fish and Wildlife Service (USFWS) lists species as threatened or endangered when they meet criteria detailed under the Endangered Species Act (ESA) of 1973, as amended (16 U.S.C. §1531 et seq.). Additionally, Mississippi Wildlife Fisheries and Parks (MWFP) and NOAA National Marine Fisheries Service (NMFS) identify and list protected species. Section 7(a) (2) of the ESA requires that each federal agency ensure that any action

authorized, funded, or carried out by the agency is not likely to jeopardize the continued existence of any endangered or threatened species or result in the destruction or adverse modification of Critical Habitat of those species. When the action of a federal agency may affect a protected species or its Critical Habitat, that agency is required to consult with either the NMFS or the USFWS, depending upon the protected species that may be affected. ESA Section 7 coordination is underway and the appropriate recommendations would be incorporated into the proposed alternative. Compliance with the Migratory Bird Treaty Act and Bald and Golden Eagle Protection Act are also discussed in this section.

Relevant federally protected species that are known to occur or could occur in Jackson County and that could occur in or near the proposed alternative project area or could pass through the proposed alternative project area are listed in Table 3.3-7. A brief discussion of the state imperiled diamond back terrapin is also provided in the environmental consequences.

Table 3.3-7: Federally threatened, endangered, and proposed species in the project area for

WCNH/Birds Proposed Alternative A (Preferred)

Common Name	Scientific Name	Federal Status	Habitat
Birds			
Piping Plover	Charadrius melodus	Threatened	Beaches and mudflats in southeastern coastal areas. Critical Habitat, MS-15, exists in Jackson County but is not in the proposed alternative area.
Red Knot	Calidris canutus rufa	Threatened	Marine intertidal habitats including inlets, estuaries, and bays feeding in mud and sand flats on beaches and barrier islands
Mississippi Sandhill Crane	Grus canadensis pulla	Endangered	Open wetland habitats surrounded by shrubs or trees. Critical Habitat has been established on and adjacent to the Mississippi Sandhill Crane National Wildlife Refuge (USFWS 2013).
Fishes			
Gulf Sturgeon	Acipenser oxyrinchus desotoi	Threatened	Migrates from large freshwater coastal rivers to brackish and marine coastal bays and estuaries. Graveline beach is adjacent to Critical Habitat Unit 8, but there would be no in-water work in Critical Habitat.
Mammals			
West Indian Manatee	Trichechus manatus	Endangered	Fresh and salt water in large coastal rivers, bays, bayous and estuaries
Reptiles			
Hawksbill Sea Turtle	Eretmochelys imbricate	Endangered	Coral reefs, open ocean, bays, estuaries
Leatherback Sea Turtle	Dermochelys coriacea	Endangered	Open ocean, coastal waters
Kemp's ridley Sea Turtle	Lepidochelys kempii	Endangered	Nearshore and inshore coastal waters, often in salt marshes; neritic zones with muddy or sandy substrate (NOAA Fisheries 2014b)
Green Sea Turtle	Chelonia mydas	Threatened	Shallow coastal waters with SAVs and algae, nests on open beaches
Loggerhead Sea Turtle	Caretta	Threatened	Open ocean; also inshore areas, bays, salt marshes, ship channels and mouths of large rivers
Alabama Red-belly Turtle	Pseudemys alabamensis	Endangered	Fresh and brackish habitats, river banks, submerged and emergent aquatic vegetation; upland habitat for nesting (MDWFP 2001; USFWS 2013)

Common Name	Scientific Name	Federal Status	Habitat		
Black Pinesnake	Pituophis melanoleucus lodingi	Threatened	uplands with well-drained sandy soils in areas of longleaf pine and hardwood tree species (USFWS 2013).		
Gopher Tortoise	Gopherus polyphemus	Threatened	Well-drained, sandy soils, which allow easy burrowing; an abundance of diverse herbaceous ground cover; and an open canopy and sparse shrub cover, which allows sunlight to reach the ground floor (USFWS 2013).		
Plants					
Louisiana Quillwort	Isoetes		Mineral soil, usually light gray in color, in bottomlands that are periodically washed free of leaves and debris. Streams along which quillworts grow may have flow year around.		

Birds

Mississippi Sandhill Crane: The Mississippi Sandhill crane utilizes open wetland habitats surrounded by shrubs or trees. Critical Habitat has been designated on and adjacent to the Mississippi Sandhill Crane National Wildlife Refuge (USFWS 2013).

Piping Plover (*Charadrius melodus*): The piping plover does not nest in Mississippi; however, this species uses Gulf Coast beaches and barrier islands for wintering (MDWFP 2001). Plovers use sparsely vegetated sand beaches, mudflats, and salt marshes for roosting and foraging.

Red Knot (*Calidris canutus rufa*): In coastal Mississippi, the red knot is mainly a migratory species that uses coastal beaches and marine intertidal areas as stopover feeding locations or staging areas on the way to and from their wintering grounds in South America and breeding areas in the Arctic. Foraging on ocean beaches, mud and sand flats, and salt marshes occurs from March to April during the northward spring migration and September and October during the southward autumn migration (Niles et al. 2007; USFWS 2013). Red knots have been observed wintering on the Gulf Coast and are observed from October to March (USFWS 2013). The nonbreeding diet of this species includes marine invertebrates such as snails, crustaceans, and small mollusks including the coquina clam (*Donax variabilis*), which is common on Gulf coast beaches, and the dwarf surf clam (*Mulinia lateralis*) (Niles et al. 2007; USFWS 2013). Roosting and resting habitat includes areas above the high tide line such as reefs and high sand flats (USFWS 2013).

Fishes

Gulf sturgeon (*Acipenser oxyrinchus desotoi*): This anadromous species migrates from coastal bays and estuaries to large coastal rivers in the spring for spawning and then returns to brackish and marine environments from October through March for foraging. The riverine spawning habitats for Gulf sturgeon in the State of Mississippi include the Mississippi, Pearl, and Pascagoula rivers (Ross et al. 2009; MDWFP 2001) but not the Biloxi and Tchoutacabouffa rivers (USFWS, GSMFC, and NMFS 1995; NMFS and USFWS 2009). The marine wintering areas where individuals have been observed are nearshore and barrier island habitats from the Pearl River east to the barrier islands (Ross et al. 2009). Winter habitat is mainly around Cat, Ship, Horn, and Petit Bois islands with nearshore observations likely due to migratory movements to and from these offshore islands (Rogillio et al. 2007; Ross et al. 2009). The coastal Mississippi Sound waters of the State of Mississippi are designated as Critical Habitat.

Gulf Sturgeon Designated Critical Habitat: The proposed alternative area is adjacent to Gulf sturgeon Critical Habitat at the mouth of Graveline Bayou/along Graveline beach (Unit 8-Lake

Ponchartrain-Mississippi Sound). There would be no in-water work during the implementation of the proposed alternative.

Mammals

West Indian Manatee (*Trichechus manatus*): This species uses both fresh and saltwater habitats such as coastal rivers, bays, bayous, and estuaries. The manatee is an occasional visitor to Mississippi's coasts, although migration into the area is poorly understood. After wintering in Florida, and perhaps Mexico, manatees migrate northward during spring, including to Mississippi and Alabama waters, although these migrations are not well understood (Fertl et al. 2005). Manatees frequently seek out freshwater sources such as rivers and river mouths and have been known to be found near estuaries (Fertl et al. 2005). SAVs are the typical manatee forage material; however, manatees can also consume other aquatic vegetation, algae, and terrestrial vegetation (Fertl et al. 2005). There are no proposed restoration activities in open water.

Reptiles

Hawksbill Sea Turtle (*Eretmochelys imbricata*): Although this species uses various habitats such as the open ocean, bays, and estuaries throughout different life stages, it is mainly associated with coral reefs. This species nests in Florida from April to November (NOAA Fisheries 2014a). It likely does not nest in Mississippi and observations are rare in the state (MDWFP 2001; NOAA Fisheries 2014a). The main dietary items of this species are sponges and other invertebrates (NOAA Fisheries 2014a). There are no proposed restoration activities in open water.

Leatherback Sea Turtle (*Dermochelys coriacea*): This species mainly inhabits the offshore open ocean; however, it does use nearshore coastal waters during nesting or feeding. Nesting for this species occurs in Florida from April through November. Their main forage item is jellyfish. This species migrates long distances from nesting to feeding areas. While not common, there have been sporadic observations of leatherback sea turtles in Mississippi waters (MDWFP 2001). There are no proposed restoration activities in open water.

Kemp's Ridley Sea Turtle (*Lepidochelys kempii*): Typical habitat for this species includes nearshore and inshore coastal waters and often salt marshes and neritic zones with muddy or sandy substrate (NOAA Fisheries 2013b). This species has been observed in nearshore waters of the Mississippi Sound during migration and foraging and has been accidentally caught by shore-based fishermen (MDWFP 2001; Shaver and Rubio 2008). Females typically nest from May through July (NOAA Fisheries 2014b). Males potentially use Gulf of Mexico habitats all year and females presumably use the Mississippi Sound and barrier island habitats for foraging when not nesting (NOAA Fisheries 2014b). Kemp's ridley sea turtles do not nest in Mississippi (MDWFP 2001). There are no proposed restoration activities in open water.

Green Sea Turtle (*Chelonia mydas*): This species typically prefers shallow coastal waters with SAVs and algae for foraging and nests on open beaches (NOAA Fisheries 2015). Nesting typically does not occur on mainland beaches and there is likely no Mississippi nesting at all (MDWFP 2001; NOAA Fisheries 2015). This species migrates long distances in the open ocean from nesting to feeding areas. Observations of this species in Mississippi are rare (MDWFP 2001). There are no proposed restoration activities in open water.

Loggerhead Sea Turtle (*Caretta caretta*): Loggerhead habitat for foraging and migration includes open ocean, inshore areas, bays, salt marshes, ship channels, and mouths of large rivers. This sea

turtle feeds on mollusks, fish, crustaceans, and other marine organisms. This species typically nests at night from late April through September (NOAA Fisheries 2014c). Although loggerheads occasionally use barrier islands for nesting, mainland nesting is rare (MDWFP 2001). Preferences for nesting beaches include high-energy coarse-grained beaches adjacent to the ocean that are narrow and steeply sloped (NOAA Fisheries 2014c). This species has been observed in nearshore waters of the Mississippi Sound during migration and foraging and has been accidentally caught by shore-based fishermen (MDWFP 2001). There was one atypical nesting event on Graveline beach. There are no proposed restoration activities in open water.

Alabama Red-Belly Turtle (*Pseudemys alabamensis*): The habitat of the Alabama red-belly turtle includes fresh and brackish habitats, river banks, submerged and emergent aquatic vegetation, and upland habitat for nesting (MDWFP 2001; USFWS 2013). This species is mainly a freshwater species associated with river and stream channels and associated wetlands. Nesting occurs from mid-May to mid-July (MDWFP 2001).

Mississippi Diamondback Terrapin (*Malaclemys terrapin pileata*): The Mississippi diamondback terrapin (*Malaclemys terrapin pileata*) utilizes pocket beaches adjacent to marsh for nesting habitat (Frey 2014). Diamondback terrapins have a diet of fish, snails, worms, clams, crabs and marsh plants and live in brackish water habitats such as estuaries and tidal marshes, preferring marshes with nearby channels. Juveniles may spend first few years under mats of flotsam or vegetation (Ernst et al. 1994). Clutches are laid from April to August. The Mississippi diamondback terrapin is ranked by the Mississippi Department of Wildlife, Fisheries, and Parks (MDWFP) as S2: Imperiled in Mississippi. (Mississippi Natural Heritage Program 2015). Restoration activities will not be done on pocket beaches.

Black Pine Snake (*Pituophis melanoleucus lodingi*): Suitable habitat includes open canopy longleaf pine forest with herbaceous ground cover and well-drained sandy soils and, less so, hardwood forests (USFWS 2010). Much of the habitat in the proposed alternative area is not suitable because of dense canopy cover or due to existing disturbance.

Gopher Tortoise (*Gopherus polyphemus*): The Gopher Tortoise uses well-drained to excessively well-drained upland soils. Tortoises require soils that are sandy enough to permit construction of burrows and open canopies that allow sufficient herbaceous plant growth and sunny areas in which to nest. In Mississippi, these areas often support a mixture of longleaf pine and scrub oaks.

Plants

Louisiana Quillwort (*Isoetes louisianensis*): The Louisiana Quillwort has been observed in 10 counties in 174 streams within 17 watersheds (USFWS 2012a) throughout the State of Mississippi with the largest colony found in the DeSoto National Forest (USFWS 2012a). This species is found in all three coastal Mississippi counties (MDWFP 2001; USFWS 2012a) although none have been found near the proposed alternative area (MDWFP 2001). In coastal Mississippi, Louisiana Quillwort habitat includes perennial streams and banks in bottomland hardwood habitats likely with bald cypress and possibly the presence of stream macrophytes such as *Sparganium* spp. and *Orontium* spp. (USFWS 2012a). Earlier sources indicate that suitable habitat for this species consists of sand or gravel bars located in intermittent streams and associated riparian areas (MDWFP 2001). Louisiana Quillworts are sensitive to changes in hydrology, sedimentation, and alterations to the surrounding overstory (USFWS 2012a).

PDARP programmatic ESA consultations were developed with the National Marine Fisheries Services (NMFS, 2016) and the U.S. Fish and Wildlife Service (USFWS, 2016). Potential impacts to threatened or endangered species and their Critical Habitat are presented in Table 3.3-8. The MS TIG has begun coordination under the programmatic ESA consultations. The proposed alternative area in the southeast is adjacent to the Mississippi sound which is designated Critical Habitat for Gulf sturgeon. None of the restoration activities would be completed in open water. Thus, there would be no effect as a result of any restoration activity to in water species (and associated Critical Habitat), including Gulf sturgeon, West Indian manatee, and sea turtles; for this reason, they are not included in the environmental consequences discussion in Table 3.3-8.

Table 3.3-8: Protected Species Environmental Consequences for the WCNH/Birds Proposed Alternative

7.1	<u>A</u>					
Species /Critical Habitat	Applicable Habitats	Restoration Activities for Applicable Habitats	Potential Impacts to Species/Critical Habitat			
Alabama Red-Belly Turtle (Pseudemys alabamensis)	Estuarine Marsh	Acquisition/PreservationDebris Removal	The restoration activities for this habitat type would have no adverse effect on this species.			
Piping plover (Charadrius melodus) and red knot (Calidris canutus rufa)	Beach	Access Restriction	Since the only restoration activity on beach type habitat is access restriction, no adverse impacts are anticipated.			
Black pine snake (Pituophis melanoleucus lodingi)	Fire-suppressed pine savanna	Acquisition/Preservation Chemical treatment Mechanical treatment Prescribed fire Debris Removal Road removal/repair and culvert placement	It is not likely that this exact habitat exists in the proposed alternative area. However, if the habitat does exist prescribed fire and mechanical treatment of upland areas may affect species habitat. Surveys should be conducted in areas where the species is likely to occur. Survey results would be considered in the design of the management and or restoration measures to either avoid or minimize impacts to the species.			
Gopher tortoise (Gopherus polyphemus)	 Fire-suppressed pine savanna Beech Magnolia Forest 	Acquisition/Preservation Chemical treatment Mechanical treatment of undesirable vegetation Prescribed fire Debris Removal Road removal/repair and culvert placement	Prescribed fire and mechanical treatment of upland areas may affect species habitat. Surveys would be conducted in areas where the gopher tortoise is likely to occur. Survey results would be considered in the design of the management and or restoration measures to either avoid or minimize impacts to the species.			
Louisiana quillwort (Isoetes louisianensis)	Coastal Plain Small Stream Forest	 Acquisition/Preservation Chemical treatment Mechanical treatment of undesirable vegetation Debris Removal Road removal/repair and culvert placement 	Chemical treatment, mechanical treatment, debris removal, and road removal/repair could result in an impact to vegetation. Restoration activity areas would be surveyed for the species and if found avoided in the implementation of restoration measures and activities.			
Mississippi Sandhill Crane (<i>Grus</i> canadensis pulla)	 Coastal Plain Small Stream Forest Estuarine 	 Acquisition/Preservation Chemical treatment Mechanical treatment of undesirable vegetation Debris Removal 	Chemical treatment, mechanical treatment, debris removal, and road removal/repair could result in a noise impact causing the species to temporarily vacate the area. The species			

Species /Critical Habitat	Applicable Habitats	Restoration Activities for Applicable Habitats	Potential Impacts to Species/Critical Habitat
	Marsh	 Road removal/repair and culvert placement 	could return after restoration activities have ceased.
Mississippi diamondback terrapin (Malaclemys terrapin pileata)	Beach	Access Restriction	Since the only restoration activity on beach type habitat is access restriction, no adverse impacts are anticipated.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The MS TIG would continue to consult with the appropriate regulatory agency to further avoid or minimize impacts to these species in the planning site-specific restoration activities and management measures. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to protected species:

Piping Plover and Red Knot

- Provide all individuals working on a restoration activities associated with the proposed alternative with information in support of general awareness of piping plover or red knot presence and means to avoid birds and their critical or otherwise important habitats.
- Minimize vegetation planting in preferred habitats and avoid removal of wrack year-around along the shoreline.
- During recreational use, enforce leash or "no pet" policies in critical or important habitats.

Gopher Tortoise

• If suitable habitat is present, coordinate with the local USFWS Ecological Services Field Office to discuss the need for surveys to identify any gopher tortoise burrows and to develop conservation measures to avoid or minimize impacts. Measures could include establishing a protective buffer (size determined by USFWS and the state trust resource agency) if burrows are within the proposed alternative area and cannot be avoided, implementing standard procedures to relocate the tortoise within the proposed alternative site but away from the areas of restoration.

Protected Plants

- If suitable habitat is present, coordinate with the local USFWS Ecological Services Field Office to discuss the need for surveys to identify protected plants and to develop conservation measures to avoid or minimize impacts.
- Enhance and protect plants on site and in adjacent habitats to the maximum extent possible.

Protected Species

• Provide all individuals working on restoration activities associated with the proposed alternative with information in support of general awareness of and means to avoid impacts to protected species and their habitats present at the specific project site.

ESA Section 7 coordination is underway and the appropriate recommendations would be incorporated into the proposed alternative. Because no effects to manatee are expected, the Implementing Trustees determined that no take of manatee under ESA or MMPA would occur.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which undertaken. Without NRDA funding restoration actions were for acquisition preservation/management, these properties may be developed in the foreseeable future. If these properties were developed, there would likely be adverse impacts to habitat that could be utilized by Habitats that protected species could use would not be protected from development under the No Action alternative and would not be managed for increased habitat benefits. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.3.3 Migratory Birds

Affected Environment

Migratory bird species groups that could occur in the alternative project area include wading birds, shorebirds, seabirds, raptors, goatsuckers, waterfowl, doves and pigeons, and rails and coots (see Table 3.3-9).

Table 3.3-9: Species Groups Present in the Project Area for WCHN/Birds Proposed Alternative A

SPECIES	BEHAVIOR	SPECIES/HABITAT IMPACTS
Wading birds (herons, egrets, ibises)	Foraging, feeding, resting, roosting, nesting	Wading birds primarily forage and feed at the water's edge. As such, they may be impacted locally and temporarily by the proposed alternative. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. These birds primarily nest and roost in trees or shrubs (e.g., pines, Baccharis), and could utilize areas that will be managed by mechanical treatment and prescribed fire.
Shorebirds (plovers, oystercatchers, stilts, sandpipers)	Foraging, feeding, resting, roosting, nesting	Shorebirds forage, feed, rest, and roost in the action area. As such, they may be impacted locally and temporarily by the proposed alternative. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. In the proposed alternative area, these birds would primarily nest on beaches. Access restriction would include placement of barriers at the western edge of the beach in order to reduce nest disturbance. Placement of barriers would be done so as not to impact nesting. Chemical treatment and/or mechanical treatment of common reed could also be conducted in the area; care would be take to complete activities away from nesting birds or when nesting is not occurring on the beach.
Seabirds (terns, gulls, skimmers, double-crested cormorant, American white pelican, brown pelican)	Foraging, feeding, resting, roosting,	Seabirds forage, feed, rest, and roost in the action area. Terns and skimmers could utilize the beach habitat in the proposed alternative area. As such, they may be impacted locally and temporarily by the proposed alternative. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. Chemical treatment and/or mechanical treatment of common reed could also be conducted in the area; care would be take to complete activities away from nesting birds or when nesting is not occurring on the beach.
Raptors (osprey, hawks, eagles, owls)	Foraging, feeding, resting, roosting, nesting	Raptors forage, feed, rest and nest in the action area. As such, they may be impacted locally and temporarily by the proposed alternative.

SPECIES	BEHAVIOR	SPECIES/HABITAT IMPACTS
		It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. There is an existing Osprey nest in the northeastern part of the proposed alternative area. Work in the area could include debris removal. Debris removal would be completed so as not to disturb osprey nesting. Ospreys are relatively tolerant of human activity in the vicinity of their nests.
Goatsuckers	Foraging, feeding, resting, roosting, nesting	Goatsuckers forage, feed, rest, and roost in the action area. However, they are nocturnal/crepuscular and therefore not active during the project work period. They nest in thickets and woodlands.
Waterfowl (ducks, loons, and grebes)	Foraging, feeding, resting, roosting, nesting	Waterfowl forage, feed, rest, and roost in the action area. As such, they may be impacted locally and temporarily by the proposed alternative. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. These birds primarily roost and nest in low vegetation. There would be no restoration activities in open water or estuarine marsh.
Doves and pigeons	Foraging, feeding, resting, roosting	Doves and pigeons could forage, feed, rest, and roost in the action area. However, they are unlikely to utilize habitat in the firesuppressed pine savanna or the estuarine zone for nesting; no impacts to nesting are anticipated.
Rails and coots	Foraging, feeding, resting, roosting, nesting	Rails and coots forage, feed, rest, and roost in the action area. As such, they may be impacted locally and temporarily by the proposed alternative. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting if disturbed by the proposed alternative. These birds primarily roost and nest in marshes, which are within the action area. There would be no restoration activities where these species nest.

Migratory Bird Treaty Act

The Migratory Bird Treaty Act of 1918 (MBTA) implements various treaties and conventions among the United States, Canada, Japan, Mexico, and the former Soviet Union for the protection of migratory birds. Under MBTA, unless permitted by regulations, it is unlawful to pursue; hunt; take; capture or kill; attempt to take, capture, or kill; possess; offer to sell or sell; barter; purchase; deliver; or cause to be shipped, exported, imported, transported, carried, or received any migratory bird, part, nest, egg, or product, manufactured or not. USFWS regulations broadly define "take" under MBTA to mean "pursue, hunt, shoot, wound, kill, trap, capture, or collect, or attempt to pursue, hunt, shoot, wound, kill, trap, capture, or collect".

Bald and Golden Eagle Protection Act

The Bald and Golden Eagle Protection Act (BGEPA) of 1940 (16 U.S.C. 668-668c) prohibits anyone, without a permit issued by the Secretary of the Interior, from "taking" bald eagles, including their parts, nests, or eggs. BGEPA provides criminal penalties for persons who "take, possess, sell, purchase, barter, offer to sell, purchase or barter, transport, export or import, at any time or any manner, any bald eagle [or any golden eagle], alive or dead, or any part, nest, or egg thereof." Golden eagles are not present along the Gulf Coast.

Environmental Consequence for WCNH/Birds Proposed Alternative A (Preferred)

Migratory birds could use areas at and around the proposed alternative project area for foraging, feeding, resting, and nesting. Nesting species include raptors (forest edge near marsh), wading birds (pine trees/shrubs adjacent to estuarine marsh), marsh birds (estuarine marsh), waterfowl (estuarine marsh), and shorebirds (beach); Table 3.3-10. For all planned restoration activities, precommencement nesting surveys for migratory birds and raptors within the restoration activity area

would be conducted and if evidence of nesting is found, CP resource managers would coordinate with the USFWS to develop and implement appropriate conservation measures, such as those described below. Due to the implementation of best management practices no "take" of nesting birds is anticipated. There are no golden eagles in the proposed alternative footprint. Raptor nest surveys would be completed within the restoration activity area where raptor nesting habitat exists. If evidence of nesting is found, CP resource managers coordinate with the USFWS to develop and implement appropriate conservation measures, therefore no impacts to golden or bald eagles are anticipated. Potential adverse effects to birds include elevated noise levels due to the use of mechanical equipment for vegetation clearing, and from noise and smoke during prescribed burning. These species are mobile and would likely exit the area during management activities. Foraging and resting birds may temporarily be displaced during management activities. Bird roosting would not be affected because management activities would occur during daylight hours. Therefore, impacts are expected to be short-term, localized, and minor.

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to migratory bird species including bald eagles:

Migratory Birds

- Use care to avoid birds when operating machinery or vehicles near birds.
- Avoid working in migratory bird nesting habitats during breeding, nesting, and fledging
 (approximately mid-February through late August). If proposed alternative activities must
 occur during this timeframe and breeding, nesting, or fledging birds are present, contact the
 state trust resource agency to obtain the most recent guidance to protect nesting birds or
 rookeries, and their recommendations will be implemented.
- Conservation areas may already be marked to protect bird nesting areas. Stay out of existing marked areas.
- If vegetation clearing is necessary, clear vegetation outside the migratory bird nesting season (approximately mid-February through late August) or have a qualified biologist inspect for active nests. If no active nests are found, vegetation may be removed. If active nests are found, vegetation may be removed after the nest successfully fledges.

Bald Eagles

- If bald eagle breeding or nesting behaviors are observed or a nest is discovered or known, have all activities avoid the nest by a minimum of 660 feet. If the nest is protected by a vegetated buffer where there is *no* line of sight to the nest, then the minimum avoidance distance is 330 feet. Maintain this avoidance distance from the onset of breeding/courtship behaviors until any eggs have hatched and eaglets have fledged (approximately 6 months).
- If a similar activity (such as driving on a roadway) is closer than 660 feet to a nest, maintain a distance buffer as close to the nest as the existing tolerated activity. If a vegetated buffer is present and there is no line of sight to the nest and a similar activity is closer than 330 feet to a nest, then maintain a distance buffer as close to the nest as the existing tolerated activity.

• In some instances, activities conducted within 660 feet of a nest may result in disturbance, particularly for the eagles occupying the Mississippi barrier islands. If an activity appears to cause initial disturbance, stop the activity and move all individuals and equipment away until the eagles are no longer displaying disturbance behaviors. Contact the USFWS's Migratory Bird Permit Office to determine how to avoid impacts or if a permit may be needed.

The MS TIG has begun coordination and review of the proposed alternative for impacts to bald eagles and migratory birds in accordance with the Bald and Golden Eagle Protection Act (BGEPA) of 1940 (16 U.S.C. 668-668c) and the Migratory Bird Treaty Act (MBTA) of 1918 (16 U.S.C. 703–712) to ensure appropriate conservation measures and best practices would be incorporated into the project.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. Although migratory birds and bald/golden eagles would still be protected under the No Action alternative, if these properties were developed, there would likelybe impacts tohabitats that these species use. It is likely that these impacts would be minimized with the use of required Best Management Practices. Noise disturbance would increase if development takes place. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.3.2 Wildlife

Affected Environment

Section 3.6 of the PDARP/PEIS discusses the biota of the northern Gulf of Mexico. For the proposed alternative project area, faunal species include those associated with natural estuarine marsh, transition areas and uplands adjacent to estuarine marsh, and beach habitats. These include various species of mammals, birds, fish, reptiles, infauna, epifauna, and other aquatic invertebrates. The mixing of freshwater from tributaries with saline water from the Mississippi Sound allows for a range of fish species in the waters of Graveline Bay/Bayou including redfish (*Sciaenops ocellatus*), freshwater catfish (order Siluriformes), flounder (Paralichthys spp.), speckled trout (Cynoscion nebulosus), white trout (Cynoscion arenarius), southern kingfish (Menticirrhus americanus), sheepshead (Archosargus probatocephalus), and black drum (Pogonias cromis), as well as crab and shrimp species. The estuarine emergent wetland habitat supports an array of neonate and juvenile fish and aquatic invertebrates. Other fish and marine mammals such as Atlantic bottlenose dolphins (Tursiops truncatus) could also occur in the Mississippi Sound adjacent to the proposed alternative area. The upland areas support a range of species including, but not limited to, white-tailed deer (Odocoileus virginianus), wild turkey (Meleagris gallopavo), squirrels (Sciurus spp.), and rabbit (Oryctolagus cuniculus). The MDMR plans for the CP within the proposed alternative project area include protecting habitats and the ecological integrity of the tidal marsh and adjacent uplands in order to benefit wildlife and habitat.

Environmental Consequences for WCHN/Birds Proposed Alternative A

Acquisition/Preservation: Prevention of development of habitats would be a long-term, benefit to wildlife species that currently inhabit or transiently utilize the preserved habitats.

<u>Access Restriction</u>: Access restriction would provide protection of shorebird habitat and would provide a long-term benefit to shorebirds, wading birds, pelicans, seagulls, and other species that routinely use the beach for loafing, foraging and nesting.

<u>Chemical Treatment</u>: Chemical treatment would result in a short-term, minor impact to wildlife species in and near treatment areas due to equipment noise and exposure to chemicals. There would be a long-term benefit to habitats and wildlife that utilizes the habitat.

Mechanical Treatment and Prescribed Fire: Mechanical treatment and prescribed fire would be the most intrusive to wildlife, however, these techniques would be applied to areas that have dense woody shrub layers which preclude utilization by several bird and mammal species. There would be a short term, minor to moderate impact to species in the area during mechanical treatment and prescribed fire. Many species would leave the area during the operations. Mechanically treated and/or prescribed fire areas would become open habitat and be colonized with native pine savanna species over several seasons. Once restored, these communities are one of the most diverse habitats and would result in increased diversity in insect, bird, and small mammal populations. There would be a long-term benefit to wildlife resulting from mechanical treatment and/or prescribed fire.

<u>Debris Removal</u>: Debris removal could result in short-term, minor impacts from equipment noise or disturbance during removal operations. There would be short-term benefits as a result of debris removal.

<u>Road Removal/Repair and Culvert Placement</u>: Road removal/repair and culvert placement would result in short-term, minor impacts to wildlife from equipment noise or disturbance during removal operations. Removing roadways would provide a long-term benefit by increasing habitat connectivity.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. If these properties were developed, there would likely be impacts tohabitats that wildlife species use causing disturbances in all life stages of certain wildlife. Human disturbance, such as noise would likely increase with development and could cause adverse impacts to wildlife. Wildlife habitat would not be enhanced under the No Action like it would in proposed Alternative A. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.4 Socioeconomic Resources

<u>Introduction to Affected Environment (Socioeconomic Resources)</u>: The section provides a discussion of socioeconomic resources and environmental justice, tourism and recreational use, cultural

resources, land and marine management, and public health and safety. PDARP/PEIS Section 3.2 is incorporated by reference here.

<u>Programmatic Review of Environmental Consequences (Socioeconomic Resources)</u>: Sections 6.4.1.5.3 and 6.4.1.10.1.3 of the PDARP describe the impacts to Socioeconomic Resources for the relevant restoration approaches and are incorporated by reference and briefly described here.

PDARP/PEIS consequences related to economic effects: Acquisition and preservation could have long-term, minor to moderate adverse economic effects if acquisition prevents or limits development. Acquisition could permanently limit the amount and type of development permitted, and the management and intensity of use on these properties would likely change. Ownership changes and/or permitted uses could affect property taxes and have broader regional economic impacts. Land acquisition could have a minor to moderate impact on socioeconomic resources due to changes in visitor spending and tax impacts. The transfer of fee title to lands are transactions negotiated or arranged between willing parties and, as such, are not expected to give rise to adverse socioeconomic impacts to those who choose to engage in such transactions.

PDARP/PEIS consequences related to recreation and tourism: The acquisition of lands to protect habitat could result in impacts to recreation and tourism opportunities depending on site-specific land management practices applied. Closures, such as fencing or other mechanisms to protect nest sites, could result in short-term (seasonal) prohibitions on public access. Restrictions on public access in areas where public access had previously been allowed could reduce recreational opportunities. Over the long term, these techniques could result in healthy populations and provide wildlife enthusiasts with increased wildlife viewing opportunities. Conservation or acquisition of natural land resources can have indirect benefits on fish and wildlife habitat, potentially resulting in increased fishing and hunting opportunities. Seasonal or permanent employment could increase in order to provide labor for the installation, maintenance, and implementation of management projects such as hunting or trapping. Minor, short-term adverse impacts could result due to restoration activities. However, improvements in habitat associated with this approach may draw additional visitors to the area with associated visitor spending, increasing sales and tax receipts on retail purchases.

<u>PDARP/PEIS</u> consequences related to cultural resources: Creating, enhancing, or restoring bird nesting habitat may result in minor (temporary disturbance) to moderate (disturbance without loss of cultural information) impacts on cultural and historic resources depending on the scale of the action and site-specific characteristics. Discovery or recovery of cultural or historic resources would allow their future protection.

Table 3.3-10 summarizes the socioeconomic resources' environmental consequences associated with the proposed alternative which are discussed in detail in this section.

Table 3.3-10: Proposed Alternative Impacts to Socioeconomic Resources

Table 3.3-10: Proposed Alternative Impacts to Socioeconomic Resources							
	Acquisition/Preservation	Access restriction	Chemical treatment	Mechanical treatment	Prescribed fire	Debris removal	Road removal/repair and culvert placement
Socioeconomic Resources and	d Environme	ntal Justice					
Adverse Impact Duration	short to long-term						
Adverse Impact Intensity	minor to moderate						
Beneficial Impact Duration							
Tourism and Recreational Us	se						
Adverse Impact Duration			short-term	short-term	short-term		
Adverse Impact Intensity			minor	minor	minor		
Beneficial Impact Duration	long-term						
Cultural Resources							
Adverse Impact Duration	TBD	TBD	TBD	TBD	TBD	TBD	TBD
Adverse Impact Intensity	TBD	TBD	TBD	TBD	TBD	TBD	TBD
Beneficial Impact Duration	TBD	TBD	TBD	TBD	TBD	TBD	TBD
Land and Marine Manageme	ent						
Adverse Impact Duration	long-term						
Adverse Impact Intensity Beneficial Impact Duration	minor to moderate						
Public Health and Safety, inc	luding flood	and shoreling	e protection				
Adverse Impact Duration			short-term		short-term		
Adverse Impact Intensity			minor		minor		
Beneficial Impact Duration	long-term						

As appropriate in a tiered analysis, the evaluation the proposed alternative focuses on the specific resources with a potential to be affected. Infrastructure, fisheries and aquaculture, marine transportation, aesthetics and visual resources would have negligible to minor adverse effects or would provide benefits. To avoid redundant or unnecessary information, a summary of environmental consequences for these resources is provided here.

Infrastructure: Infrastructure on the site includes access roads for logging/timber management, gas pipelines and utility corridors. There could be short-term, minor impacts to gas pipelines or utility corridors from activities associated with mechanical treatment and prescribed fire. Care would be take to identify utility corridors as part of project planning and prior to implementation or restoration measures. Portions of poorly maintained roads within fire-suppressed pine savanna habitat would be removed as a result of implementing the proposed alternative. These are largely private logging roads. The impacts resulting from these actions road repair/removal and culvert placement are covered in the site-specific analysis for physical and biological resources, but the proposed activities would not affect public infrastructure.

Fisheries and Aquaculture: There would be no activities in open water or estuarine marsh. Acquisition and restoration measures could benefit oyster reefs in Graveline Bay and Graveline

Bayou by a net reduction in sediment movement resulting from preservation and restoration versus a development/build out scenario of lands proposed for acquisition.

Marine Transportation: There would be no restoration activity that would occur in open water; the proposed alternative would not have an impact on marine transportation.

Aesthetics and Visual Resources: Prescribed fire would result in a change in viewshed. There may be temporary short-term, minor impacts as a result due to presence of smoke. The land may look scorched after a prescribed burn until vegetation regrows. Depending on weather conditions, burn units can revegetate ("green up") within days to weeks. Revegetation after burning would result in a viewshed of natural vegetation with increased diversity of flowering plants and fauna. Removal of unmaintained roads and debris would enhance the aesthetic character of the land for the public that utilizes the area.

For socioeconomic and environmental justice, the following resources are further analyzed in this section:

- Socioeconomics and Environmental Justice
- Tourism and Recreational Use
- Cultural Resources
- Land and Marine Management
- Public Health and Safety

3.3.1.4.1 Socioeconomic Resources and Environmental Justice

Affected Environment

PDARP Section 3.2 discusses socioeconomic resources of the Gulf Coast and is incorporated by reference here. The affected environment for the proposed alternative includes the population of Census Tract 409 and 411, specifically the residents close to the Graveline Bay. The population of Jackson County was 139,668 in 2010 and accounted for 4.7% of the state's total population, while Census Tract 409 (population 11,240 in 2010) accounted for 8% of the county population, and Census Tract 411 (population 6,700 in 2010) accounted for 5% of the county population (Table 3.3-12). In 2010, median household income in Jackson County was \$49,145, which was 25% higher than the median household income in the State of Mississippi (\$39,464). Median household income of Census Tract 409 in 2014 was \$60,212, which is 23% higher than that of the county and 53% higher than the median household income of the state. Median household income of Census Tract 411 in 2014 was \$41,985, which is 15% lower than that of the county and 6% higher than the median household income of the state. (U.S. Census Bureau, 2010-2014 American Community Survey 5-Year Estimates)

Table 3.3-11: Population data (http://www.census.gov/2010census/popmap/)

Торіс	Mississippi		Jackson County		Census 7		Census Tract 411	
2010 Total Population	2,967,297		139,668		11,240		6,700	
White alone	1,754,684	59%	100,735	72%	9,163	82%	3,761	56%
Black or African American alone	1,098,385	37%	30,034	22%	1,321	12%	2,566	38%

Торіс	Mississippi		Jackson County		Census Tract 409		Census Tract 411	
Asian alone	25,742	<1%	3023	2.2%	251	2.2%	62	<1%
American Indian and Alaska Native alone	15,030	<1%	565	<1%	56	<1%	41	<1%
Native Hawaiian and Other Pacific Islander alone	1,187	<1%	79	<1%	11	<1%	3	<1%
Some Other Race alone	38,162	1.3%	2610	1.9%	155	1.3%	108	1.6%
Two or More Races	34,107	1.1%	2622	1.9%	283	2.5%	159	2.3%

Acquisition and preservation of property in fee and an in-perpetuity set-aside would permanently restrict development on acquired parcels. The change in ownership would affect property taxes paid to local governments and could result in a broader regional economic impact resulting from changes in visitor spending in the area. There could be minor increases in spending resulting from recreational access to the proposed alternative project area as it increases in size and would also be expected to enhance opportunities to hike, or view wildlife in the area. Land acquisition could have a minor to moderate impact on socioeconomic resources due to changes in visitor spending and tax impacts. The transfer of fee title to lands would be transactions negotiated or arranged between willing parties and, as such, are not expected to give rise to adverse socioeconomic impacts to those who choose to engage in such transactions. Executive Order 12898 directs federal agencies to make achieving environmental justice part of its mission by identifying and addressing, as appropriate, disproportionately high adverse human health or environmental effects of its activities on minority and low-income populations. There would be no disproportionate impacts on minority, low-income, or underserved populations from the implementation of proposed Alternative A.

No Action

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. If these properties were developed, there would likely be increased property taxes paid to local governments. There would be no benefits from additional recreational visitor spending that could result from implementation of proposed Alternative A. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.4.2 Tourism and Recreational Use

Affected Environment

The public has access to the Graveline Bay CP for recreational activities including boating, kayaking, fishing, bird-watching and pedestrian access, though hiking opportunities are limited. The Octavia Street boat ramp affords public access to CP properties as well as Graveline Bayou and its tributaries. Fishing, crabbing, and waterfowl hunting are also done in the area.

Acquisition and preservation would result in a long-term benefit to tourism and recreational opportunities and would open an additional 1,410 acres, that were previously inaccessible, to recreational activities and would enhance the limited hiking opportunities that are currently available on the existing Graveline Bay CP. Implementation of the proposed alternative would also expand areas for fishing, bird-watching, and camping. There would be long-term benefits that would result from the implementation of proposed Alternative A. There could be a short-term, minor, adverse impacts to recreation to prevent public exposure to smoke during prescribed fire. There could be minor, short-term, adverse impacts to recreation due to restricted access during mechanical or chemical treatment operations. The public would be notified and access would be curtailed during short seasonal windows.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. The No Action alternative would not increase tourism or recreation in the area that could be expected from proposed Alternative A. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.4.3 Cultural Resources

Affected Environment

Cultural resources include historic properties listed in, or eligible for listing in the National Register of Historic Places (36 C.F.R. §60[a-d]). The National Historic Preservation Act of 1966 (NHPA), as amended and recodified (54 U.S.C. § 300308), defines an historic property as "any prehistoric or historic district, site, building, structure, or object included on, or eligible for inclusion on the National Register [of Historic Places]." Under the statute and implementing regulations, historic properties include significant traditional religious and cultural properties important to Indian tribes. Historic properties include built resources (bridges, buildings, piers, etc.), archaeological sites, and Traditional Cultural Properties, which are significant for their association with practices or beliefs of a living community that are both fundamental to that community's history and a piece of the community's cultural identity. Although often associated with Native American traditions, such properties also may be important for their significance to ethnic groups or communities. Historic properties also include submerged resources.

This proposed alternative is currently being reviewed under Section 106 of the NHPA to identify any historic properties located within the proposed alternative area and to evaluate whether the proposed alternative would affect any historic properties. The MS TIG is currently conducting a literature review of the proposed alternative component areas. Previously recorded archaeological sites, shipwrecks, historical standing structures, National Register of Historic Places (NRHP) properties, National Register Districts and National Historic Landmarks are being reviewed. The preliminary review of the previously recorded archaeological sites using MDAH records revealed archaeological sites located within the vicinity of the proposed alternative component areas. The types of sites include shell middens and charted shipwrecks.

The National Historic Preservation Act of 1966 (NHPA) charges the federal government with protecting the cultural heritage and resources of the nation. This proposed alternative would be implemented in accordance with all applicable laws and regulations concerning the protection of cultural and historic resources. Cultural and historic resources would be considered when preparing site-specific restoration measures and management actions. Where there is a likelihood disturbance of cultural resources, CP resource managers would conduct appropriate surveys to inform the methods and location of restoration and management actions. For site-specific restoration measures and management actions, environmental compliance would be conducted by evaluating each restoration measure/management action proposed to be conducted on the parcel(s) against the environmental threshold criteria evaluated under this programmatic analysis. Restoration measures/management actions would be designed to avoid cultural resources to the extent practicable. Graveline CP resource managers would work with the Mississippi State Historic Preservation Office and the DOI to determine compliance measures if resources are likely in the area or encountered during implementation.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. Cultural resources would still be protected under the No Action. Development of the area could result in the adverse impacts to cultural resources. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.4.4 Land and Marine Management


Affected Resources

Land and marine management consideration for the Graveline Bay CP include Coastal Zone Management Act consideration, CP Planning initiatives and local land use planning. Governing the nature of land use development of the project component areas is the 1972 Coastal Zone Management Act (CZMA), which provides for management of the nation's coastal resources and balances economic development with environmental conservation. The overall program objectives of CZMA remain balanced to "preserve, protect, develop, and where possible, to restore or enhance the resources of the nation's coastal zone." The water bottoms are considered state-owned and part of the Public Trust Tidelands.

The Graveline Bay Preserve is designated as a CP in the Mississippi CPs Program. It contains 2,339-acres and is bounded by Graveline Bay and Bayou. MDMR manages the area as a CP for conservation purposes to protect ecological integrity of tidal marsh and surrounding areas (MDMR 2015a).

According to the Future Land Use Map for Jackson County (Neel-Schaffer 2009) the future land use surrounding Graveline bay is General Agriculture to the east and northeast, Single Family Residential to the south, and Residential Estate to the west and northwest. A Land Development Suitability Model for the Graveline area developed by MDMR was utilized in the development of the Gautier

Comprehensive Plan. It shows that most of the areas listed in this RP/EA as beach, estuarine marsh, open water, and most of the coastal plains small stream forest, as being "Water" which is not suitable for development. Most of the beech-magnolia forest and fire suppressed pine savanna are Levels 6, 7, and 8, with 8 being the most suitable for development.



Source: Mississippi Department of Marine Resources

Figure 3.3-3: MDMR Land Development Suitability Model

The City of Oceans Springs is to the west of the alternative project area and the City of Gautier is to the east. The 2010 Ocean Springs Comprehensive Plan shows Graveline Bay and Bayou and areas to the south, west, and northeast as Southeast Growth Area. These are areas which might be appropriate for annexation. The Comprehensive plan describes this growth area as:

...much of the Southeast Growth Area contains wetlands, and the area was flooded extensively during Hurricane Katrina. However, as the eastward expansion of population from Ocean Springs continues, especially as families seek more affordable housing (Planning Works, 2010).

The Ocean Springs School District recently constructed a \$37 M High School near Graveline Bayou. It opened in 2012 and was named the third best school in the state in 2016. The district is widely recognized as one of the highest quality school districts on the Gulf Coast and it is an attractor for residential development and economic development in the Ocean Springs. The new high school is expected to increase development in the area (Planning Works, 2010). The only planned acquisition in the Ocean Springs City limits is to the south of Graveline Bay (Graveline beach). To the north of Graveline Bay and in the City of Ocean Springs, the planned acquisitions are primarily fire-suppressed pine flatwoods.

The acquisition and management of up to 1,410 acres of land in the City of Ocean Springs and City of Gautier new growth areas could require zoning change or variance to designate areas as conservation lands. Acquisition and restoration would affect planned land use by removing the land from residential development. The proposed action is consistent with CP planning initiatives. There would be a long-term, minor to moderate, adverse effect to land and marine management depending on the number of willing sellers and the size of parcels acquired and preserved.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future; however, development would likely have no effect on land and marine management, as existing developments would be completed and would be consistent with existing land use plans. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.1.4.5 Public Health and Safety

Affected Resources

The proposed alternative area consists of Graveline Bay, Graveline Bayou, and surrounding uplands. The surrounding communities that use the area for recreation make up the public health and safety affected resource.

Most of the proposed alternative area is a floodplain. A large portion of the area is mapped as Zone VE. This includes beach areas, open water and mostly estuarine marsh. Zone VE is defined as Coastal flood zone with velocity hazard. Some estuarine marsh, streams, and riparian areas are mapped as Zone AE. Zone AE is defined as "Base Flood Elevations Determined". Upland areas are mostly Zone X. Zone X are defined as "Areas of 0.2% annual change flood; areas of 1% annual chance flood with average depths of less than 1 foot or with drainage areas less than 1 square mile; and areas protected by levees from 1% annual chance flood". Proposed alternative activities would not result in a detectable change to natural and beneficial floodplain values. Restored hydrology from road removal/repair and culvert placement would enhance floodplain functions. The floodplain in acquired and managed parcels would be maintained for flood storage capacity and would preclude residential development and flood risk. Prescribed fire and chemical treatment would also expose the public to smoke and potentially chemicals, respectively.

Environmental Consequences for WCNH/Birds Proposed Alternative A (Preferred)

There would be a short-term, minor adverse impacts to public health and safety. Exposure to smoke during prescribed burns would adversely impact public health, but these impacts are expected to be minor since prescribed burns are typical in this region and short term. Burn plans that include public notification of burns and controlled access into the site during burns would be developed to minimize the risk and potential exposure of the public to smoke. Fire breaks would restrict fire to designated areas and crews will be on site to ensure that fire does not jump the fire breaks. Safety plans would be part of the controlled burn plans.

Chemical treatment would require use of herbicide that could be hazardous if spilled or handled improperly. Personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities. Most of the applications would be in remote areas where there is limited public access.

The proposed alternative area is designated as floodplain. Preventing development in the floodplain/the transition of native habitats to new impervious surface provides a flood risk/public safety benefit. The proposed alternative would have a beneficial effect to the surrounding communities. It would promote healthy lifestyles by allowing recreational use on previously private parcels of land.

No Action

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Without NRDA funding for acquisition and preservation/management, these properties may be developed in the foreseeable future. If these properties were developed, there would likely be no impacts to public health and safety since local building codes and ordinances would be followed. The No Action Alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.3.2 Site-Specific NEPA Review for WCNH/Birds Proposed Alternative A (Preferred)

Section 3.3.1 is a discussion of environmental consequences analysis for proposed Alternative A for WCNH/Birds Restoration Type at a programmatic level. The exact parcels and associated restoration measures and management activities on those parcels are not known at this time. The environmental consequences are based on the range of restoration measures and management activities contemplated on parcels in the proposed alternative project area. The programmatic analysis provides maximum impacts to each of the resource categories based on the MS TIG's knowledge of the proposed alternative project area and the restoration activities and management measures likely needed to restore the project area. The MS TIG is proposing the selection of Alternative A (Preferred). Section 3.1.2 also presents a process that the MS TIG would follow to complete the requirements of NEPA and other environmental statutes as site-specific restoration measures and management activities are planned for Alternative A, if selected.

3.4 Grand Bay Land Acquisition and Habitat Management-Background and Project Description

The proposed Grand Bay Land Acquisition and Habitat Management project includes acquiring privately owned inholdings within the boundaries of the Grand Bay National Wildlife Refuge (NWR), Grand Bay National Estuarine Research Reserve (NERR), and the Grand Bay Savanna CP (Figure 3.4-1). Public and private lands within these boundaries total 28,262 acres. The U.S. Fish and

Wildlife Service (USFWS) manages the NWR³⁹, the Mississippi Department of Marine Resources manages lands on the NERR⁴⁰ and the CP⁴¹. The project location consists of parcels adjacent to and near Grand Bay in Jackson County, Mississippi. The project is located in Jackson County, Mississippi in the boundaries of Grand Bay NWR, NERR and Grand Bay Savanna CP (Figure 3.4-1). The proposed project alternatives consider a number of measures:

- Alternative B acquisition of up to 8,000 acres of land from willing sellers at appraised value in the NWR, NERR and CP boundaries
- Alternative C habitat management on up to 17,500 acres of current public lands within the NWR, NERR and CP boundaries
- Alternative D a combination of both acquisition (up to 8,000 acres) and habitat management (up to 17,500 acres) on both current public lands and acquired parcels in the NWR, NERR and CP boundaries

The proposed Grand Bay Land Acquisition and Habitat Management project has several objectives including: acquisition of properties to protect habitat; contiguous ownership of large tracts for connectivity and to facilitate large-scale, well-established habitat management techniques; and restoration of the structure and function of target habitats within the project boundary (Figure 3.4-4; Table 3.4-13). These actions help restore injuries to wetlands, coastal, and nearshore habitats in Mississippi as well as bird species injured by the Spill.

Habitat that could be acquired includes a diverse array of nearshore coastal and wetland habitats. Grand Bay coastal wetland and nearshore habitats include coastal marsh, beach, freshwater marsh, pine savannas and flatwoods, forested freshwater scrub-shrub, and open water including tidal creeks and bayous (Figure 3.4-5; Table 3.4-13). Habitat in the project area is utilized for foraging, nesting and/or loafing by bird species that were injured in the Spill. Restoration activities and management measures conducted under this project would provide benefits to wading bird species injured by the *DWH* oil spill.

A total of 448 acres of developed land is also present within the proposed project area. Residential and commercial development has been proceeding rapidly in the coastal portion of Jackson County, Mississippi, converting forest plantations and farm fields into developed lots with houses, businesses, and institutions (USFWS, 2008). Publicly-owned developed land within the project area consists of 390 acres, comprised of roads, the I-10 rest area, I-10 weigh station, and Grand Bay NERR visitor center. Privately-owned developed land within the proposed project area consists of 58 acres comprised mostly of single family residences, associated outbuildings, agricultural buildings and a commercial development.

³⁹ Land Protection Plan and Final Environmental Assessment for the Expansion of Grand Bay National Wildlife Refuge (USFWS 2012) and Grand Bay National Wildlife Refuge Comprehensive Conservation Plan (USFWS 2008)

⁴⁰ Grand Bay National Estuarine Research Reserve Final Environmental Impact Statement/Reserve Management Plan (MDMR 1998)

³ Mississippi Gulf Ecological Management Sites (MDMR 2016)




Figure 3.4-1: The Grand Bay Land Acquisition and Management Project Area for Proposed Alternatives

Table 3.4-1: Habitats and Ownership within the proposed project area

Habitat	Publicly Owned (acres)	Privately Owned (acres)	Total Acreage of Habitat
Forested Freshwater Scrub- Shrub	1,895	1,416	3,311
Coastal Marsh	7,003	2,077	9,080
Savannas and Flatwoods	2,741	3,535	6,276
Freshwater Marsh	730	1,207	1,937
Beach	21	0	21
Open Water	6,443	746	7,189
Other (Roadways, development, etc.)	390	58	448
Total	19,223	9,039	28,262

Section 5.5.2.2 of the PDARP/PEIS describes seven restoration approaches for the WCNH Restoration Type. Section 5.5.12.2 describes eight restoration approaches for the Birds Restoration Type. The restoration approaches proposed by the MS TIG that address the goals and objectives for this project include:

- Protect and conserve marine, coastal, estuarine and riparian habitats
- Restore and conserve bird nesting and foraging habitat

Restoration measures and benefits would include acquisition to reduce the threat of further development and to provide for large-scale management efforts, habitat enhancement, decreased habitat fragmentation and increased habitat connectivity to other large conservation parcels in the area. Appropriate management practices for the landscape, such as large-scale fire management, is less effective in landscapes where publicly and privately owned parcels are interspersed since management cannot continue from public to private properties without management agreements.

The MS TIG is proposing to allocate \$6 M toward this project.⁴²

Restoration Measures-Methodology and Timing

The proposed alternatives include management of habitats within the project boundary that are currently in public ownership and in newly acquired parcels (See Figure 3.4-4). The Implementing Trustee would begin negotiations with willing sellers (e.g., title surveys, appraisals, etc.) after RP/EA approval. Additional data collection on target habitats needed to facilitate restoration and management (e.g., habitat inventories, identification of appropriate restoration measures and management activities, etc.) would also be conducted following approval of the project. Restoration measures and management activities would be implemented on a site-specific basis and may vary across the project area depending on the current condition of habitats. Habitat restoration measures and management activities could include chemical treatment, mechanical treatment, and prescribed fire, described below. Proposed restoration measures and management activities by habitat type are summarized in Table 3.4-2.

Table 3.4-2: Restoration Measures and Management Activities by Habitat

Habitat	Total Publicly owned	Total Privately Owned Parcels	Total Acres of Habitat	Acquisition/Preservation	Chemical Treatment	Mechanical Treatment	Prescribed Fire
Developed	390	58	448				
Forested Freshwater Scrub-Shrub	1,895	1,416	3,311	X	X	X	
Coastal Marsh	7,003	2,077	9,080	X	X	X	
Savannas and Flatwoods	2,741	3,535	6,276	X	X	X	X
Freshwater Marsh	730	1,207	1,937	X	X	X	
Beach	21	0	21	X			
Water	6,443	746	7,189				

⁴² The project budget of \$6 M would not complete the acquisition of the entire 8,000 acres and/or the 17,500 acres of management. There are other *DWH* funded projects which propose acquisition and habitat management in the Grand Bay proposed project alternative area and there could be additional *DWH* funds for these activities in Grand Bay in the future.

Acquisition and Preservation: Protection of habitats is consistent with the MS TIG goal to increase connectivity of coastal habitats. Lands would be purchased in fee from willing sellers at appraised value. Acquisition and preservation includes the purchase of land and preservation into perpetuity, facilitating protection of habitats on the parcels through prevention of development. Acquisition of parcels would only be made at appraisal value. Acquisition and preservation would apply to up to 8,000 acres of various habitats including forested freshwater scrub-shrub, coastal marsh, pine savannas and flatwoods, freshwater marsh and beach as listed in Table 3.4-2. Acquired properties would then be held in trust and managed into perpetuity. The proposed project time frame is limited to 15 years. Acquisition and preservation would apply to Alternative B and Alternative D (Preferred).

Invasive species Management: Invasive species management will focus on prevention, control and eradication of known exotic invasive plant species in the project area for the proposed alternatives. Example species include, but are not limited to, Chinese privet (*Ligustrum sinense*), Chinese tallow (*Sapium sebiferum*), common reed (*Phragmites australis*), Cogon grass (*Imperata cylindrica*), Japanese climbing fern (*Lygodium japonicum*), Japanese honeysuckle (*Lonicera japonica*) and others. A number of techniques are commonly utilized on the NWR and NERR and at the nearby Sandhill Crane NWR to accomplish this, incorporated by reference here (USFWS 2007, USFWS 2008, GBNERR 2016). For example, prescribed fire is used for both reduction of fuel loads and invasive species management in savannas to promote grassy-herbaceous ground cover. For the purposes of discussion and to facilitate a programmatic impact analysis, invasive species management techniques will be divided into three categories; 1. Chemical Treatment; 2. Mechanical Treatment, and 3. Prescribed Fire, which are described below. Resource managers could use an integrated approach including a variety of techniques for site specific restoration and management measures depending on existing habitat conditions.

- 1) Chemical Treatment: Chemical treatments could include basal-bark application, cut stump treatments, foliar spray applications, or stem injection of herbicides to target eradication or control of invasive plant species. These applications are typically completed seasonally in target areas. Activities could require the vehicular transport of personnel into areas, use of approved herbicides, use of established safety and containment procedures, and the targeted application of herbicide in small areas. Personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities. Treatments are typically done in areas that range from several acres up to 50 acres for a large-scale treatment by trained personnel. Within the proposed project area, chemical treatment would be limited to small areas within 3,311 acres of forested freshwater scrub shrub habitat, 6,276 acres of pine savannas and flatwoods, and 1,937 acres of freshwater marsh. Chemical treatment would be applicable to Alternative C and Alternative D (Preferred).
- 2) Mechanical Treatment: Mechanical treatment is often used in combination with prescribed fire to reduce woody vegetation and trees in target habitats. Use of these techniques result in an increase in savanna species including sun-loving graminoids (grass-like plants) and forbs (flowering plants) (desired conditions in this area). Mechanical treatment could include removal of trees using commercial tree contracts, chain saws, bulldozing, use of a bulldozer or gyrotrac with roller chopper to remove shrubs and small trees or drum chopping to push over and crush small, pre-commercial pines and shrubs. In wet areas, soft track or wide track equipment would be used to

distribute the equipment weight and minimize ground disturbance. Alternatively, crews may remove material with chainsaws or by hand. Replanting could also be part of habitat restoration and management operations. Mechanical treatment is used both at small and large scales successfully: several thousand acres of undesirable vegetation has been cleared in the Sandhill Crane NWR (USFWS 2007). Mowing, tilling and disking are also used to prevent the spread of invasive species such as cogon grass. Mechanical treatment would be used within 6,276 acres of savannas and flatwoods within the project area. Operations could occur over several seasons depending on the success of acquisitions and other restoration priorities. Mechanical clearing would be applicable to Alternative C and Alternative D (Preferred).

3) **Prescribed Fire**: Native habitats within the southeastern United States, including those within the project boundary, evolved in the midst of reoccurring, natural fires (USFWS 2007, USFWS 2008, GBNERR 2016). These habitats therefore depend on a reoccurring fire schedule. Habitat management agencies in the project area therefore successfully use prescribed fires to restore and maintain high quality, natural habitats. Prescribed fires reduce woody vegetation and tree encroachment in pine savanna habitat and can be effective in helping prevent the spread of certain exotic invasive species (e.g., Cogon grass and Chinese tallow), when used in combination with other methods (e.g., chemical and mechanical treatment). This project proposes to implement a schedule of prescribed fires on publicly owned property within the project boundary to accomplish habitat restoration and management goals. Wire grass, for example, is a fire-dependent savanna species. Only after being burned during the growing season will this grass produce seeds. Their complex system of underground roots and shoots helps them survive the fire. By increasing species such as this, the project is also expected to provide services to wildlife that use them, such as many declining populations of grassland bird species that rely on savanna habitat.43 Historically, natural fire occurred on a three to five-year interval. Fires were of low intensity, fueled by grasses and pine litter. Prescribed fire and associated management within the project boundary would simulate these historic, natural fires.

Site preparation for a prescribed fire often involves compression of vegetation using equipment like roller choppers, gyrotracks, and excavators and/or other mechanical treatments included above to create habitat conditions which facilitate desired burns. Clearing, plowing and disking may be used to prepare fire breaks, zones devoid of fuel that border burn units and help manage burn boundaries. Fire could be applied using handheld drip torches to initiate prescribed fire. Aerial ignition from helicopters could also be used. Prescribed burns would follow standardized planning protocols and methodologies, such as considering environmental factors (certain weather, fuel and moisture conditions that would make the fire manageable 44) and burning on a 2-3 year rotation during the growing season (Spring and Summer months, when possible). Prescribed fires could range in size depending on habitats and logistics; average

102

⁴³ https://www.fws.gov/refuge/Grand Bay/what we do/resource management.html

⁴⁴ https://www.fws.gov/mississippisandhillcrane/fire.html

prescribed burns at Grand Bay NWR are 79 acres, compared to 59 acres at Mississippi Sandhill Crane NWR. Twenty percent of the Grand Bay fires reach 100 acres or more, compared to 13% at the Mississippi Sandhill Crane NWR (USFWS 2005). For Alternatives C and D prescribe fire would applied on up to 6,276 acres of savanna and flatwoods.

Best Practices: The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS to avoid and minimize impacts to resources. Best practices listed in the PDARP/PEIS are intended to evolve as an adaptive management component of implementing the PDARP/PEIS; as such, the appendix to the PDARP/PEIS is a living document. As new best practices are established, existing best practices are refined, or new techniques and information are informed by implementation, these measures will be added to or updated in the relevant websites identified in the appendix of the PDARP. In this capacity, new projects will have available the current range of best practices to support project design and implementation. In addition to PDARP/PEIS best practices, the MS TIG could develop best practices for site-specific restoration measures and management activities in different locations due to differences in relevant site conditions.

3.4.1 Grand Bay Land Acquisition and Habitat Management Alternatives B-D: Affected Environment and Environmental Consequences

This section discusses proposed Grand Bay Land Acquisition and Habitat Management Alternatives B, C, and D (listed below). Proposed Alternative D is one of two preferred alternatives for WCNH/Birds. The other alternative, Proposed Alternative A (Preferred): Graveline Land Acquisition and Management was discussed above in Section 3.3.

- Alternative B: Grand Bay Land Acquisition (Up to 8,000 acres)
- Alternative C: Grand Bay Habitat Management (Up to 17,500 acres)
- Alternative D (Preferred): Grand Bay Land Acquisition and Habitat Management (Alt. B + C)

Alternative B: Grand Bay Land Acquisition (Up to 8,000 acres)

The proposed action for Alternative B would include the acquisition of privately owned land from willing sellers of up to 8,000 acres of habitat within the boundaries of the Grand Bay NWR, Grand Bay NERR and Grand Bay Savanna CP (Figure 3.4-4). Habitat management activities described in Section 3.4 would not be implemented on newly acquired lands. Habitat management activities as currently planned and implemented under existing management plans and policies would continue on publicly owned lands.

Alternative C: Grand Bay Habitat Management (Up to 17,500 acres)

The proposed action for Alternative C would include habitat management of 17,500 acres of publicly owned lands within the boundaries of the Grand Bay NWR, Grand Bay NERR and Grand Bay Savanna CP (Figure 3.4-4). Privately owned lands within the NWR, NERR and the CP boundaries would not be acquired. Habitat management activities described in Section 3.4 above would be implemented to enhance habitat only on the existing publicly owned lands.

Alternative D (Preferred): Grand Bay Land Acquisition (up to 8,000 acres) and Habitat Management (up to 17,500 acres)

The proposed action for Alternative D (Preferred) would include acquisition of up to 8,000 acres of land and habitat management of up to 17,500 acres of currently owned and newly acquired lands within the boundaries of the Grand Bay NWR, Grand Bay NERR and the Grand Bay Savanna CP (Figure 3.4-4).

Project Location

The project location consists of parcels adjacent to and near Grand Bay in Jackson County, Mississippi. The project is located in Jackson County, Mississippi in the boundaries of Grand Bay NWR, NERR and Grand Bay Savanna CP (Figure 3.4-4).

3.4.1.1 Overview of Affected Environment and Environmental Consequences

This analysis incorporates by reference the MS portions of that affected environment as relevant from Sections 3.5.1 and 3.3.6 of the PDARP/PEIS. Likewise, the PDARP/PEIS provides programmatic evaluation of the environmental consequences of the restoration approaches "Protect and conserve marine, coastal, estuarine and riparian habitats" and restore and conserve bird nesting and foraging habitat" which are considered in this Draft RP/EA. PDARP/PEIS evaluations from sections 6.4.1.5 and 6.4.10 are incorporated by reference here. Tiering from that analysis, this section presents the Affected Environment of Grand Bay and environmental consequences of the proposed actions in context of the site-specific affected environment.

The programmatic analysis looked at a series of resources as part of the biological, physical, and socioeconomic environment. As appropriate in a tiered analysis, the evaluation of each alternative focuses on the specific resources with a potential to be affected by the proposed project. To avoid redundant or unnecessary information, resources that are not expected to be affected are evaluated summarily in the respective sections. These resources include, noise, marine and estuarine fauna, infrastructure, fisheries and aquaculture, marine transportation, and aesthetics and visual resources which will be discussed in Sections 3.4.1.2, 3.4.1.3, and 3.4.1.4.

3.4.1.2 Physical Environment

<u>Programmatic Review of Environmental Consequences (Physical Environment)</u>: Sections 6.4.1.5 and 6.4.1.10 of the PDARP/PEIS describe the impacts to Physical Resources for the relevant restoration approaches and are incorporated by reference and briefly described here.

PDARP/PEIS consequences related to geology and substrates, water resources, and air quality: Specific restoration activities identified as part of land management plans could result in short-term, minor to moderate adverse effects on geology, substrates, and water resources. Fire management may have short-term adverse impacts on soils, substrates, and air quality. Land acquisition could permit public access for recreational use which could result in short-term, minor to moderate adverse effects through increased soil compaction, rutting, or erosion caused by human presence and activity within the conservation area. Increased public use could result in short-term, minor effects on surface water through increased sedimentation. Fee title land acquisition could reduce disturbance of geology and substrates by protecting lands from development pressure. This would be a long-term beneficial effect that will extend the life of the project.

<u>PDARP/PEIS</u> consequences related to hydrology and water quality: Where protected lands overlap ground water recharge zones, surface water, or brackish-water resources, water sources and water quality could be further protected from future degradation by helping to reduce runoff. Similarly, where protected land overlaps wetlands or shorelines, the protection of natural hydrologic processes could indirectly help limit development and associated effects on water quality, including via saltwater intrusion. These would be long-term beneficial effects.

Environmental consequences for the proposed alternatives are within the general range impacts as described in the PDARP/PEIS with some variances related to specific actions. As appropriate in a tiered analysis, the evaluation of the proposed alternative focuses on the specific resources with a potential to be affected. Noise impacts for the proposed alternative would be negligible to minor. To avoid redundant or unnecessary information, noise is evaluated here.

Noise: Restoration measures and management activities that would have adverse noise impacts would occur primarily in savannas and flatwoods (mechanical treatment associated with prescribed fire. There would be short-term, minor, adverse noise impacts from equipment and operations associated with mechanical treatment, establishment of fire breaks, prescribed fire operations, and road repair/removal and culvert replacement. Restoration activities would occur sporadically and seasonally and would be dependent on successful acquisitions. The operations would be short-term and are remote. Noise receptors in the area of the work would be buffered by forested areas between the receptor and the site of noise-producing activity. Acquisition and preservation of developable areas would provide a long-term benefit by reducing ambient noise pollution when compared to a build out scenario if property were developed. In addition, the following best practice would be implemented for the proposed alternative to the extent practicable: Minimize construction noise to the maximum extent practicable when working near protected species and their habitats.

For the physical environment, the following resources are further analyzed below:

- Geology and substrates
- Water Quality and Hydrology
- Air Quality and Greenhouse Gas Emissions

3.4.1.2.1. Geology and Substrates

Affected Environment

Section 3.3.3 of the PDARP/PEIS discusses the geomorphic zones of the northern Gulf of Mexico. The Grand Bay Land Acquisition and Habitat Management project area for proposed alternatives is located within the Gulf Coastal Plain and the Mississippi Alluvial Plain physiographic regions. Landforms and substrates are generally comprised of Holocene sediments. These sediments are composed of sand, silt, and clay with comparatively high organic matter content. The coastal estuaries of Mississippi are composed of mostly sandy fine-grained sediment, silt and clays (Schmid 2015).

Seismic activity in the project area for proposed alternatives is low. Since the late 1800s, about ten earthquakes large enough to be detected have occurred in the Gulf of Mexico. These earthquakes were mostly small-magnitude events (magnitudes of 3 to 4 on the Richter scale).

Data from the Mississippi State Geological Survey generally indicates that surface soils in the project area for the proposed alternatives consist of Holocene age coastal deposits of loam, sand, gravel, and

clay. The United States Department of Agriculture (USDA) Natural Resource Conservation Service (NRCS) Web Soil Survey identifies 32 soil-mapping units within the footprint of the proposed project. These soil map units located within the project footprint area are listed on Table 3.4-3 (NRCS 2016). Of these soils Atmore loam, 1 to 3 percent slopes; Axis mucky sandy clay loam, frequently flooded; Bayou sandy loam, 0 to 1 percent slopes; Croatan and Johnston soils, frequently flooded; Daleville loam, ponded; Daleville silt loam, 0 to 1 percent slopes; Handsboro mucky silt loam, frequently flooded; Harleston fine sandy loam, 0 to 2 percent slopes; Hyde silt loam; Johns loamy fine sand, 0 to 2 percent slopes; Kinston, Chastain, and Mantachie soils, frequently flooded; Myatt loam, 0 to 1 percent slopes, occasionally flooded; Nugent and Jena soils, frequently flooded; Ocilla loamy sand, 0 to 2 percent, occasionally flooded; Smithton loam, 0 to 1 percent slopes, occasionally flooded; and Stough loam, 0 to 2 percent slopes are listed as hydric (NRCS 2016a). Soils characteristics are listed in Table 3.4-3.

Table 3.4-3: Soils Characteristics

Soil Type	Texture	Drainage Class
Kinston, Chastain, and Mantachie soils, frequently Flooded	Fine Sandy Loam (upper) Sandy Clay Loam (lower)	Poorly Drained
Atmore loam, 1 to 3 percent Slopes	Loam (upper) Loam (lower)	Poorly Drained
Lenoir silt loam, 0 to 1 percent Slopes	Silt Loam (upper) Clay (lower)	Somewhat Poorly Drained
Daleville silt loam, 0 to 1 percent Slopes	Silt Loam (upper) Clay Loam (lower)	Poorly Drained
Daleville loam, ponded	Loam (upper) Clay Loam (lower)	Poorly Drained
Eustis loamy sand, 0 to 5 percent slopes	Loamy Sand (upper) Loamy Sand (lower)	Somewhat Excessively Drained
Eustis loamy sand, 5 to 12 percent slopes	Loamy Sand (upper) Loamy Sand (lower)	Somewhat Excessively Drained
Bigbee loamy sand, 0 to 5 percent slopes, occasionally flooded	Loamy Sand (upper) Fine Sand (lower)	Somewhat Excessively Drained
Myatt loam, 0 to 1 percent slopes, occasionally flooded	Loam (upper) Sandy Clay Loam (lower)	Poorly Drained
Hyde silt loam	Silt Loam (upper) Silt Clay Loam (lower)	Very Poorly Drained
Smithton loam, 0 to 1 percent slopes, occasionally flooded	Loam (upper) Sandy Loam (lower)	Poorly Drained
Johns loamy fine sand, 0 to 2 percent slopes	Loamy Fine Sand (upper) Sandy Clay Loam (lower)	Somewhat Poorly Drained
Vancleave loamy sand, 0 to 2 percent slopes	Loamy Sand (upper) Sandy Loam (lower)	Moderately Well Drained
Vancleave loamy sand, 2 to 5	Loamy Sand (upper)	Moderately Well Drained

Soil Type	Texture	Drainage Class
percent slopes	Sandy Loam (lower)	
Escambia very fine sandy loam, 0 to 2 percent slopes	Very Fine Sandy Loam (upper) Loam (lower)	Somewhat Poorly Drained
Malbis fine sandy loam, 0 to 2 percent slopes	Fine Sandy Loam (upper) Loam (lower)	Well Drained
Ocilla loamy sand, 0 to 2 percent slopes, occasionally flooded	Loamy Sand (upper) Loamy Sand (lower)	Somewhat Poorly Drained
Benndale fine sandy loam, 3 to 8 percent slopes	Fine Sandy Loam (upper) Loam (lower)	Well Drained
Prentiss silt loam, 0 to 2 percent Slopes	Silt Loam (upper) Loam (lower)	Moderately Well Drained
Stough loam, 0 to 2 percent Slopes	Loam (upper) Sandy Loam (lower)	Somewhat Poorly Drained
Freest sandy loam, 2 to 5 percent slopes	Sandy Loam (upper) Loam (lower)	Moderately Well Drained
Nugent and Jena soils, frequently flooded	Loamy Sand (upper) Stratified Sand to Fine Sandy Loam (lower)	Excessively Drained
Wadley loamy sand, 0 to 5 percent slopes	Loamy Sand (upper) Sandy Clay Loam (lower)	Somewhat Excessively Drained
Croatan and Johnston soils, frequently flooded	Muck (upper) Fine Sandy Loam (lower)	Very Poorly Drained
Udorthents	Loamy Sand (upper) Sandy Loam (lower)	Moderately Well Drained
Axis mucky sandy clay loam, frequently flooded	Mucky Sandy Clay Loam (upper) Sandy Loam (lower)	Very Poorly Drained
Handsboro mucky silt loam, frequently flooded	Mucky Silt Loam (upper) Muck (lower)	Very Poorly Drained
Maurepas muck, frequently Flooded	Muck	Very Poorly Drained
Bayou sandy loam, 0 to 1 percent slopes	Sandy Loam (upper) Sandy Loam (lower)	Poorly Drained
Harleston fine sandy loam, 0 to 2 percent slopes	Fine Sandy Loam (upper) Sandy Loam (lower)	Moderately Well Drained
Harleston fine sandy loam, 2 to 5 percent slopes	Fine Sandy Loam (upper) Sandy Loam (lower)	Moderately Well Drained
Columbus loam, 0 to 2 percent slopes, occasionally flooded	Loam (upper) Loam (lower)	Moderately Well Drained

 $Environmental\ Consequences\ for\ WCNH/Birds\ Proposed\ Alternatives\ B,\ C\ and\ D\text{-}\ (Preferred)$

Table 3.4-4 lists environmental consequences of the project activities to substrates (soils). There would be no affect to geologic resources by implementing the proposed alternatives.

Table 3.4-4: Proposed Alternatives -Environmental Impacts to Geology and Substrates

Restoration Measure	Alternative B: Grand Bay Land Acquisition			Alternative C: Grand Bay Habitat Management			Alternative D (Preferred): Grand Bay Land Acquisition and Habitat Management		
	Adverse	Adverse	Beneficial	Adverse	Adverse	Beneficial	Adverse	Adverse	Beneficial
	Impact	Impact	Impact	Impact	Impact	Impact	Impact	Impact	Impacts
	Duration	Intensity	Duration	Duration	Intensity	Duration	Duration	Intensity	Duration
Acquisition/	long-						long-		
Preservation	term	Minor					term	minor	
Chemical				short-			short-		
Treatment				term	minor		term	minor	
Mechanical				short-	minor to		short-	minor to	
Treatment				term	moderate		term	moderate	
Prescribed				short-			short-		
Fire				term	moderate		term	moderate	

<u>Acquisition/Preservation</u>: Acquisition and preservation would open new areas to recreational activities including hiking, fishing, bird watching, and camping. The increased public use could result in a long-term, minor, and adverse impact to soils due to potential compaction, but these would be limited to relatively small areas. There would be no adverse impacts to geology from acquisition and preservation. Impacts would be applicable to proposed Alternative B and D.

<u>Chemical Treatment</u>: Treatment activities could require the use of ATVs, pickups or other small equipment that could result in soil disturbance, rutting, and compaction. The use of equipment would result in a short-term, minor adverse impact to soils. Removal of nuisance species and replanting could result in short-term, minor, adverse impacts to soils. There would be no adverse impacts to geology from chemical treatment. Impacts would be applicable to proposed Alternative C and D.

<u>Mechanical Treatment</u>: Activities include but would not be limited to use of brush-hog, mowing, disking, and use of chainsaws. In addition, use of gyro-tracs and in some cases bobcats or bulldozers o lay down or remove vegetation could be used. Turning over soils, soil compaction, disturbance and/or rutting from equipment use could result in short-term, minor to moderate, adverse impacts, depending on the size of the operation, soils wetness and season of the operation. To minimize these effects, care would be taken in the selection of equipment used and timing of operations, particularly in wetter conditions. There would be no adverse impacts to geology from mechanical treatment. Impacts would be applicable to proposed Alternative C and D.

Prescribed Fire: Preparations for prescribed fires could include installation of fire breaks, and use of light to heavy equipment to fell or lay down woody underbrush. Fire breaks would be constructed around the boundary of the burn unit by clearing and or disking. Soils would be turned and mineral soils layers exposed. Soil could be disturbed and compacted during the burn operations due to equipment use. Vegetation laydown/removal operations using light to heavy equipment could result in soil disturbance or rutting. In wet areas, soft track or wide track vehicles could be used to distribute the equipment weight and minimize impact. Alternatively, crews may remove material with chainsaws. There could be short-term, moderate, adverse impacts to mineral soil exposure, rutting, and soil disturbance during the site preparation and burn operations. There would be no adverse impacts to geology from prescribed fire. Impacts would be applicable to proposed Alternative C and D.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to geology and substrates (soils):

- Allow revegetation of fire breaks or actively revegetate with native species or annual grasses, if prolonged period of greening up is anticipated.
- Develop and implement spill prevention and response plan, including conducting daily inspections during chemical treatment, mechanical treatment, and prescribed fire operations to ensure there are no leaks of antifreeze, hydraulic fluid, pesticide, or other substances.
- To the extent practicable, for equipment use in wet areas, soft tracked or wide tracked equipment should be used to distribute the equipment weight and minimize impacts to soils. Alternatively, crews may remove vegetative materials with chainsaws.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. Under Alternatives B and D, if development were to occur, there would likely be impacts to soils Acquiring the parcels would prevent them from being developed or from structures being constructed on them. Acquiring the parcels would place them under the purview of resource managers and management plans that would help conserve and protect the resource. Under Alternative C, the No Action alternative would not provide the additional benefits to soils described above. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.2.2 Hydrology and Water Quality

Affected Environment

Section 3.3.2 of the PDARP/PEIS addresses river flows on the northern gulf geography and water quality. Section 6.14.2 discusses future sea level rise, storm surge and storm intensity projections and is incorporated by reference here. In the project area for the proposed alternatives, the affected resources consist of shallow water within bays, bayous, and wetlands within Grand Bay. Mississippi's water quality standards specify the appropriate levels for which various water quality parameters or indicators support a water body's designated use(s). Each use assessed for a water body is determined to be either "Attaining" or "Not Attaining" in accordance with the applicable water quality standards and U.S. Environmental Protection Agency (EPA) guidelines for assessments pursuant to §305(b). A water body's use is said to be impaired when based on current and reliable site-specific data of sufficient quantity, quality, and frequency of collection it is not attaining its designated use(s). Where data and information of appropriate quality and quantity indicate non-attainment of a designated use or uses for an assessed water body, the water body will be placed on the Mississippi 2014 Section 303(d) List of Impaired Water Bodies (MDEQ 2014).

The proposed alternatives are in a region with abundant annual rainfall, receiving more than 64 inches per year (USFWS 2008). The proposed alternatives are in the Mississippi Coastal Streams watershed, the Pascagoula Watershed, and the Escatawpa Watershed. These three watersheds include portions of George, Greene, Jackson, Wayne, Perry, Hancock, Harrison, Pearl River, and Stone counties; however, the project area for the proposed alternatives is exclusively in Jackson County. Major tributaries within the Mississippi Coastal Streams watershed include Bayou Casotte, Wolf River, Rotten Bayou, DeLisle Bayou, Bayou La Croix, Bayou Bacon/Jourdan River, Turkey Creek/Bernard Bayou, Biloxi River, and Tuxachanie Creek. Major tributaries within the Pascagoula River watershed include Okatoma Creek, Leaf River, Black Creek, Red Creek, Pascagoula River, Escatawpa River, Chickasawhay River, Thompson Creek, and Tallahala Creek.

Major rivers carry high sediment loads into the Mississippi Sound. Inland fresh water drainage from these and other smaller rivers create an estuarine environment. Variable salinity levels can affect the productivity and survival of organisms living in the area, as well as economic and recreational activities. Pollution from agriculture, improperly treated sewage, roadways, accidental oil spills, industry discharges, and other sources also affect the health of the habitats. Grand Bay is influenced by freshwater flow from Southwest Bayou, Middle Bayou, Clay Bayou, Bayou Cumbest and Bayou Heron. The Grand Bay Land Acquisition and Habitat Management proposed alternatives are located in waters classified by the State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters (MDEQ 2012) as "shellfish harvesting 45", "recreation 46", and "fish and wildlife 47" (Bang's Lake), and "recreation" and "fish and wildlife 48" for all other areas in the project location. Bayou Cumbest, which drains directly into Grand Bay, is listed as impaired on the State of Mississippi 303(d) list (MDEQ 2014) for Organic Enrichment/Low Dissolved Oxygen.

Floodplains

A large portion of the proposed alternative area is mapped as Zone VE. Zone VE is defined as Coastal flood zone with velocity hazard. This includes beach areas, open water and most estuarine marsh. Some estuarine marsh, streams, and riparian areas in the proposed alternative project area are mapped as Zone AE. Zone AE is defined as "Base Flood Elevations Determined". Upland areas in the proposed alternative project area are mostly Zone X. Zone X are defined as "Areas of 0.2% annual change flood; areas of 1% annual chance flood with average depths of less than 1 foot or with drainage areas less than 1 square mile; and areas protected by levees from 1% annual chance flood".

-

⁴⁵ Waters in the shellfish harvesting classification are for propagation and harvesting shellfish for sale or use as a food product.

⁴⁶ Waters in the recreation classification are to be suitable for recreational purposes, including such water contact activities as swimming and water skiing.

⁴⁷ Waters in the fish and wildlife classification are intended for fishing and for propagation of fish, aquatic life, and wildlife

⁴⁸ Waters that meet the Fish and Wildlife criteria are also be suitable for secondary contact recreation.

Wetlands

The project area for the proposed alternatives is a mosaic of wetlands and uplands extending from the open water, salt pannes in Grand Bay up to mesic and wet savanna and flatwoods near I-10. Wetlands in the proposed alternative project area include forested freshwater scrub shrub wetlands, coastal marsh, wet savannas and flatwoods, and freshwater marsh (See Habitats in Section 3.4.1.3).

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred) Environmental consequences affecting hydrology, water quality, floodplains and wetlands are discussed below.

Hydrology

Table 3.4-5 lists the environmental consequences of each project activity to hydrology in the project area for the proposed alternatives.

Table 3.4-5: Proposed Alternatives -Environmental Impacts to Hydrology and Water Quality

Table 5.4-5. I Toposed Alternatives -Environmental impacts to rivurology and water Quanty									
Restoration Measure	Alternative B: Grand Bay Land Acquisition			Alternative C: Grand Bay Habitat Management			Alternative D (Preferred): Grand Bay Land Acquisition and Habitat Management		
	Adverse	Adverse	Beneficial	Adverse	Adverse	Beneficial	Adverse	Adverse	Beneficial
	Impact	Impact	Impact	Impact	Impact	Impact	Impact	Impact	Impact
	Duration	Intensity	Duration	Duration	Intensity	Duration	Duration	Intensity	Duration
	Hydrology								
Acquisition/									
Preservation			long-term						long-term
Chemical				short-			short-		
Treatment				term	minor		term	minor	
Mechanical				short-	minor to		short	minor to	
Treatment/				term	moderate		term	moderate	
Prescribed Fire									
				Water	Quality				
Acquisition/									
Preservation			long-term						long-term
Chemical				short-			short-		
Treatment				term	minor		term	minor	
Mechanical				short-	minor to		short-	minor to	
Treatment/				term	moderate		term	moderate	
Prescribed Fire									

<u>Acquisition/Preservation</u>: Acquisition and preservation would open new areas to recreational activities including hiking, fishing, bird watching, and camping. Preservation of lands would have indirect, long-term, benefits to both hydrology and water quality by preventing development and disturbances, and stormwater infrastructure. Impacts would be applicable to proposed Alternative B and D.

<u>Chemical Treatment</u>: There could be short-term, minor impacts to hydrology as a result of minor rutting/soil disturbance and temporary changes in hydrologic patterns from vehicular transport of personnel to treatment areas. Impacts would be applicable to proposed Alternatives C and D.

<u>Mechanical Treatment/Prescribed Fire</u>: Mechanical treatment would apply to up to 6,276 acres of savannas and flatwoods. Since large equipment may be needed, soil disturbance, rutting, compaction could have short-term, minor to moderate, adverse impact to hydrology. There could be small, temporary changes to stormwater flows and runoff retention patterns due to rutting by equipment and

vegetation removal resulting in a short-term, minor to moderate, adverse impact to hydrology. There would be short-term, minor to moderate, adverse impacts to hydrology resulting from mechanical treatment of woody underbrush and construction of fire breaks. There could be small, temporary changes to stormwater flows and runoff retention patterns due to rutting by equipment and vegetation removal. Soft track or wide track equipment would be used in wet areas to the extent practicable. Alternately, crews may remove vegetative material with chainsaws. Impacts would be applicable to proposed Alternative C and D.

Water Quality

Table 3.4-17 lists the environmental consequences of each project activity to water quality in the project area for the proposed alternatives.

<u>Acquisition/Preservation</u>: Acquisition and preservation would open new areas to recreational activities including hiking, fishing, bird watching, and camping. Access via motorized vehicles would be limited. Preservation of lands would have indirect, long-term benefits by preventing development and disturbances, which could reduce surface water runoff and result in long-term water quality benefits to Grand Bay. Impacts would be applicable to proposed Alternative B and D.

Chemical Treatment: Chemical treatment activities would include the use of herbicides. There could be unavoidable spills near the intended application area. However, best practices would be used to prevent any harmful chemicals from entering the environment. Implementation of best practices that the MS TIG would consider, described in section 3.4.1.2.1 above includes development and implementation of a spill prevention and response plan, including conducting daily inspections during chemical treatment to ensure there are no leaks of pesticides or other substances. Personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities. As such, this activity would have short-term, minor, adverse impact, if any, on water quality and wetlands (described below). Impacts would be applicable to proposed Alternative C and D.

Mechanical Treatment/Prescribed Fire: Mechanical treatment would apply to up to 6,276 acres of savannas and flatwoods. Since large equipment may be needed, soil disturbance, rutting, compaction and any resulting rutting and compaction could have short-term, minor to moderate, adverse impact to water quality. There could be small, temporary changes to stormwater flows and runoff retention patterns and resulting sediment movement due to rutting by equipment and vegetation removal resulting in a short-term, minor to moderate, adverse impact to water quality. Similar impacts could result from mechanical treatment of woody underbrush and construction of fire breaks. Soft tracked or wide tracked equipment would be used in wet areas to the extent practicable. Alternately, crews may remove vegetative material with chainsaws. In addition, appropriate erosion control plans would be developed as necessary to prevent sediment movement from the mechanical treatment/prescribed fire area. Impacts would be applicable to proposed Alternative C and D.

Floodplains

Acquisition and preservation of land in perpetuity would prevent land development in floodplains. There would be a long-term benefit to floodplains. Chemical treatment, mechanical treatment, and prescribed fire operations would not result in a detectable change to natural and beneficial floodplain values. Impacts would be applicable to proposed Alternative B and D.

Wetlands

Table 3.4-6 is a summary of proposed alternatives impacts to wetlands.

Table 3.4-6: Proposed Alternatives -Environmental Impacts to Wetlands

Restoration Measure	Alternative B: Grand Bay Land Acquisition			Alternative C: Grand Bay Habitat Management			Alternative D (Preferred): Grand Bay Land Acquisition and Habitat Management		
	Adverse	Adverse	Beneficial	Adverse	Adverse	Beneficial	Adverse	Adverse	Beneficial
	Impact	Impact	Impact	Impact	Impact	Impact	Impact	Impact	Impact
	Duration	Intensity	Duration	Duration	Intensity	Duration	Duration	Intensity	Duration
				Wetla	nds				
Acquisition/									
Preservation			long-term						long-term
Chemical				short-			short-		
Treatment				term	minor		term	minor	
Mechanical				short-	minor to		short-	minor to	
Treatment				term	moderate	long-term	term	moderate	long-term
Prescribed Fire				short-	minor to		short-	minor to	
				term	moderate	long-term	term	moderate	long-term

<u>Acquisition/Preservation</u>: There would be a long-term benefit to wetlands from acquisition and preservation. Wet savannas and flatwood, forested freshwater scrub shrub, freshwater marsh, and coastal marsh areas that are acquired would not be impacted for development. Impacts would be applicable to proposed Alternative B and D.

<u>Chemical Treatment</u>: Chemical treatment activities would require the use of herbicides and equipment during application. Personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities. Only chemicals approved for use in wetlands would be used. Equipment traffic in wetlands would be avoided to the extent practicable. Best practices would be used during the application of herbicides. Accidental spillage could result in short-term, minor, adverse impacts to wetland habitat. However, best practices would be used to prevent any harmful chemicals from entering the environment and for clean up if a spill occurred. Impacts would be applicable to proposed Alternative C and D.

Mechanical Treatment: Mechanical treatment in wetland areas would be done in a manner that would minimize impacts to soil to the extent practicable. In wet areas, soft track or wide track equipment would be used to distribute the equipment weight and minimize impact. Alternatively, crews may remove material with chainsaws. If required, a USACE permit would be obtained likely a Nationwide 27 (Aquatic Habitat Restoration, Establishment, and Enhancement Activities), as well as a MDMR Coastal Wetlands Permit (if required). Nationwide 27 allows for mechanized land clearing to remove non-native invasive, exotic, or nuisance vegetation and other related activities. If there is any clearing within wetlands or stream boundaries, damage to vegetation, soil compaction and any resulting erosion could have a short-term, minor to moderate impact to wetlands. USACE permit and/or MDMR Coastal Wetlands permit conditions (if required) would be adhered to in all operations. There would be long-term benefits to wet savannas and flatwoods from mechanical clearing including establishment of more native flora and increased diversity in flora and fauna. Impacts would be applicable to proposed Alternative C and D.

<u>Prescribed Fire</u>: Prescribed fire would apply to up to 6,276 acres of savannas and flatwoods, a portion of which, are likely wetlands. Intermittent fires were historically a critical perturbation in for this

habitat. There would be short-term, minor to moderate impacts resulting from mechanical treatment of woody underbrush and construction of fire breaks if the fire breaks are in wetlands or streams; the impacts, permit requirements and minimization measures are discussed above in mechanical treatment. There would be long-term beneficial effects to wet fire-suppressed pine savannas including a re-establishment of wetland communities, and increased diversity in flora and faunal populations that colonized the prescribed burn unit. Impacts would be applicable to proposed Alternative C and D.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to water quality and hydrology:

- In the execution of land acquisition and the design of habitat management measures the MS TIG would consider resiliency measures to facilitate habitat migration due to sea level rise (CEQ, 2016).
- Develop and implement an erosion control plan to minimize erosion during and after construction and where possible use vegetative buffers (100 feet or greater), revegetate with native species or annual grasses, and conduct work during dry seasons.
- For chemical treatment, personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities.
 Personnel will apply herbicide in accordance with the direction and guidance provided on the appropriate U.S. Environmental Protection Agency (EPA) labels and state statutes during land-based activities.
- Soft track or wide track equipment would be used in wet areas to the extent practicable. Alternatively, crews may remove vegetative material with chainsaws.
- Avoid and minimize, to the maximum extent practicable, placement of dredged or dill
 material in wetlands and other aquatic resources. Design construction equipment corridors to
 avoid and minimize impacts to wetlands and other aquatic resources to the maximum extent
 practicable. If required, a USACE permit would be obtained; likely a Nationwide 27
 (Aquatic Habitat Restoration, Establishment, and Enhancement Activities) as well as MDMR
 Coastal Wetlands Permit (if required). USACE permit and/or MDME Coastal Wetlands
 permit conditions (if required) would be adhered to in all operations.
- Designate a vehicle staging area removed from any natural surface water resource or wetland
 to perform fueling, maintenance, and storage of construction vehicles and equipment. Inspect
 vehicles and equipment daily prior to leaving the storage area to ensure that no petroleum or
 oil products are leaking.
- Develop and implement a spill prevention and response plan, including conducting daily inspections of all construction and related equipment to ensure there are no leaks of antifreeze, hydraulic fluid, or either substances and cleaning and sealing all equipment that would be used in the water to rid it of chemical residue.
- Control dust related to construction site activities through a Soil Erosion Sediment Control Plan that includes spraying of a suppressing agent on dust piles (non-hazardous, biodegradable).

• Cover trucks hauling loose materials.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. Under Alternatives B and D, if development were to occur, there would likely be adverse impacts to hydrology, water quality, floodplains, and wetlands. Adverse hydrologic affects could include increased runoff rates due to impervious surfaces related to development. Increases in sediment entering waterways could result in adverse effects to water quality. Floodplain and wetland function could be adversely affected by development of parcels proposed for acquisition, preservation and management under proposed WCNH/Birds Alternatives B and D. Under Alternative C, the No Action alternative would not provide the additional benefits to hydrology, water quality, floodplains, and wetlands described above. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.2.3. Air Quality and Greenhouse Gas Emissions

Affected Environment

The following section is a discussion of air quality for the project area for the proposed alternatives. EPA has set national ambient air quality standards (NAAQS) for six principal air pollutants (also called criteria pollutants): Ground-Level Ozone (O3), Particulate Matter (PM), Nitrogen Dioxide (NO2), Sulfur Dioxide (SO2), Carbon Monoxide (CO), and Lead (Pb). MDEQ is the state agency responsible for development and maintenance of state specific air emission standard for Mississippi, and monitors all of these pollutants. In Jackson County, the following parameters are monitored: Ozone, Particulate Matter, Nitrogen Oxides, and Sulfur Dioxide. According to MDEQ 2015 Air Quality Data Summary⁴⁹ the entire state of Mississippi, including Jackson County, is meeting all of the NAAQS.

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred)

The environmental consequences for this section is divided into two discussions: 1- environmental consequences from equipment operation/best practices and; 2- environmental consequences resulting from prescribed fire/best practices.

<u>1-Environmental Consequences Resulting From Equipment Operation/Best Practices:</u> The following project implementation activities would produce emissions during equipment operation: chemical treatment, mechanical treatment, and road removal/culvert replacement. Because these restoration activities would occur seasonally, and would be limited in scope and distribution, the impacts on air quality or to emissions of greenhouse gases would be short-term and minor.

Best Practices

_

⁴⁹http://www.deq.state.ms.us/mdeq.nsf/pdf/Air 2015AirQualityDataSummary/\$File/2015%20Air%20Quality%20Data%2 0Summary.pdf

Unavoidable short-term and minor adverse impacts from equipment emissions would be offset through the following best practices measures to the extent practicable:

- Shut down idling construction equipment, if feasible.
- Encourage the use of the proper size of equipment for the job to maximize energy efficiency.
- Use of ultra-low sulfur diesel fuel in off-road construction equipment with engine horsepower (HP) rating of 60 HP and above.
- <u>2- Environmental Consequences Resulting from Prescribed Fire/Best Practices</u>: The use of controlled burns is included in this project as a restoration activity to provide major long-term benefits for native species habitats, water and soil quality, and nutrient cycling⁵⁰. However, short-term moderate adverse impacts to air quality and greenhouse gases may occur during the controlled burn events because fire produces smoke, which is primarily composed of water vapor and carbon dioxide but also contains carbon monoxide, nitrogen oxide, hydrocarbons, particulate matter, and trace minerals. According to the National Coalition of Prescribed Fire Councils Guide to Smoke Management (September 2007 version)⁵¹, the primary concerns of smoke as an air pollutant are as follows:
 - 1) Carbon Dioxide: The emission factor for carbon dioxide for prescribed burning is 2,000-3,500 pounds/ton (pounds of emissions/ton of organic matter burned).
 - 2) Carbon monoxide: The emission factor for carbon monoxide for prescribed burning is 20-500 pounds/ton. It is classified as a criteria pollutant by EPA. As a result of rapid dilution and its instability, carbon monoxide emissions from prescribed burning are not a concern to the general public.
 - 3) Water vapor: The emission factor for water vapor for prescribed burning is 50-1500 pounds/ton. The only possible concern about water vapor is visibility reduction in the vicinity of the fire.
 - 4) Particulate matter: The emission factor for particulate matter for prescribed fire is 20-180 pounds/ton. Particulates are a criteria pollutant and can impact health and visibility. Particulates are presently the major pollutant of concern from prescribed burning. They represent a health risk by inhalation and also reduce visibility.
 - 5) Hydrocarbons: The emission factor for hydrocarbons for wildland fire is 10-40 pounds/ton. While hydrocarbons are not a criteria pollutant, they may impact health and visibility and in some cases, may contribute to excessive ozone concentrations.
 - 6) Nitrogen oxides: The emission factor for nitrogen oxides for wildland fire is 1-9 pounds/ton. Nitrogen oxides are a criteria pollutant and can impact health and visibility. The low emission factor reduces concern of ambient air quality standards on a local level; however, nitrogen oxides can affect ozone formation.
 - 7) Secondary emissions: Secondary emissions are pollutants which are formed in the atmosphere by photochemical transformation of primary emissions. They include oxidants such as ozone which is a criteria pollutant. Specific emission factors from prescribed burning are unknown but are believed to be relatively small.

⁵⁰ https://www.nrcs.usda.gov/Internet/FSE DOCUMENTS/16/stelprdb1046311.pdf

⁵¹http://www.garxfire.com/pdf%20files/The National Coalition of Prescribed Fire Councils Guide to Smoke Management.pdf

8) Air Toxics: There is an emerging concern about the potential emission of air toxics (acetaldehyde, acrolein; 1,3 butadiene; formaldehye; and polycyclic organic matter (POM). POM includes eight major categories of compounds including polycyclic aromatic hydrocarbons (PAHs) which include numerous chemicals which can be emitted from fire.

Adverse impacts to air quality by controlled burns would be minimized by the frequency and timing of the events; typically, they would be conducted every 1-3 years on managed burn areas according to the management plan. Unavoidable short-term moderate adverse impacts from controlled burns would be offset through the development of a Prescribed Burn Plan, which would include some or all of the following Best Smoke Management Practices (BSMPs) and would be part of the management plan. These BSMPs were developed by USDA Forest Service/NRCS⁵² (October 2011) to mitigate the impacts of smoke to public health (See Section 3.3.1.4.5), public safety and nuisance, and visibility. These six BSMPs have applicability depending on the type of burn, fuel to be burned, and level of effort needed to address air quality concerns. BSMPs are utilized by the individual fire manager and may be an expectation of a state-wide smoke management program and any applicable conservation plans which are in place for the project area.

Table 3.4-7: Summary of Basic Smoke Management Practices

Basic Smoke Management Practice	Benefit achieved with the BSMP	When the BSMP is Applied		
Evaluate Smoke Dispersion Conditions	Minimize smoke impacts	Before, During, After		
Monitor Effects on Air Quality	Be aware of where the smoke is going and degree it impacts air quality	Before, During, After		
Record-Keeping/Maintain a Burn/Smoke Journal	Retain information about the weather, burn and smoke. If air quality problems occur, documentation helps analyze and address air regulatory issues	Before, During, After		
Communication- Public Notification	Notify neighbors and those potentially impacted by smoke, especially sensitive receptors	Before, During		
Consider Emission Reduction Techniques	Reducing emissions can reduce downwind impacts	Before, During		
Share the Airshed – Coordination of Area Burning	Coordinate multiple burns in the area to manage exposure of the public to smoke	Before, During, After		

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. Under Alternatives B and D, if development were to occur, there would likely be adverse impacts to air quality due the potential

_

⁵² https://www.nrcs.usda.gov/Internet/FSE DOCUMENTS/16/stelprdb1046311.pdf

of development, the additional traffic and other air pollution related to development, and removal of vegetation that benefits air quality. Under the No Action alternative, prescribed fire would not take place as an additional management activity, resulting in no additional short-term, minor to moderate impacts to air quality from burning. This short term impact however would be offset by the potential for development with its resultant potential for long-term impacts. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.3 Biological Environment

<u>Introduction to Affected Environment (Biological Environment)</u>: Biological environment resources discussed in the section include habitats, wildlife, and protected species. PDARP/PEIS sections 3.4, 3.5, and 3.6 are incorporated by reference here. The affected environment for the proposed alternatives biological environment is described in respective sections below.

<u>Programmatic Review of Environmental Consequences (Biological Environment)</u>: Sections 6.4.1.5 and 6.4.1.10 of the PDARP describe the impacts to Biological Resources for the restoration approaches being considered for the proposed alternative. Specific restoration activities identified as part of land management plans could result in short-term, minor to moderate adverse effects on conservation areas. Consequences reviewed in the PDARP/PEIS are incorporated here and summarized.

<u>PDARP/PEIS</u> consequences related to Invasive species: Activities that may occur on conserved lands may result in introduction of invasive species. Use of best practices would help prevent the introduction of invasive species. Implementation of land management plans, located within or near restoration activities, could result in disturbed, removed, or altered habitats, which could cause minor to moderate, short- and long-term adverse effects on species that use those habitats for forage or nesting purposes.

<u>PDARP/PEIS</u> consequences related to <u>public</u> access: Land acquisition could permit public access for recreational use. This public use, depending on management stipulations, could result in long-term, minor to moderate adverse effects on area species through increased human presence and activity on acquired habitats.

<u>PDARP/PEIS</u> consequences related to habitat migration: Conservation of habitat through fee title acquisition and improved management could have a long-term benefit to any habitat on the property acquired or protected. Conservation would also allow for upland migration of beach, wetland, or other habitats as the sea level rises and could limit development encroachment.

<u>PDARP/PEIS</u> consequences related to habitat: Conservation of habitat through fee title acquisition could have a long-term benefit to fish, birds, and terrestrial wildlife through the protection of coastal, riparian, or terrestrial habitat. These habitats can be important for food supply and various life stages of some species. Benefits of the proposed restoration approach include conservation of bird nesting and foraging habitat that would increase bird health and reproduction by preventing habitat loss through land conversion.

<u>PDARP/PEIS</u> consequences related to access restriction: Restrictions on seasonal or overall human use that could result from changes in land management would reduce habitat degradation.

Improvements in habitat associated with this approach may draw additional visitors to the area, resulting in potential indirect adverse impacts from human presence. Human disturbance can lead to failure of nests, increased egg and chick predation, or even total colony abandonment.

<u>PDARP</u> consequences related to vegetation management: Managing vegetation is a common restoration technique to enhance habitat for specific bird species. Reducing vegetation on beaches, for example, can provide nesting and foraging habitat for birds such as such as snowy plover, least tern, black skimmer, and American oystercatcher. Conversely, adding vegetation can provide habitat for other bird species such as wading birds and brown pelicans. Common vegetation management methods include mechanical treatments, application of pesticides or herbicides, and biological control to manage plant species.

Environmental consequences for the proposed alternative are within the general range impacts as described in the PDARP/PEIS with some variances related to specific actions. As appropriate in a tiered analysis, the evaluation the proposed alternative focuses on the specific resources with a potential to be affected. Marine and estuarine fauna are not expected to be affected by the proposed alternative as there is no in-water work. To avoid redundant or unnecessary information, marine and estuarine fauna are evaluated summarily here.

Marine and Estuarine Fauna (Submerged Aquatic Vegetation, Nearshore Benthic Invertebrates, Marine Mammals, Essential Fish Habitat): There would be no in-water work. Estuarine marsh would be acquired and preserved, but there are limited management activities planned. Acquisition and preservation of habitat would prevent development in and adjacent to this habitat and preclude habitat removal or stresses that could result from shoreline development.

For the biological environment, the following resources are further analyzed in this section:

- Habitats
- Protected Species
- Migratory Birds
- Wildlife

3.4.1.3.1. Habitats

Affected Environment

Section 3.5 of the PDARP/PEIS provides a discussion of habitats of the northern Gulf of Mexico; Section 3.7.4 covers invasive species. Grand Bay is part of the Mississippi coastal bays and estuaries system, which also includes St. Louis Bay, Biloxi Bay, Back Bay of Biloxi, Pascagoula Bay, and Graveline Bay. Grand Bay is comprised of estuarine and non-estuarine wetland marsh habitat. The estuarine system is semi-enclosed with areas open access to the Gulf of Mexico, resulting in seawater that is occasionally diluted with freshwater runoff and flow. However, large volumes of freshwater do not regularly enter the Grand Bay system and salinities in the Grand Bay system are regularly recorded above 30 ppt (Grand Bay National Estuarine Research Reserve 2013). This open water estuarine area supports oyster reefs and seagrass habitats. The intertidal areas support a variety of marsh types and extensive, unvegetated salt flats. The non-tidal areas include wet pine savannas, coastal bayhead and cypress swamps, freshwater marshes, and maritime forests (Grand Bay National Estuarine Research Reserve 2013).

For the purposes of this Draft RP/EA, the MS TIG has grouped habitats and incorporated by reference the descriptions of those habitats provided in previous plans including:

- Grand Bay National Wildlife Refuge Comprehensive Conservation Plan (USFWS 2008)
- Grand Bay NERR Management Plan 2013-2018 (GBNERR 2012)
- Land Protection Plan and Final Environmental Assessment for the Expansion of Grand Bay National Wildlife Refuge (USFWS 2012)

Within the project area for the proposed alternatives, coastal wetland and nearshore habitats include forested freshwater scrub-shrub, coastal marsh, savannas and flatwoods, freshwater marsh, beach, and open water (Figure 3.4-4).

<u>Forested Freshwater Scrub-Shrub</u>: Approximately 3,311 acres of forested freshwater scrub-shrub exists within the project area for the proposed alternatives, 1,416 acres within acquisition parcels. This habitat was described in detail in the Grand Bay National Wildlife Refuge Comprehensive Conservation Plan (USFWS 2008), and is comprised of the following habitats: pine scrub, short scrub, tall scrub and pocosin.

<u>Coastal Marsh</u>: Approximately 9,080 acres of coastal marsh exists in the project area for the proposed alternatives, 2,007 acres within acquisition parcels. This habitat was described in detail in the Grand Bay National Wildlife Refuge Comprehensive Conservation Plan (USFWS 2008) and the Grand Bay NERR Management Plan (MDMR 1998). This habitat is comprised of estuarine marsh, tidal marsh, and intertidal marsh.

<u>Savannas and Flatwoods</u>: Approximately 6,276 acres of savannas and flatwoods exists in the project area for the proposed alternatives, 3,535 within acquisition parcels. This habitat was described in detail in the Grand Bay National Wildlife Refuge Comprehensive Conservation Plan (USFWS 2008) and the Grand Bay NERR Management Plan (MDMR 1998). This habitat is comprised of the following habitat types: pine savannas and flatwoods, mesic pine savanna, wet pine savanna, mesic pine flatwoods, pond cypress savannas, and maritime forest.

<u>Freshwater Marsh</u>: Approximately 1,937 acres of freshwater marsh exists in the project area for the proposed alternatives, 1,207 acres within acquisition parcels. This habitat was described in detail in the Grand Bay NERR Management Plan (MDMR 1998).

Beach: Approximately 21 acres of beach exists in the project area for the proposed alternatives.

Open Water: Approximately 7,189 acres of open water exists in the project area for the proposed alternatives.

Invasive Species EO 13112 applies to all federal agencies whose actions may affect the status of invasive species, requires agencies to identify such actions, and to the extent practicable and permitted by law, requires agencies to 1.) take actions specified in the Order to address the problem consistent with their authorities and budgetary resources and 2.) not authorize, fund, or carry out actions that they believe are likely to cause or promote the introduction or spread of invasive species in the United States or elsewhere unless pursuant to guidelines that it has prescribed, the agency has determined and made public its determination that the benefits of such actions clearly outweigh the potential harm caused by invasive species; and that all feasible and prudent measures to minimize risk of harm will be taken in conjunction with the actions. The proposed alternative habitat management

is primarily invasive species management with restoration actions and measures including chemical treatment, mechanical treatment and prescribed fire. Best practices that would be used to control or eliminate invasive species are discussed in the environmental consequences section below.




Figure 3.4-4: Grand Bay Land Acquisition and Habitat Management - Habitats

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred) A summary of proposed restoration activities and their potential adverse and beneficial impacts are listed in Table 3.4-8 and discussed in this section.

Table 3.4-8: Proposed Alternatives-Habitat Impacts

Table 3.4-8: Proposed Alternative	s-manitat mipacis			
Habitats	Acquisition/Preservation	Chemical treatment	Mechanical treatment	Prescribed fire
Forested Freshwater Scrub-Shrub				
Adverse Impact Duration		short-term	short-term	
Adverse Impact Intensity		minor	minor	
Beneficial Impact Duration	long-term	long-term	long-term	
Coastal Marsh				
Adverse Impact Duration		short-term	short-term	
Adverse Impact Intensity		minor	minor	
Beneficial Impact Duration	long-term	long-term	long-term	
Savannas and Flatwoods				
Adverse Impact Duration		short-term	short-term	short-term
Adverse Impact Intensity		minor	minor to moderate	minor to moderate
Beneficial Impact Duration	long-term	long-term	long-term	long-term
Freshwater Marsh				
Adverse Impact Duration		short-term	short-term	
Adverse Impact Intensity		minor	minor	
Beneficial Impact Duration	long-term	long-term	long-term	

<u>Acquisition and Preservation:</u> There would long-term benefits to acquiring and preserving habitats; see table 3.4-8. Benefits would be applicable to proposed Alternative B and D.

<u>Chemical Treatment:</u> For all habitats, chemical treatment would be in small areas. There would be short-term minor impacts associated with accessing habitats and, if applicable, short-term impacts from any accidental spills. Care would be taken to obtain permits and handle chemicals as per manufactures instruction, particularly in aquatic systems. There would be long-term benefits from chemical treatment including control, prevention or elimination of Cogon grass, Chinese tallow, privet, Japanese climbing fern and other nuisance species and the resulting increase in diversity of native flora. Chemical treatment may be applied in combination with mechanical treatment and prescribed fire (discussed below). Impacts and benefits would be applicable to proposed Alternative B and D; see Table 3.4-8.

Mechanical Treatment: In forested freshwater scrub-shrub, coastal marsh, and freshwater marsh, mechanical clearing activities would likely be limited to clearing by hand or with small tools such as chainsaws. Physical disturbance from site access and dragging of vegetation, etc. would result in short-term, minor impacts. There would be a long-term benefit from mechanical treatment including control, eradication or prevention of the spread of nuisance species including Chinese tallow, privet, and other woody shrubs/invasive species; long-term benefits would also include a resulting increase in diversity of plant community flora.

For savanna and flatwoods, mechanical treatment activities include but would not be limited to use of brush-hog, use of chainsaws, use of gyro-tracs and in some cases bobcats or bulldozers to lay down or remove vegetation. These treatments could be used alone or in combination and also in preparation for prescribed fire. These would be short-term, minor to moderate impacts depending on the sizes of the treatment and intensity of treatment needed. There would be long-term benefits including increased diversity of flora, once nuisance species are controlled, eradicated or prevention measures are underway.

Mechanical treatment in wetter savanna and flatwoods would be done in a manner that would minimize impacts to soil to the extent practicable. In wet areas, soft track or wide track equipment would be used to distribute the equipment weight and minimize impact. Alternatively, crews may remove material with chainsaws. If required, a U.S. Army Corps of Engineers permit would be obtained likely a Nationwide 27 (Aquatic Habitat Restoration, Establishment, and Enhancement Activities), as well as a MDMR Coastal Wetlands Permit (if required). Nationwide 27 allows for mechanized land clearing to remove non-native invasive, exotic, or nuisance vegetation and other related activities. If there is any clearing within wetlands or stream boundaries, damage to vegetation, soil compaction and any resulting erosion could have a short-term, minor to moderate impact to wetlands. USACE permit and/or MDMR Coastal Wetlands permit conditions (if required) would be adhered to in all operations. Impacts from mechanical treatment would be applicable to proposed Alternative C and D.

Prescribed fire: Prescribed fire would apply to up to 6,276 acres of savanna and flatwoods. The preferred prescribed fire regime would be completed on a two-year rotation, with 50% of the prescribed burns occurring during the growing season, if possible given that weather conditions, seasonal wetness, availability of trained staff, invasive species presence and other factors are considerations in maintaining the fire frequency. These activities would largely be applied in areas that were colonized by woody invasive and understory shrubs such as gallberry (*Ilex glabra*), privet, saw palmetto, Chinese tallow, and other species. Impacts to soils and wetland were discussed in previous section. Prescribed fire could result in short-term, minor to moderate, adverse impacts, to existing habitats depending on the size of the operation. There would be long-term benefits to savanna and flatwoods from prescribed fire by creating conditions that would result in the reestablishment of diverse plant communities. There could also be incidental burning of freshwater marsh, when prescribed fire escapes during burning of adjacent habitats. These are periodic, unplanned occurrences. Resource managers typically allow the burns to spread through the marsh. Impacts would be applicable to proposed Alternative C and D.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to reduce the spread of invasive species:

 Prior to bringing any equipment (including personal gear, machinery, vehicles, or vessels) to the work site, inspect each item for mud or soil, seeds, and vegetation. If present, clean the equipment, vehicles, or personal gear until they are free from mud, soil, seeds, and vegetation. • Inspect the equipment, vehicles, and personal gear each time they are being prepared to go to a site or prior to transferring between sites to avoid spreading exotic, nuisance species.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. Under alternatives B and D, if development were to occur, there would likely be adverse impacts to habitats including habitat removal and/or fragmentation. If the parcels were to be developed, the habitats would be altered or removed completely constituting an adverse impact. When compared to Alternatives C and D, the No Action alternative would not provide the habitat benefits associated with management activities. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.3.2 Protected Species

Affected Environment

Section 3.6 of the PDARP/PEIS discusses biota of the northern Gulf of Mexico. This section covers endangered species in the Grand Bay Land Acquisition and Habitat Management project area for the proposed alternatives. The USFWS lists species as threatened or endangered when they meet criteria detailed under the Endangered Species Act (ESA) of 1973, as amended (16 U.S.C. §1531 et seq.). Additionally, Mississippi Wildlife Fisheries and Parks (MWFP) and NOAA National Marine Fisheries Service (NMFS) identify and list protected species. Section 7(a) (2) of the ESA requires that each federal agency ensure that any action authorized, funded, or carried out by the agency is not likely to jeopardize the continued existence of any endangered or threatened species or result in the destruction or adverse modification of Critical Habitat of those species. When the action of a federal agency may affect a protected species or its Critical Habitat, that agency is required to consult with either the NMFS or the USFWS, depending upon the protected species that may be affected. ESA Section 7 coordination is underway and the appropriate recommendations would be incorporated into the proposed project. Compliance with the Migratory Bird Treaty Act and Bald and Golden Eagle Protection Act are also discussed in this section.

Relevant federally protected species that are known to occur or could occur in Jackson County and that could occur in or near the project area for or could pass through the project area are listed in Table 3.4-9. A brief discussion of the state imperiled diamond back terrapin is also provided in the environmental consequences.

Table 3.4-9: Proposed Alternatives-Protected Species

able 5.4-9. I Toposed Afternatives-1 Totected Species								
Common Name	Scientific Name	Federal Status	Habitat					
Birds								
Piping Plover	Charadrius melodus	Threatened	Beaches and mudflats in southeastern coastal areas. Critical Habitat, MS-15, exists in Jackson County					
Red Knot	Calidris canutus rufa	Threatened	Marine intertidal habitats including inlets, estuaries, and bays feeding in mud and sand flats on beaches and barrier islands					
Mississippi Sandhill Crane	Grus canadensis pulla	Endangered	Open wetland habitats surrounded by shrubs or trees. Critical Habitat has been established on and					

Common Name	Scientific Name	Federal Status	Habitat
			adjacent to the Mississippi Sandhill Crane National Wildlife Refuge (USFWS 2013).
Red-cockaded Woodpecker	Picoides borealis	Endangered	This species excavates nesting and roosting cavities in living pine trees, and is the only species known to do so exclusively. Cavities have been found in most species of southern pines, but longleaf pine appears to be the preferred species. Older, mature trees are selected for cavity excavation.
Wood Stork	Mycteria americana	Threatened	Freshwater and estuarine wetlands, primarily nesting in cypress or mangrove swamps. They feed in freshwater marshes, narrow tidal creeks, or flooded tidal pools. Particularly attractive feeding sites are depressions in marshes or swamps where fish become concentrated during periods of falling water levels.
Fishes			
Gulf Sturgeon/Critical Habitat	Acipenser oxyrinchus desotoi	Threatened	Migrates from large freshwater coastal rivers to brackish and marine coastal bays and estuaries. The Deer Island Subtidal Reef and the Grand Bay Intertidal and Subtidal Reef project components have structures within Critical Habitat Unit 8.
Mammals			
West Indian Manatee	Trichechus manatus	Endangered	Fresh and salt water in large coastal rivers, bays, bayous and estuaries
Louisiana Black Bear	Ursus americanus luteolus	Endangered	Habitat is bottomland hardwoods along some of the major river systems
Reptiles/Amphibians			
Hawksbill Sea Turtle	Eretmochelys imbricata	Endangered	Coral reefs, open ocean, bays, estuaries
Leatherback Sea Turtle	Dermochelys coriacea	Endangered	Open ocean, coastal waters
Kemp's ridley Sea Turtle	Lepidochelys kempii	Endangered	Nearshore and inshore coastal waters, often in salt marshes; neritic zones with muddy or sandy substrate (NOAA Fisheries 2014b)
Green Sea Turtle	Chelonia mydas	Threatened	Shallow coastal waters with SAVs and algae, nests on open beaches
Loggerhead Sea Turtle	Caretta coretta	Threatened	Open ocean; also, inshore areas, bays, salt marshes, ship channels and mouths of large rivers
Alabama Red-belly Turtle	Pseudemys alabamensis	Endangered	Fresh and brackish habitats, river banks, submerged and emergent aquatic vegetation; upland habitat for nesting (MDWFP 2001; USFWS 2013)
Black Pinesnake	Pituophis melanoleucus lodingi	Threatened	uplands with well-drained sandy soils in areas of longleaf pine and hardwood tree species (USFWS 2013).
Gopher Tortoise	Gopherus polyphemus	Threatened	Well-drained, sandy soils, which allow easy burrowing; an abundance of diverse herbaceous ground cover; and an open canopy and sparse shrub cover, which allows sunlight to reach the ground floor (USFWS 2013)
Plants			
Louisiana Quillwort	Isoetes louisianensis	Endangered	Mineral soil, usually light gray in color, in bottomlands that are periodically washed free of leaves and debris. Streams along which quillworts grow may have flow year-round.

Birds

Mississippi Sandhill Crane: The Mississippi Sandhill crane utilizes open pine savanna and wetland habitats. Critical Habitat has been established on and adjacent to the Mississippi Sandhill Crane National Wildlife Refuge (USFWS 2013).

Piping Plover (*Charadrius melodus*): The piping plover does not nest in Mississippi; however, this species uses Gulf Coast beaches and barrier islands for wintering (MDWFP 2001). Plovers use sparsely vegetated sand beaches, mudflats, and salt marshes for roosting and foraging.

Red Knot (*Calidris canutus rufa*): In coastal Mississippi, the red knot is mainly a migratory species that uses coastal beaches and marine intertidal areas as stopover feeding locations or staging areas on the way to and from their wintering grounds in South America and breeding areas in the Arctic. Foraging on ocean beaches, mud and sand flats, and salt marshes occurs from March to April during the northward spring migration and September and October during the southward autumn migration (Niles et al. 2007; USFWS 2013). Red knots have been observed wintering on the Gulf Coast and are observed from October to March (USFWS 2013). The nonbreeding diet of this species includes marine invertebrates such as snails, crustaceans, and small mollusks including the coquina clam (*Donax variabilis*), which is common on Gulf coast beaches, and the dwarf surf clam (*Mulinia lateralis*) (Niles et al. 2007; USFWS 2013). Roosting and resting habitat includes areas above the high tide line such as reefs and high sand flats (USFWS 2013).

Red-cockaded Woodpecker (Picoides borealis): In Mississippi, this species has been recorded primarily from the southern two-thirds of the state. It has not been found in the Delta and only sporadically occurs in the northern counties. The Red-cockaded Woodpecker is a species of southern pine forests. The preferred nesting habitat is open, park-like, mature pine woodlands with few or no hardwood trees present. Preferred feeding habitats are pine stands with trees 23 cm (9 in.) and greater in diameter. These may or may not include a significant hardwood component. The Red-cockaded woodpecker excavates nesting and roosting cavities in living pine trees, and is the only species known to do so exclusively. Cavities have been found in most species of southern pines, but longleaf pine (*Pinus palustris*) appears to be the preferred species. Older, mature trees are selected for cavity excavation (MS Museum of Natural Science 2014).

Wood Stork (*Mycteria americana Linnaeus*): In Mississippi, wood storks have been observed most frequently along the western edge of the state in those counties bordering the Mississippi River and with increasing frequency in some counties along the eastern edge of the state, although they may occur almost anywhere there are sloughs or swamps to provide feeding habitat. The Wood Stork occurs primarily in freshwater wetlands, including ponds, bayheads, flooded pastures, oxbow lakes, and ditches. Nesting usually occurs in baldcypress trees in swamps, although breeding has also been observed in mangroves (MS Museum of Natural Science 2014).

Fishes

Gulf sturgeon (*Acipenser oxyrinchus desotoi*): This anadromous species migrates from coastal bays and estuaries to large coastal rivers in the spring for spawning and then returns to brackish and marine environments from October through March for foraging. The riverine spawning habitats for Gulf sturgeon in the State of Mississippi include the Mississippi, Pearl, and Pascagoula rivers (Ross et al. 2009; MDWFP 2001) but not the Biloxi and Tchoutacabouffa rivers (USFWS, GSMFC, and NMFS 1995; NMFS and USFWS 2009). The marine wintering areas where individuals have been observed

are nearshore and barrier island habitats from the Pearl River east to the barrier islands (Ross et al. 2009). Winter habitat is mainly around Cat, Ship, Horn, and Petit Bois islands with nearshore observations likely due to migratory movements to and from these offshore islands (Rogillio et al. 2007; Ross et al. 2009). The coastal Mississippi Sound waters of the State of Mississippi are designated as Critical Habitat.

Gulf Sturgeon Designated Critical Habitat: The project area extends into Gulf sturgeon Critical Habitat in Mississippi coastal waters and near the shoreline (Unit 8-Lake Ponchartrain-Mississippi Sound). Critical Habitat was designated in 2003 by the National Marine Fisheries Service (NMFS) and was based on seven primary constituent elements (PCEs) essential for its conservation. None the restoration activities would be in open water. Therefore, the proposed alternative is expected to have No Effect on Gulf sturgeon and consultation with the U.S. Fish and Wildlife Service will not be requested.

Mammals

West Indian Manatee (*Trichechus manatus*): This species uses both fresh and saltwater habitats such as coastal rivers, bays, bayous, and estuaries. The manatee is an occasional visitor to Mississippi's coasts, although migration into the area is poorly understood. After wintering in Florida, and perhaps Mexico, manatees migrate northward during spring, including to Mississippi and Alabama waters, although these migrations are not well understood (Fertl et al. 2005). Manatees frequently seek out freshwater sources such as rivers and river mouths and have been known to be found near estuaries (Fertl et al. 2005). SAVs are the typical manatee forage material; however, manatees can also consume other aquatic vegetation, algae, and terrestrial vegetation (Fertl et al. 2005). None of the restoration activities would be in open water.

Louisiana Black Bear (Ursus americana luteolus): The Mississippi Department of Wildlife, Fisheries, and Parks (MDWFP) estimates the Mississippi population of Louisiana black bear to be around 50 animals. Most of the bears observed in Mississippi are believed to be males that have traveled from other states; only one was reported sighted in Hancock County from 1996 – 2006 (Young 2006). This sighting was in northern Hancock County in the Pearl River drainage system. Large contiguous bottomland forest habitat is preferred by the species and does exist adjacent to the proposed project elements. However, the bears typically prefer larger tracts of bottomland forest with no human disturbance and having good cover (Young 2006). The proposed project areas do not have hardwood forest that is preferred by Louisiana black bear. There is no known breeding population of bears in this area, and any presence would likely be transitory animals following a river corridor for foraging and cover.

Reptiles

Hawksbill Sea Turtle (*Eretmochelys imbricata*): Although this species uses various habitats such as the open ocean, bays, and estuaries throughout different life stages, it is mainly associated with coral reefs. This species nests in Florida from April to November (NOAA Fisheries 2014a). It likely does not nest in Mississippi and observations are rare in the state (MDWFP 2001; NOAA Fisheries 2014a). The main dietary items of this species are sponges and other invertebrates (NOAA Fisheries 2014a).

Leatherback Sea Turtle (*Dermochelys coriacea*): This species mainly inhabits the offshore open ocean; however, it does use nearshore coastal waters during nesting or feeding. Nesting for this species occurs in Florida from April through November. Their main forage item is jellyfish. This

species migrates long distances from nesting to feeding areas. While not common, there have been sporadic observations of leatherback sea turtles in Mississippi waters (MDWFP 2001).

Kemp's ridley Sea Turtle (*Lepidochelys kempii*): Typical habitat for this species includes nearshore and inshore coastal waters and often salt marshes and neritic zones with muddy or sandy substrate (NOAA Fisheries 2013b). This species has been observed in nearshore waters of the Mississippi Sound during migration and foraging and has been accidentally caught by shore-based fishermen (MDWFP 2001; Shaver and Rubio 2008). Females typically nest from May through July (NOAA Fisheries 2014b). Males potentially use Gulf of Mexico habitats all year and females presumably use the Mississippi Sound and barrier island habitats for foraging when not nesting (NOAA Fisheries 2014b). Kemp's ridley sea turtles do not nest in Mississippi (MDWFP 2001).

Green Sea Turtle (*Chelonia mydas*): This species typically prefers shallow coastal waters with SAVs and algae for foraging and nests on open beaches (NOAA Fisheries 2015). Nesting typically does not occur on mainland beaches and there is likely no Mississippi nesting at all (MDWFP 2001; NOAA Fisheries 2015). This species migrates long distances in the open ocean from nesting to feeding areas. Observations of this species in Mississippi are rare (MDWFP 2001).

Loggerhead Sea Turtle (*Caretta caretta*): Loggerhead habitat for foraging and migration includes open ocean, inshore areas, bays, salt marshes, ship channels, and mouths of large rivers. This sea turtle feeds on mollusks, fish, crustaceans, and other marine organisms. This species typically nests at night from late April through September (NOAA Fisheries 2014c). Although loggerheads occasionally use barrier islands for nesting, mainland nesting is rare (MDWFP 2001). Preferences for nesting beaches include high-energy coarse-grained beaches adjacent to the ocean that are narrow and steeply sloped (NOAA Fisheries 2014c). This species has been observed in nearshore waters of the Mississippi Sound during migration and foraging and has been accidentally caught by shore-based fishermen (MDWFP 2001).

Alabama Red-Belly Turtle (*Pseudemys alabamensis*): The habitat of the Alabama red-belly turtle includes fresh and brackish habitats, river banks, submerged and emergent aquatic vegetation, and upland habitat for nesting (MDWFP 2001; USFWS 2013). This species is mainly a freshwater species associated with river and stream channels and associated wetlands. Nesting occurs from mid-May to mid-July (MDWFP 2001).

Black Pine Snake (*Pituophis melanoleucus lodingi*): Suitable habitat includes open canopy longleaf pine forest with herbaceous ground cover and well-drained sandy soils and, less so, hardwood forests (USFWS 2010).

Gopher Tortoise (*Gopherus polyphemus*): The Gopher Tortoise uses well-drained to excessively well-drained upland soils. Tortoises require soils that are sandy enough to permit construction of burrows and open canopies that allow sufficient herbaceous plant growth and sunny areas in which to nest. In Mississippi, these areas often support a mixture of longleaf pine and scrub oaks.

Plants

Louisiana Quillwort (*Isoetes louisianensis*): The Louisiana Quillwort has been observed in 10 counties in 174 streams within 17 watersheds (USFWS 2012a) throughout the State of Mississippi with the largest colony found in the DeSoto National Forest (USFWS 2012a). This species is found in all three coastal Mississippi counties (MDWFP 2001; USFWS 2012a) although none have been found

near the proposed project area (MDWFP 2001). In coastal Mississippi, Louisiana Quillwort habitat includes perennial streams and banks in bottomland hardwood habitats likely with bald cypress and possibly the presence of stream macrophytes such as *Sparganium* spp. and *Orontium* spp. (USFWS 2012a). Earlier sources indicate that suitable habitat for this species consists of sand or gravel bars located in intermittent streams and associated riparian areas (MDWFP 2001). Louisiana Quillworts are sensitive to changes in hydrology, sedimentation, and alterations to the surrounding overstory (USFWS 2012a).

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred) PDARP programmatic ESA consultations were developed with the National Marine Fisheries Services (NMFS 2016) and the U.S. Fish and Wildlife Service (USFWS 2016). Potential impacts to threatened or endangered species and their Critical Habitat are presented in Table 3.4-22. The MS TIG has begun coordination under the programmatic ESA consultations. The project area in the southeast is adjacent to the Mississippi sound which is designated Critical Habitat for Gulf sturgeon. None of the restoration activities would be completed in open water. Thus, there would be no effect as a result of any restoration activity to in water species (and associated Critical Habitat), including Gulf sturgeon, West Indian manatee, and sea turtles; for this reason, they are not included in the environmental consequences discussion in Table 3.4-10.

Table 3.4-10: Proposed Alternatives-Protected Species Impacts

Species /Critical Habitat	Applicable Habitats	Restoration Activities for Applicable Habitats	Potential Impacts to Species/Critical Habitat
Alabama Red-Belly Turtle (Pseudemys alabamensis)	Freshwater Marsh Savannas and flatwoods	Acquisition/Preservation Chemical/Mechanical Treatment Prescribed Fire	For areas that have potential for occurrence of the species, the site would be surveyed and species, if present, would be avoided in the planning and implementation of the activity.
Piping plover (Charadrius melodus) and red knot (Calidris canutus rufa)	Beach	Access Restriction	Acquisition and preservation are the only measure planned for the beach. No adverse impacts to piping plover are anticipated.
Black pine snake (Pituophis melanoleucus lodingi)	Savanna and Flatwoods	Acquisition/Preservation Chemical treatment Mechanical treatment of undesirable vegetation Prescribed fire	If habitat exists prescribed fire and mechanical clearing of upland areas may affect species habitat. Surveys should be conducted in areas where the species is likely to occur. Survey results would be considered in the design of the management and or restoration measures to either avoid or minimize impacts to the species.
Gopher tortoise (Gopherus polyphemus)	Savannas and Flatwoods	Acquisition/Preservation Chemical treatment Mechanical treatment Prescribed fire	Prescribed fire and mechanical clearing of upland areas may affect species habitat. Surveys would be conducted in areas where the gopher tortoise is likely to occur. Survey results would be considered in the design of the management and or restoration measures to either avoid or minimize impacts to the species.
Louisiana quillwort (Isoetes louisianensis)	Savanna flatwoods, Forested freshwater scrub- shrub	Acquisition/Preservation Chemical treatment Mechanical treatment Prescribed fire	Chemical treatment, mechanical treatment, could result in an impact to vegetation. Restoration activity areas that are likely to contain the species would be surveyed; if the species was found it would be avoided in the implementation of restoration measures and activities.
Mississippi Sandhill Crane (<i>Grus</i> canadensis pulla)	Savanna flatwoods, Forested freshwater scrub-	Acquisition/Preservation Chemical treatment Mechanical treatment Prescribed fire	Chemical treatment, mechanical treatment, prescribed fire. Noise impact causing the species to temporarily vacate the area. The species could return after restoration activities have ceased.

Species /Critical Habitat	Applicable Habitats	Restoration Activities for Applicable Habitats	Potential Impacts to Species/Critical Habitat
	shrub		
Mississippi diamondback terrapin (Malaclemys terrapin pileata)	Beach	Access Restriction	There are no restoration activities planned for beach habitat other than access restriction; no adverse impacts are anticipated.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The MS TIG would continue to consult with the appropriate regulatory agency to further avoid or minimize impacts to these species in the planning site-specific restoration activities and management measures. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to protected species.

Gopher Tortoise

• If suitable habitat is present, coordinate with the local USFWS Ecological Services Field Office to discuss the need for surveys to identify any gopher tortoise burrows and to develop conservation measures to avoid or minimize impacts. Measures could include establishing a protective buffer (size determined by USFWS and the state trust resource agency) if burrows are within the project area and cannot be avoided, implementing standard procedures to relocate the tortoise within the project site but away from the areas of restoration.

Protected Plants

- If suitable habitat is present, coordinate with the local USFWS Ecological Services Field Office to discuss the need for surveys to identify protected plants and to develop conservation measures to avoid or minimize impacts.
- Enhance and protect plants on site and in adjacent habitats to the maximum extent possible.

Protected Species

• Provide all individuals working on a project with information in support of general awareness of and means to avoid impacts to protected species and their habitats present at the specific project site.

ESA Section 7 coordination is underway and the appropriate recommendations would be incorporated into the proposed site-specific restoration activities and management measures as applicable. Because there is no in-water work, no effects to manatee are expected, and the Implementing Trustees determined that no take of manatee under ESA or MMPA would occur.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. Under Alternatives B and

D, if development were to occur, there would likely be adverse impacts to habitat that could be utilized by protected species. Habitats that protected species could use would not be protected from development under the No Action alternative when compared to Alternatives B and D; however, no impacts would occur to protected species or designated critical habitats without conducting required consultations. There would be no benefits to habitat from management activities that would be provided under Alternatives C and D. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.3.3 Migratory Birds

Affected Environment

Migratory bird species groups that could occur in the proposed alternatives project area include wading birds, shorebirds, seabirds, raptors, goatsuckers, waterfowl, doves and pigeons, and rails and coots (Table 3.4-11).

Table 3.4-11: Migratory Bird Species Groups Present in Project Area and Example Behaviors

SPECIES	BEHAVIOR	SPECIES/HABITAT IMPACTS
Wading birds (herons, egrets, ibises)	Foraging, feeding, resting, roosting, nesting	Wading birds primarily forage and feed at the water's edge. As such, they may be impacted locally and temporarily by the project. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. These birds primarily nest and roost in trees or shrubs (e.g., pines, <i>Baccharis</i>), and could utilize areas that will be managed by mechanical treatment and prescribed fire.
Shorebirds (plovers, oystercatchers, stilts, sandpipers)	Foraging, feeding, resting, roosting, nesting	Shorebirds forage, feed, rest, and roost in the action area. As such, they may be impacted locally and temporarily by the project. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. In the project area, these birds would primarily nest on beaches. There are no planned activities near shorebird nesting habitats.
Seabirds (terns, gulls, skimmers, double-crested cormorant, American white pelican, brown pelican)	Foraging, feeding, resting, roosting	Seabirds forage, feed, rest, and roost in the action area. Terns and skimmers could utilize the beach habitat in the project area. As such, they may be impacted locally and temporarily by the project. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. There are no activities planned near seabird nesting habitat.
Raptors (osprey, hawks, eagles, owls)	Foraging, feeding, resting, roosting, nesting	Raptors forage, feed, rest and nest in the action area. As such, they may be impacted locally and temporarily by the project. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. There are Osprey nests in cleared pipeline rights-of-way and possibly in snags located near open water. Chemical treatment, mechanical treatment and prescribed fire could be completed in the vicinity of raptor nests. Surveys would be conducted before commencing restoration activities.
Goatsuckers	Foraging, feeding, resting, roosting, nesting	Goatsuckers forage, feed, rest, and roost in the action area. However, they are nocturnal/crepuscular and therefore not active during the project work period. They nest in thickets and woodlands.
Waterfowl (ducks, loons, and grebes)	Foraging, feeding, resting, roosting, nesting	Waterfowl forage, feed, rest, and roost in the action area. As such, they may be impacted locally and temporarily by the project. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. These birds primarily roost and nest in low vegetation. There would be no restoration activities in open water or estuarine marsh.
Doves and pigeons	Foraging, feeding, resting, roosting	Doves and pigeons could forage, feed, rest, and roost in the action area. However, they are unlikely to utilize habitat in the savanna flatwood area.

SPECIES	BEHAVIOR	SPECIES/HABITAT IMPACTS
Rails and coots	Foraging, feeding, resting, roosting, nesting	Rails and coots forage, feed, rest, and roost in the action area. As such, they may be impacted locally and temporarily by the project. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting if disturbed by the project. These birds primarily roost and nest in marshes, which are within the action area. There would be no in-water restoration activities where these species nest.

Migratory Bird Treaty Act: The Migratory Bird Treaty Act of 1918 (MBTA) implements various treaties and conventions among the United States, Canada, Japan, Mexico, and the former Soviet Union for the protection of migratory birds. Under MBTA, unless permitted by regulations, it is unlawful to pursue; hunt; take; capture or kill; attempt to take, capture, or kill; possess; offer to sell or sell; barter; purchase; deliver; or cause to be shipped, exported, imported, transported, carried, or received any migratory bird, part, nest, egg, or product, manufactured or not. USFWS regulations broadly define "take" under MBTA to mean "pursue, hunt, shoot, wound, kill, trap, capture, or collect, or attempt to pursue, hunt, shoot, wound, kill, trap, capture, or collect".

Bald and Golden Eagle Protection Act: The Bald and Golden Eagle Protection Act (BGEPA) of 1940 (16 U.S.C. 668-668c) prohibits anyone, without a permit issued by the Secretary of the Interior, from "taking" bald eagles, including their parts, nests, or eggs. BGEPA provides criminal penalties for persons who "take, possess, sell, purchase, barter, offer to sell, purchase or barter, transport, export or import, at any time or any manner, any bald eagle ... [or any golden eagle], alive or dead, or any part, nest, or egg thereof." Golden eagles are not present along the Gulf Coast.

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred) Migratory birds could use areas at and around the project area for foraging, feeding, resting, and nesting. Nesting species include raptors, wading birds, marsh birds, waterfowl and shorebirds; Table 3.4-23. For all planned restoration activities, pre-commencement nesting surveys for migratory birds and raptors within the restoration activity area would be conducted and if evidence of nesting is found, resource managers would coordinate with the USFWS to develop and implement appropriate conservation measures, such as those described below. Due to the implementation of best management practices no "take" of nesting birds is anticipated. There are no golden eagles in the project footprint. Raptor nest surveys would be completed within the restoration activity area where raptor nesting habitat exists. If evidence of nesting is found, resource managers coordinate with the USFWS to develop and implement appropriate conservation measures, therefore no impacts to golden or bald eagles are anticipated. Potential adverse effects to birds include elevated noise levels due to the use of mechanical equipment for vegetation clearing, and from noise and smoke during prescribed burning. These species are mobile and would likely exit the area during management activities (no impacts to overall population). Foraging and resting birds may temporarily be displaced during management activities. Bird roosting would not be affected because management activities would occur during daylight hours. Therefore, impacts are expected to be short-term, localized, and minor.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to migratory bird species including bald eagles:

Migratory Birds

- Use care to avoid birds when operating machinery or vehicles near birds.
- Avoid working in migratory bird nesting habitats during breeding, nesting, and fledging
 (approximately mid-February through late August). If project activities must occur during
 this timeframe and breeding, nesting, or fledging birds are present, contact the state trust
 resource agency to obtain the most recent guidance to protect nesting birds or rookeries, and
 their recommendations will be implemented.
- Conservation areas may already be marked to protect bird nesting areas. Stay out of existing marked areas.
- If vegetation clearing is necessary, clear vegetation outside the migratory bird nesting season (approximately mid-February through late August) or have a qualified biologist inspect for active nests. If no active nests are found, vegetation may be removed. If active nests are found, vegetation may be removed after the nest successfully fledges.

Bald Eagles

- If bald eagle breeding or nesting behaviors are observed or a nest is discovered or known, have all activities avoid the nest by a minimum of 660 feet. If the nest is protected by a vegetated buffer where there is no line of sight to the nest, then the minimum avoidance distance is 330 feet. Maintain this avoidance distance from the onset of breeding/courtship behaviors until any eggs have hatched and eaglets have fledged (approximately 6 months).
- If a similar activity (such as driving on a roadway) is closer than 660 feet to a nest, maintain a distance buffer as close to the nest as the existing tolerated activity. If a vegetated buffer is present and there is no line of sight to the nest and a similar activity is closer than 330 feet to a nest, then maintain a distance buffer as close to the nest as the existing tolerated activity.
- In some instances, activities conducted within 660 feet of a nest may result in disturbance, particularly for the eagles occupying the Mississippi barrier islands. If an activity appears to cause initial disturbance, stop the activity and move all individuals and equipment away until the eagles are no longer displaying disturbance behaviors. Contact the USFWS's Migratory Bird Permit Office to determine how to avoid impacts or if a permit may be needed.

The MS TIG has begun coordination and review of the project for impacts to bald eagles and migratory birds in accordance with the Bald and Golden Eagle Protection Act (BGEPA) of 1940 (16 U.S.C. 668-668c) and the Migratory Bird Treaty Act (MBTA) of 1918 (16 U.S.C. 703–712) to ensure appropriate conservation measures and best practices would be incorporated into the project.

No Action

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. Habitats that these species could use would not be as protected from development under the No Action alternative when compared to Alternatives B and D. However, under Alternatives B and D, even if development were to occur, migratory birds and bald/golden eagles would still be protected under federal statute.

Enhancements to potential habitat that these species utilize would not take place under the No Action alternative when compared to Alternatives C and D. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.3.4 Wildlife

Affected Environment

Section 3.6 of the PDARP/PEIS discusses the biota of the northern Gulf of Mexico. For the proposed alternative project area, the Grand Bay National Wildlife Refuge Comprehensive Conservation Plan (USFWS, 2008) is incorporated by reference. That plan provides a discussion of a number of species including grassland birds, migratory birds, waterfowl, marshbirds, landbirds, amphipians, reptiles, and the Mississippi sandhill crane. Goals and objectives for these species within that plan include:

- Grassland birds: providing pine savanna habitat for the benefit of these species;
- Other migratory birds: improving knowledge base for management by increasing baseline knowledge of the distribution, abundance and use of the refuge by a variety of birds, including waterfowl, marsh birds, and landbirds;
- Amphibians and reptiles: continuing monitoring their presence through surveys and considering projects that might benefit their populations while pursuing primary Mississippi sandhill crane-oriented goals and objectives of refuge;
- Fire management: proactively using prescribed fire for habitat management and fuel reduction objectives in a rapidly developing area with ever more constraints that must be observed by fire managers;
- Manage and protect migratory birds;
- Achieve goals (savanna restoration, fire, roll chopping, etc.) to meet refuge purpose of establishing breeding pairs of Mississippi sandhill cranes;
- After fire, conduct migratory bird surveys in savanna.

Management actions to achieve the goals and objectives are also outlined in the Grand Bay NWR Comprehensive Conservation Plan.

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred) Table 3.4-12 summarizes the environmental consequences to wildlife from the proposed alternatives. A discussion is provided below.

Table 3.4-12: Summarized proposed alternative impacts on wildlife

Restoration Measure	Alternative B: Grand Bay Land Acquisition			Alternative C: Grand Bay Habitat Management			Alternative D (Preferred): Grand Bay Land Acquisition and Habitat Management		
	Adverse	Adverse	Beneficial	Adverse	Adverse	Benefici	Adverse	Adverse	Beneficial
	Impact	Impact	Impact	Impact	Impact	al Impact	Impact	Impact	Impact
	Duration	Intensity		Duration	Intensity		Duration	Intensity	
Acquisition/									
Preservation			long-term						long-term
Chemical				short-	minor to	long-	short-	minor to	
Treatment				term	moderate	term	term	moderate	long-term
Mechanical				short-	minor to	long-	short-	minor to	
Treatment				term	moderate	term	term	moderate	long-term
Prescribed Fire				short-	minor to	long-	short-	minor to	
				term	moderate	term	term	moderate	long-term

<u>Acquisition/Preservation</u>: Prevention of development of habitats would be a long-term, benefit to wildlife species that currently inhabit or transiently utilize the preserved habitats. Impacts would be applicable to proposed Alternative B and D.

Chemical and Mechanical Treatment/Prescribed Fire: Chemical treatment is often used in combination with fire or mechanical clearing. Invasive species management approaches would result in a short-term, minor to moderate impacts to wildlife species in and near treatment areas due to equipment noise, mechanical clearing, exposure to chemicals and prescribed burning. treatment and prescribed fire would be the most intrusive; however, these techniques would be applied to areas that have dense woody shrub layers which preclude utilization by a number of bird and mammal species. There would be a short term, minor to moderate impact to species in the area during mechanical treatment and prescribed fire. Many species would leave the area during the operations, but would likely return to utilize the restored habitats. Mechanically treated and/or prescribed fire areas would become open habitat and be colonized with native pine savanna species over several seasons. These communities are one of the most diverse habitats and would result in increased diversity in insect, bird, and small mammal populations. Improved savanna and flatwoods would provide high quality habitat for grassland birds. Fire management applied to up to 6,276 acres of savannas and flatwoods would not only meet resource manager fuel reduction objectives, but would also enhance habitat for the Mississippi sandhill crane habitat, and benefit other migratory birds. Adverse and beneficial impacts from invasive species management treatments including chemical, mechanical and prescribed fire would be applicable to proposed Alternative C and D.

No Action

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. Under alternatives B and D, if development were to occur, there would likely be adverse impacts to habitats that wildlife species use. The development resulting from the No Action alternative could cause additional human disturbance, such as noise would increase with development and could cause adverse impacts to wildlife. Wildlife habitat would not be enhanced under the No Action alternative like it would in Alternatives C and D. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.4 Socioeconomic Environment

<u>Introduction to Affected Environment (Socioeconomic Resources):</u> The section provides a discussion of socioeconomic resources and environmental justice, tourism and recreation, cultural resources, land and marine management, and public health and safety. PDARP/PEIS Section 3.2 is incorporated by reference here.

<u>Programmatic Review of Environmental Consequences (Socioeconomic Resources)</u>: Sections 6.4.1.5 and 6.4.1.10 of the PDARP describe the impacts to Human Use and Socioeconomic Resources for the relevant restoration approaches and are incorporated by reference and briefly described here.

<u>PDARP/PEIS</u> consequences related to economic effects: Acquisition and preservation could have long-term, minor to moderate adverse economic effects if acquisition prevents or limits development. Acquisition could permanently limit the amount and type of development permitted, and the management and intensity of use on these properties would likely change. Ownership changes and/or permitted uses could affect property taxes and have broader regional economic impacts. Land acquisition could have a minor to moderate impact on socioeconomic resources due to changes in visitor spending and tax impacts. The transfer of fee title to lands are transactions negotiated or arranged between willing parties and, as such, are not expected to give rise to adverse socioeconomic impacts to those who choose to engage in such transactions.

PDARP/PEIS consequences related to recreation and tourism: The acquisition of lands to protect habitat could result in impacts to recreation and tourism opportunities depending on site-specific land management practices applied. Closures, such as fencing or other mechanisms to protect nest sites, could result in short-term (seasonal) prohibitions on public access. Restrictions on public access in areas where public access had previously been allowed could reduce recreational opportunities. Over the long term, these techniques could result in healthy populations and provide wildlife enthusiasts with increased wildlife viewing opportunities. Conservation or acquisition of natural land resources can have indirect benefits on fish and wildlife habitat, potentially resulting in increased fishing and hunting opportunities. Seasonal or permanent employment could increase in order to provide labor for the installation, maintenance, and implementation of management projects such as hunting or trapping. Minor, short-term adverse impacts could result due to restoration activities. However, improvements in habitat associated with this approach may draw additional visitors to the area with associated visitor spending, increasing sales and tax receipts on retail purchases.

<u>PDARP/PEIS</u> consequences related to cultural resources: Creating, enhancing, or restoring bird nesting habitat may result in minor (temporary disturbance) to moderate (disturbance without loss of cultural information) impacts on cultural and historic resources depending on the scale of the action and site-specific characteristics. Discovery or recovery of cultural or historic resources would allow their future protection.

As appropriate in a tiered analysis, the evaluation the proposed alternative focuses on the specific resources with a potential to be affected. Infrastructure, fisheries and aquaculture, marine transportation, aesthetics and visual resources would have negligible to minor adverse effects or would provide benefits. To avoid redundant or unnecessary information, a summary of environmental consequences for these resources is provided here.

Infrastructure: There would be no impact to infrastructure from land acquisition activities associated with Alternative B. Infrastructure on the site includes logging roads for timber management, gas pipelines and utility corridors. There could be short-term, minor adverse impacts to gas pipelines or utility corridors from activities (minor clearing, temporary crossings) associated with mechanical treatment and prescribed fire (Alternative C and D; preferred). Care would be take to identify utility corridors as part of project planning and prior to implementation or restoration measures.

Fisheries and Aquaculture: There would be limited low impact activities in open water or estuarine marsh. Alternatives B, C and D acquisition and restoration measures could benefit marine resources in Grand Bay project area. Alternative B, C and D could provide net reduction in sediment

movement resulting from preservation and restoration versus a development/build out scenario of lands proposed for acquisition.

Marine Transportation: There would be no restoration activity that would occur in open water; the proposed alternative would not have an impact on marine transportation.

Aesthetics and Visual Resources: There would be no impact from Alternative B, land acquisition. Prescribed fire would result in a change in viewshed (Alternatives C and D-preferred). There may be temporary short-term minor impacts as a result of smoke. The land may look scorched after a prescribed burn until vegetation regrows. Depending on weather conditions, burn units can revegetate ("green up") within days to weeks. Revegetation after burning would result in a viewshed of natural vegetation with increased diversity of flowering plants and fauna (Alternatives C and D-preferred).

For the socioeconomic environment, the following resources are further analyzed in this section:

- Socioeconomics and Environmental Justice
- Tourism and Recreational Use
- Cultural Resources
- Land and Marine Management
- Public Health and Safety

3.4.1.4.1 Socioeconomic Resources and Environmental Justice

Affected Environment

PDARP/PEIS Section 3.2 discusses socioeconomic resources of the Gulf Coast and is incorporated by reference here. The project area for the proposed alternatives is located within Jackson County, Mississippi. The Grand Bay NWR Comprehensive Conservation Plan summarizes the socioeconomic environment for the proposed alternatives and is incorporated here by reference (USFWS, 2008). Jackson County is three times more densely populated than the state (181 persons per square mile vs. 61 persons per square mile) and growing faster. In 2003, the county's estimated population was 133,928, about five percent of Mississippi's population of 2,881,281 (U.S. Census Bureau 2005). The county population grew by 1.9 percent from 2000 to 2003, compared to Mississippi's 1.3 percent growth in the same three years. From 1990 to 2000, Jackson County grew 14 percent compared to Mississippi's 10.5 percent in the same decade.

Over the last decade, residential and commercial development has been proceeding rapidly in the coastal portion of Jackson County, Mississippi, converting forest plantations and farm fields into developed lots with houses, businesses, and institutions. Open space and habitat are becoming more and more fragmented. This development is expected to continue over the foreseeable future, in part because of the desirability of living in a coastal county with beach and ocean

The affected environment includes portions of the populations of Census Tract 401.2, 413, 416, and 427; and 411, specifically the residents close to the Grand Bay. Census Tract 427 makes up most the population affected. Small portions of Census Tracts 413 and 401.2 are within the project area; and only a few parcels within Census Tract 413 are within the project area. The population of Jackson County was 139,668 in 2010 and accounted for 4.7% of the state's total population, while Census Tract 427 (population 1,016 in 2010) accounted for <1% of the county population (Table 3.4-13). In 2010, median household income in Jackson County was \$49,145, which was 25% higher than the median household income in the State of Mississippi (\$39,464). Median household income of

Census Tract 427 in 2014 was \$48,317, which is 1.6% lower than that of the county and 22% higher than the median household income of the state (U.S. Census Bureau, 2010-2014 American Community Survey 5-Year Estimates).

Table 3.4-13: Population data (http://www.census.gov/2010census/popmap/)

Topic	Mississippi		Jackson County		Census Tract 401.2		Census Tract 413		Census Tract 416		Census Tract 427	
2010 Total Population	2,967,297		139,668		7,569		6,504		2,557		1016	
White alone	1,754,684	59%	100,735	72%	7,328	97%	5,000	77%	294	11%	862	85%
Black or African American Alone	1,098,385	37%	30,034	22%	79	1%	1,322	20%	2,178	85%	122	12%
Asian alone	25,742	<1%	3023	2.2%	28	<1%	39	<1%	3	<1%	2	<1%
American Indian and Alaska Native alone	15,030	<1%	565	<1%	20	<1%	21	<1%	3	<1%	6	<1%
Native Hawaiian and Other Pacific Islander alone	1,187	<1%	79	<1%	2	<1%	5	<1%	0	0%	0	0%
Some Other Race alone	38,162	1.3%	2610	1.9%	45	<1%	38	<1%	44	1.7%	19	1.8%
Two or More Races	34,107	1.1%	2622	1.9%	67	1%	79	1.2%	35	1.4%	5	<1%

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred)

Acquisition and preservation of property in fee and the set-aside in perpetuity would permanently limit development (Alternative B). The change in ownership would affect property taxes paid to local governments and could result in a broader regional economic impact resulting from changes in visitor spending in the area. There could be minor increases in spending resulting from recreational access to the project area as it increases in size and opportunities to hike, view wildlife in the area, or attract recreation on the basis of eco-tourism in the region are enhanced. Land acquisition could have a minor to moderate impact on socioeconomic resources due to changes in visitor spending and tax impacts. The transfer of fee title to lands would be transactions negotiated or arranged between willing parties and, as such, are not expected to give rise to adverse socioeconomic impacts to those who choose to engage in such transactions. Executive Order 12898 directs federal agencies to make achieving environmental justice part of its mission by identifying and addressing, as appropriate, disproportionately high adverse human health or environmental effects of its activities on minority and low-income populations. There would be no disproportionate impacts on minority, low-income, or underserved populations from the implementation of proposed alternatives. Impacts would be applicable to proposed Alternative B and D.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes

described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. If development were to occur, there would likely be an increase in property taxes paid to local governments. There would be no benefits from additional recreational visitor spending that could result from implementation of Alternatives B, C or D. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.4.2 Tourism and Recreational Use

Affected Environment

The Grand Bay NWR Comprehensive Conservation Plan (USFWS 2008) provides an overview of tourism and recreational use on the NWR; information is incorporated here. The Grand Bay NWR receives about 700 visitors annually. Wildlife observation and photography, hunting (waterfowl, mourning doves, white-tailed deer, and feral hogs), and boating in tidal marshes are the managed recreational uses of Grand Bay NWR. All refuge roads open to the public are either paved or gravel. Bayou Heron Road and Pecan Road together are about 3 miles in length.

<u>Hunting</u>: Hunting for white-tailed deer, feral hogs, squirrel, geese, ducks, coots, and mourning doves on designated areas, subject to state regulations and conditions outlined by the refuge.

Ecotourism: Jackson County conducted the Pascagoula River Ecotourism Study in 2002–2003. The Gautier Economic Development Council formed an Ecotourism Planning Committee which published an "Ecotourism Master Plan" in 2004 (Gautier Economic Development Council 2004). This plan acknowledges Mississippi Sandhill Crane NWR as one of the premier local nature destinations that can attract tourists to the area for outdoor activities. Other local attractions are Shepard State Park (MDWFP), Pascagoula River Marsh (MDMR), Indian Point Campground and Recreational Vehicle Resort (privately owned), and Alf Dantzler Wildlife Preserve (MDMR).

<u>Boating/Fishing</u>: A public boat launch facility and bank fishing area is located at the end of Bayou Heron Road (USFWS 2004). A universally accessible fishing pier that is compliant with the Americans with Disabilities Act (ADA) is adjacent to the boat launch, along with a resurfaced ADA compliant gravel parking area. The refuge provides diverse habitats of salt marshes, bayous, grass beds, etc., for the region's important commercial and recreational species of fish. These habitats serve as nursery areas as well as breeding and feeding grounds for shrimp, red drum, speckled trout, blue crab, oysters, and crabs, among other marine and aquatic organisms.

<u>Wildlife Observation and Photography:</u> Grand Bay NWR provides limited opportunities for wildlife observation. Birding is one of the most popular forms of wildlife observation on the refuge, with viewing opportunities changing seasonally. Viewing opportunities include wintering flocks of wading birds and waterfowl in the bayou and bay, songbirds in the trees and shrubs, and harriers and hawks hunting over the savanna. Visitors may also see other common wildlife such as white-tailed deer, raccoon, snakes, and frogs.

<u>Hiking</u>: The Escatawpa Trail was developed in partnership with the Mississippi Interstate Welcome Center. The trail is a two-mile part boardwalk and part gravel trail. The trail features include universal access, and benches for resting and wildlife viewing opportunities. The trail provides

wildlife observation and photography opportunities, particularly at the Escatawpa River overlook. There is also a picnic pavilion near the trail entrance on land adjacent to the refuge.

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred)

Alternative B (Acquisition) would result in a long-term benefit to tourism and recreation opportunities by expanding the area's recreational activities including wildlife observation, hunting, boating, and hiking. Management activities such as chemical treatment, mechanical treatment, and prescribed fire would result in temporary access closure to parts of the proposed alternative project area but only during management activities. These would typically be done during the growing season and would be short in duration and would not preclude access from all parts of the NERR, NWR, or CP for most activities (Alternative B); short-term, minor, adverse impact to tourism and recreation would result. Proposed Alternative B and D would increase opportunities for recreation by increasing the area of publicly accessible lands resulting in a long-term benefit to tourism and recreation.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. Under alternatives B and D, if development were to occur, there would likely be adverse impacts to tourism or recreation since development would likely limit access to these properties for recreational purposes. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.4.3 Cultural Resources

Affected Environment

Cultural resources include historic properties listed in, or eligible for listing in the National Register of Historic Places (36 C.F.R. §60[a-d]). The National Historic Preservation Act of 1966 (NHPA), as amended and recodified (54 U.S.C. § 300308), defines an historic property as "any prehistoric or historic district, site, building, structure, or object included on, or eligible for inclusion on the National Register [of Historic Places]." Under the statute and implementing regulations, historic properties include significant traditional religious and cultural properties important to Indian tribes. Historic properties include built resources (bridges, buildings, piers, etc.), archaeological sites, and Traditional Cultural Properties, which are significant for their association with practices or beliefs of a living community that are both fundamental to that community's history and a piece of the community's cultural identity. Although often associated with Native American traditions, such properties also may be important for their significance to ethnic groups or communities. Historic properties also include submerged resources.

Many aboriginal earth and shell middens are located in the vicinity of Grand Bay NWR. The majority are multi-component earth and shell accumulations, products of hundreds of years of use as seasonal encampments and food processing sites. They are found principally along the remnant river levees of the historical Escatawpa River channel, now known as the Bayou Cumbest, Crooked Bayou, and Heron Bayou systems (USFWS 2008; MDMR 1998b).

By the late 1990s, at least six archaeological or cultural resource surveys had been conducted in the Grand Bay area, though most of these surveys did not contribute new knowledge about the region's past (MDMR 1998b). To date, the refuge has not been systematically surveyed for cultural and archaeological resources, but the presence of additional prehistoric and/or historic resources would be expected.

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred)

The National Historic Preservation Act of 1966 (NHPA) charges the federal government with protecting the cultural heritage and resources of the nation. The selected alternative would be implemented in accordance with all applicable laws and regulations concerning the protection of cultural and historic resources. Cultural and historic resources would be considered when preparing site-specific restoration measures and management actions. Where disturbance of cultural resources is likely, resource managers would conduct reviews and/or surveys to inform the methods and location of restoration and management actions. For site-specific restoration measures and management actions, environmental compliance would be conducted by evaluating each restoration activity and management measure proposed for the parcel(s) against the environmental threshold criteria evaluated under this programmatic analysis. Restoration activities/management measures would be designed to avoid cultural resources to the extent practicable. Resource managers would work with the Mississippi State Historic Preservation Office and the DOI to determine compliance measures if resources are likely in the area or encountered during implementation.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. Even if development were to occur, cultural resources would still be protected under the No Action alternative. Still, development of the area could result in the adverse impacts to cultural resources. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.4.4 Land and Marine Management

Affected Environment

The USFWS manages the Grand Bay NWR while the MDMR manages the Grand Bay NERR and Grand Bay Savanna CP. Management plans are summarized and incorporate by reference here.

Grand Bay National Estuarine Research Reserve Final Environmental Impact Statement/Reserve Management Plan: This EIS was finalized in 1998 by the Mississippi Department of Marine Resources. The purpose of this plan was to designate the area as part of the National Estuarine Research Reserve. For designation, a reserve management plan was produced and in 2013 was updated. The Grand Bay NERR Management Plan 2013-2018 frames out stewardship, resource protection, public use/access, research and monitoring, education and coastal training plans.

<u>Grand Bay National Wildlife Refuge Comprehensive Conservation Plan:</u> This plan was finalized in 2008 by USFWS. The purpose of the plan was to guide management actions and direction over a period of 15 years. Specifically, the CCP was written to:

- Provide a clear statement of the refuge's management direction;
- Provide refuge neighbors, visitors, and government officials with an understanding of the
- USFWS's management actions on and around the refuge;
- Ensure that the USFWS's management actions, including land protection and recreation/education programs, are consistent with the mandates of the National Wildlife Refuge System; and
- Provide a basis for development of the refuge's budget requests for operations, maintenance, and capital improvement needs.

<u>Land Protection Plan and Final Environmental Assessment for the Expansion of Grand Bay National Wildlife Refuge</u>: This plan was finalized in 2012 by USFWS. This plan identified the proposed acquisition boundary for the proposed expansion of NWR. It delineated approximately 8,428 acres from four areas adjacent to the refuge for restoration, enhancement, and management. The purpose of the proposed refuge expansion was to conserve valuable riverine habitat, to protect threatened and endangered species, to restore and protect key habitats (i.e. coastal savanna and longleaf pine), and to manage populations of migratory birds and other interjurisdictional trust species.

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred) The acquisition of up to 8,000 acres (Alternative B), management of up to 17,500 acres (Alternative C) or the both (Alternative D-preferred), is consistent with the current plans for the NWR, Grand Bay NERR and the Grand Bay Savanna CP. Alternative B would provide a long term-benefit to land and marine management by expanding the current public ownership in the area by as much as 8,000 acres. Alternative C would provide a long-term benefit to land and marine management by provide habitat restoration benefits to up to 17,500 acres of currently owned or newly acquired lands within the complex. Alternative D-preferred would provide a long-term benefit to land and marine management by acquisition and management of up to 8,000 acres and/or habitat management on up to 17,500 acres. The planning processes have been included public involvement. There would be a long-term benefit to land and marine management as a result of implementing Alternative B, C or D-preferred.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. However, under alternatives B and D, if development were to occur, there would likely be no effect on land and marine management because existing developments would be completed and would be consistent with existing land use plans. The No Action alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.1.4.5 Public Health and Safety

Affected Environment

Public roads in the proposed project alternative area are subject to flooding on the Grand Bay NERR. A large portion of the area is mapped as Zone VE. Zone VE is defined as Coastal flood zone with velocity hazard. This includes beach areas, open water and most estuarine marsh. Some estuarine marsh, streams, and riparian areas are mapped as Zone AE. Zone AE is defined as "Base Flood"

Elevations Determined". Upland areas are mostly Zone X. Zone X are defined as "Areas of 0.2% annual change flood; areas of 1% annual chance flood with average depths of less than 1 foot or with drainage areas less than 1 square mile; and areas protected by levees from 1% annual chance flood".

Environmental Consequences for WCNH/Birds Proposed Alternatives B, C and D-(Preferred)

There would be short-term, minor, adverse impacts to public health and safety. Exposure to smoke during prescribed burns would adversely impact public health, but these impacts are expected to be minor since prescribed burns are typical in this region and short term. Burn plans that include public notification of burns and controlled access into the site during burns would be developed to minimize the risk and potential exposure of the public to smoke. Fire breaks would restrict fire to designated areas and crews will be on site to ensure that fire does not jump the fire breaks. Safety plans would be part of the controlled burn plans.

Chemical treatment would require use of herbicide that could be hazardous if spilled or handled improperly. Personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities. Most of the applications would be in remote areas where there is limited public access.

The proposed alternative area is designated as floodplain. Preventing development in the floodplain/the transition of native habitats to new impervious surface provides a flood risk/public safety benefit. The proposed alternative would have a beneficial effect to the surrounding communities. It would promote healthy lifestyles by allowing recreational use on previously private parcels of land.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.2). Natural recovery would take much longer compared to a scenario in which restoration actions were undertaken. Land use trends in the vicinity of the project area do not indicate that the parcels are at high risk of development in the foreseeable future. However, under alternatives B and D, if development were to occur, there would likely be no effect to public health and safety because local building codes and ordinances would be followed. to The No Action Alternative would not have short-term, adverse impacts, to public safety from temporary exposure from prescribed fire associated with the implementation of proposed Alternative A. The No Action Alternative does not meet the MS TIG's goals and objectives and does not provide the restoration benefit to WCNH/Birds that would occur through the proposed alternatives.

3.4.2. Site-specific NEPA Review for WCNH/Birds Proposed Alternatives B, C & D-(Preferred)

Section 3.4.1 is a discussion of environmental consequences analysis of proposed Alternatives B, C, and D for WCNH/Birds Restoration Type at a programmatic level. The exact parcels and associated restoration measures and management activities on those parcels are not known at this time. The environmental consequences are based on the range of restoration measures and management activities contemplated on parcels for proposed alternative project areas. The programmatic analysis provides maximum impacts to each of the resource categories based on the MS TIG's knowledge of the proposed alternative project area and the anticipated impacts associated with the planned

restoration measures and management activities. The MS TIG is proposing the selection of Alternative D (Preferred). Section 3.1.2 also presents a process that the MS TIG would follow to complete the requirements of NEPA and other environmental statutes as site-specific restoration measures and management activities are planned for Alternative D, if selected.

3.5 Cumulative Impacts for WCNH/Birds Alternatives

Section 6.6 and Appendix 6B of the PDARP/PEIS are incorporated by reference into the following cumulative impacts analysis including the methodologies for assessing cumulative impacts, identification of affected resources and the cumulative impacts scenario. A development of the analysis in the context of the affected environment of the proposed WCNH/Bird alternatives (X), when added to the impacts from applicable past, present and reasonably foreseeable future actions (Y), to understand the potential cumulative impacts to an affected resource (Z), or where the effects may interact and/or be additive, that is X + Y = Z. This analysis includes the alternatives proposed for the WCNH/Birds Restoration Type in this Draft RP/EA, which include:

- Alternative A (Preferred): Graveline Bay Land Acquisition and Management
- Alternative B: Grand Bay Land Acquisition (up to 8,000 acres)
- Alternative C: Grand Bay Habitat Management (up to 17,500 acres)
- Alternative D (Preferred): Grand Bay Land Acquisition (up to 8,000 acres) and Habitat Management (up to 17,500 acres); Alternatives B and C combined

3.5.1 Identification of Resources Affected

Sections 3.2 and 3.3 provide an environmental consequences analysis for the following resources that would have minor to negligible effects, and based on their magnitude, with respect to context and intensity, would not contribute to cumulative impacts. These resources are excluded from this cumulative impacts analysis:

- Noise
- Marine and Estuarine Fauna
- Infrastructure
- Fisheries and Aquaculture
- Marine Transportation
- Aesthetics and Visual Resources

The following resources were analyzed in detail in Sections 3.2 and 3.3 for environmental consequences that could result from implementation of the proposed WCNH/Birds alternatives.

- Geology and Substrates
- Hydrology and Water Quality
- Air Quality and Greenhouse Gas Emissions
- Habitats
- Wildlife Species (including Birds)
- Protected Species

- Socioeconomics and Environmental Justice
- Cultural Resources
- Tourism and Recreational Use
- Public Health and Safety

Of the resources listed above, most were determined to have impacts that would not contribute to cumulative impacts, based on their magnitude with respect to context and intensity, and are therefore excluded from this cumulative impacts analysis. Only Air quality and greenhouse gas emissions and socioeconomics and environmental justice were carried forward for cumulative impacts analysis.

3.5.1.1 Cumulative Action Scenario

In order to effectively consider the potential cumulative impacts, the MS TIG identified local and site-specific past, current and reasonably foreseeable future actions which are considered relevant to identifying any cumulative impacts the alternatives may have on a local scale.

These actions fall within the established spatial and temporal boundaries. The cumulative impacts analysis depends on the availability of information and data about past, present and reasonably foreseeable future actions. For this Draft RP/EA, the MS TIG identified present and potentially significant future actions through outreach to local, state and/or federal experts familiar with major environmental and development initiatives that have a potential to contribute significantly to cumulative impacts. Publicly available databases⁵³ and projects considered in previous restoration plans (Phase III FERP/PEIS, Phase IV ERP/EAs, and the PDAR/PEIS) were also reviewed to develop this list of actions. The MS TIG also relied on expert judgments, primarily qualitative, about the potential for impacts, using publicly available information about the likely design and location of these actions. Table 3.5-1 provides a listing of actions that the MS TIG considered during this cumulative impacts analysis.

Table 3.5-1: Description of past, present, and reasonably foreseeable future actions considered in the cumulative impact analysis

	ve impuet unuly sis	
		Key Resource Areas with
		Potential for Cumulative
Category/Projects	Project Description	Impacts*
Restoration Related to	the DWH Spill (Early Restoration Phases I, II & III, IV, Restore Ac	ct, Gulf Environmental Benefit
Fund, North American	Wetlands Conservation Fund, National Academy of Sciences)	
NFWF GEBF Invasive	The purpose of the Invasive Species Management on Coastal State	Short-term, adverse impacts to:
Species Management	Lands project is to remove and manage invasive species on state	 air quality and greenhouse
on Coastal State Land	lands in coastal Mississippi in order to enhance natural ecosystem	gases
	functioning of these systems and ensure a sustainable coastal	
	environment. Work will include prescribed burning, mechanical	
	and chemical control of invasive vegetation, and feral hog control.	
	Assessment work is underway. The Mississippi Department of	
	Marine Resources has procured a contractor to begin writing both	
	an invasive species management plan and a prescribed fire	

http://blog.gulflive.com/mississippi-press-news/2014/06/mississippi_coastal_improvemen.html http://www.nfwf.org/whoweare/mediacenter/pr/Pages/gulf-main-pr-14-1117.aspx http://eli-ocean.org/gulf/restoration-projects-database/ https://restoration.atlas.noaa.gov/src/html/index.html http://ms.restore

Category/Projects	Project Description	Key Resource Areas with Potential for Cumulative Impacts*
	management plan. Writing of the plans will take place throughout the fall of 2016.	•
RESTORE Strategic Land Protection, Conservation, and Enhancement of Priority Gulf Coast Landscapes – Bucket 2	This project will protect lands through acquisition and conservation easement programs in areas across the Mississippi Gulf Coast. Priority areas include the Grand Bay National Wildlife Refuge and others.	Long-term adverse impacts to:
NFWF GEBF Acquisition of Priority Tracts for Coastal Habitat Connectivity	This project seeks to enhance coastal habitat connectivity and increase core conservation areas within the Mississippi CP system, the Gulf Islands National Seashore, and the Grand Bay National Wildlife Refuge. The conservation of coastal habitats is one of the fundamental steps in building and maintaining a sustainable, resilient coastal environment. This project will address this conservation need by acquiring key land parcels that provide multiple long-term benefits for the Mississippi Gulf Coast ecosystem.	Long-term, adverse impacts to:
NFWF GEBF Habitat Restoration: Federal Lands Program – Phase I	This project will enhance and restore habitat on federal lands in coastal Mississippi. Anticipated outcomes for key focal habitats include restoration of over 30,000 acres through invasive species removal, forest thinning and prescribed burning on lands contained within Grand Bay National Wildlife Refuge and other locations.	Short-term, adverse impacts to: • air quality and greenhouse gases
Other relevant environ	mental stewardship and restoration activities	
MsCIP Project: Bayou Cumbest Ecosystem Restoration and Hurricane Storm Damage Reduction	The project, which is adjacent to Grand Bay CP, was funded in 2014. This project includes the acquisition of about 61 tracts, removal of 19 structures, excavation and removal of fill material from former home sites and adjacent lands, filling drainage ditches, control of non-native species and planting native emergent wetland species. After acquisition, 148 acres would be restored; 110 to emergent wetlands and 38 to coastal scrub shrub habitat.	Long-term adverse impacts to:
MsCIP Franklin Creek Ecosystem Restoration	This project was funded in 2014 and is located within the alternative project area. It would use ditch and roadbed removal, culvert installation under U.S. 90, non-native species control mechanisms and controlled burning to restore 149 acres north and south of the highway with critical wet pine savannah habitat. The work would also remove about 30 residential structures from the floodplain. The project is planned but not currently funded.	Short-term, adverse impacts to: • air quality and greenhouse gases

The following section describes the cumulative impacts of the alternatives being considered when combined with other past, present and reasonably foreseeable future actions which were identified above. In many situations, implementation of the alternatives would likely help reduce overall long-term adverse impacts by providing a certain level of offsetting benefits, especially when considered in concert with the numerous other present and reasonably foreseeable future actions in the area.

3.5.1.2 Cumulative impact Analysis

Air quality and greenhouse gas emissions

Implementation of the proposed WCNH/Birds alternatives (A, C and D) would have short-term, minor to moderate adverse impacts on air equality and greenhouse gas emissions due to smoke generated during prescribed fire that is anticipated for habitat management. As defined in the PDARP/PEIS, the impacts on air quality could be measurable and would be limited to local and adjacent areas. Emissions of criteria pollutants could be at EPA's de minimis criteria levels for

general conformity determination under the Clean Air Act (40 CFR § 93.153). Prescribed fire activities would occur periodically according to site-specific management plans and burn plans, typically occurring every other year during the growing season. Limiting factors include wind, humidity, available personnel and other factors. Prescribed fire frequency will be intermittent and coordinated by resource managers so as not to occur simultaneously. The alternatives would not have cumulative long-term impacts on air quality or to emissions of greenhouse gases. Jackson County, Mississippi (as well as all other counties in Mississippi) is classified as in attainment, meaning criteria air pollutants do not exceed National Ambient Air Quality Standards (NAAQS). MDEQ monitors air quality at a station in Jackson County. Baldwin County, Alabama, is classified as unclassifiable/attainment⁵⁴.

Periodic prescribed fire practices would not cause an adverse cumulative impact, because it is not anticipated that the levels of particulates and emissions created by prescribed fire would be sufficient for the project area to exceed attainment criteria established by the EPA.

Long term beneficial impacts to air and greenhouse gas emissions would also be anticipated due to re-vegetation and carbon sequestration that would occur during habitat management (Alternatives A, C, and D) and as a result of acquisition (Alternatives A, B, and D) that would prevent development and provide for preservation in perpetuity.

Four projects are identified as potential contributors to cumulative impacts (four adverse and one beneficial) on air quality and greenhouse gas emissions when their impacts are combined with those of the alternatives: NFWF GEBF Invasive Species Management on Coastal State Land; NFWF GEBF Habitat Restoration: Federal Lands Program – Phase I, and MsCIP Franklin Creek Ecosystem Restoration, and industrial operations in the project area. Smoke from prescribed fire associated with these projects would result in short-term minor to moderate air quality impacts. However, three other projects considered in the cumulative impacts analysis (RESTORE Strategic Land Protection, Conservation, and Enhancement of Priority Gulf Coast Landscapes – Bucket 2, NFWF GEBF Acquisition of Priority Tracts for Coastal Habitat Connectivity, and MsCIP Project: Bayou Cumbest Ecosystem Restoration and Hurricane Storm Damage Reduction) would provide a long-term beneficial impact to air quality and greenhouse gas emissions by carbon sequestration preservation as a result of land acquisition, which would prevent development in perpetuity and prevent development d

When the proposed WCHN/Birds alternatives A-D are analyzed in combination with these past present, and reasonably foreseeable future actions, short-term cumulative adverse impacts to air quality and greenhouse gas emissions would likely occur. The alternatives would not contribute substantially to cumulative adverse impacts to air quality and greenhouse gas emissions. The alternatives, carried out in conjunction with other restoration efforts, would also have the potential to result in some long-term beneficial cumulative impacts to air quality.

Socioeconomics and Environmental Justice:

Land acquisition anticipated for Alternatives A, B and D could have a minor to moderate long-term impact on socioeconomic resources due properties being removed from the local tax base

⁵⁴ https://www.regulations.gov/document?D=EPA-HQ-OAR-2012-0918-0426

permanently. Individuals would not be adversely affected because any property transfers would be on a appraised value basis between willing parties. There could be long-term beneficial impacts due to increased visitor spending in the area as a result of increased recreational access to the project areas.

Three projects are identified as potential contributors to cumulative impacts (adverse and beneficial) on socioeconomics when their impacts are combined with those of the alternatives: RESTORE Strategic Land Protection, Conservation, and Enhancement of Priority Gulf Coast Landscapes – Bucket 2, NFWF GEBF Acquisition of Priority Tracts for Coastal Habitat Connectivity, and MsCIP Project: Bayou Cumbest Ecosystem Restoration and Hurricane Storm Damage Reduction. All of these projects involve voluntary land acquisition, which could permanently affect the local tax base, but could also provide a long-term beneficial impact by increasing visitor spending.

When the proposed WCHN/Birds alternatives (A, B and D) are analyzed in combination with these past, present, and reasonably foreseeable future actions, long-term cumulative adverse impacts to socioeconomics would likely occur. The alternatives would not contribute substantially to cumulative adverse impacts. The alternatives, carried out in conjunction with other restoration efforts, would also have the potential to result in some long-term beneficial cumulative impacts to socioeconomics.

3.6 Comparison of the Alternatives-WCNH/Birds Restoration Type

This section provides a comparison of the NEPA environmental consequences for the reasonable range of alternatives for the WCNH/Birds Restoration Type (Table 3.6-1). The proposed alternatives include four action alternatives as well as a Natural Recovery/No Action and are described in Table 3.6-1.

Table 3.6-1: Summary of the Comparison of the WCNH/Birds Restoration Type Alternatives

Comparison of WCNH/Bird Restoration Type Alternatives		
Alternative		
Alternative A (Preferred):	Graveline Bay Land Acquisition and Management Alternative A would provide the opportunity to implement WCNR/Bird conservation practices as well contribute to the habitat connectivity of the area, and preclude development on 1,410 acres in Graveline Bay.	
Alternative B:	Grand Bay Land Acquisition (up to 8,000 acres) Alternative B would include acquisition to reduce the threat of further development, decreased habitat fragmentation, and increased habitat connectivity to other large conservation parcels in Grand Bay NWR, NERR, and CP area.	
Alternative C:	Grand Bay Habitat Management (up to 17,500 acres) Alternative C would include habitat management on current public lands within the NWR, NERR and CP boundaries. Restoration measures and benefits provide for more effective large-scale management efforts and habitat enhancement.	
Alternative D:	Grand Bay Land Acquisition (up to 8,000 acres) and Habitat Management (up to 17,500 acres) Alternative D would combine the benefits from Alternative B and C.	

Comparison of WCNH/Bird Restoration Type Alternatives			
No Action Alternative:	Under the No Action Alternative, the MS TIG would not implement any projects for the WCNH/Birds Restoration Type at this time, and would instead allow natural recovery processes to occur, which could result in one of four outcomes for injured resources: 1) gradual recovery, 2) partial recovery, 3) no recovery, or 4) further deterioration. Although injured resources could presumably recover to or near baseline conditions under this scenario, recovery would take much longer compared to a scenario in which restoration actions were undertaken.		
Physical Env	rironment		
Alternative A (Preferred):	Based on the analysis impacts there would be short-term to long-term, minor to moderate and adverse impacts to soils and hydrology. There would be short-term, minor to moderate, adverse impacts to water quality, wetlands, and air quality and green house gases. There would be long-term benefits to soil, hydrology, floodplains, and wetlands. There would be short-term and long-term benefits to water quality.		
Alternative B:	There would be long-term, minor, adverse impacts to geology and substrates due to increased public use. There would be long-term benefits to hydrology, water quality, and wetlands by preventing development.		
Alternative C:	There would be short-term, minor to moderate, adverse impacts to geology and substrates due to soil disturbance during habitat management-mechanical treatment, chemical treatment, prescribed fire. There would be short-term, minor to moderate, adverse impacts to hydrology, water quality, and wetlands due to temporary changes to stormwater flows and runoff retention patterns due to rutting by equipment and vegetation removal during habitat management activities. There would be short-term moderate adverse impacts to air quality and greenhouse gases during the prescribed fire events.		
Alternative D:	Alternative D would combine the adverse and beneficial impacts of Alternative B and C.		
No Action Alternative:	This alternative is not expected to contribute to short-term or long term, or cumulative adverse impacts to physical resources. The No Action Alternative does not provide the restoration benefits to WCNH/Birds that would occur through the proposed alternatives.		
Biological Er	nvironment		
Alternative A (Preferred):	There would be short-term, minor to moderate, adverse impacts to habitat and wildlife. There would be short-term and long-term benefits to habitat and wildlife. The following federally protected species could be present within the proposed alternative project area: Alabama Red-Belly Turtle, Piping plover, <i>Black pine snake</i> ; gopher tortoise, <i>Louisiana quillwort</i> , Mississippi Sandhill Crane, and Mississippi diamondback terrapin. Coordination with the USFWS Ecological Services Field Office in Jackson, Mississippi would be completed to identify whether protected species or their habitats could be affected by site-specific restoration activities and management measures. Migratory bird species groups that could occur in the proposed alternative project area include wading birds, shorebirds, seabirds, raptors, goatsuckers, waterfowl, doves and pigeons, and rails and coots. For all planned restoration activities, pre-commencement nesting surveys for migratory birds and raptors within the site-specific project area would be conducted and if evidence of nesting is found, coordination with the USFWS would be completed to develop and implement appropriate measures so that no "take" of nesting birds is anticipated.		
Alternative B:	There would be long-term benefits to habitats and wildlife by preventing development.		

Comparison of WCNH/Bird Restoration Type Alternatives			
Alternative C:	There would be short-term, minor to moderate, adverse impacts to habitat and wildlife due to site disturbance during restoration activities.		
	The following federally protected species could be present within the proposed alternative project area: Alabama Red-Belly Turtle, Piping plover, <i>Black pine snake</i> ; gopher tortoise, <i>Louisiana quillwort</i> , Mississippi Sandhill Crane, and Mississippi diamondback terrapin. Coordination with the USFWS Ecological Services Field Office in Jackson, Mississippi would be completed to identify whether protected species or their habitats could be affected by site-specific restoration activities and management measures.		
	Migratory bird species groups that could occur in the proposed alternative project area include wading birds, shorebirds, seabirds, raptors, goatsuckers, waterfowl, doves and pigeons, and rails and coots. For all planned restoration activities, pre-commencement nesting surveys for migratory birds and raptors within the site-specific project area would be conducted and if evidence of nesting is found, Coordination with the USFWS would be completed to develop and implement appropriate measures so that no "take" of nesting birds is anticipated.		
	There would be long-term benefits to habitats by implementing activities designed to enhance habitat.		
Alternative D:	Alternative D would combine the adverse and beneficial impacts of Alternative B and C.		
No Action Alternative:	This alternative is not expected to contribute to short-term or long term, cumulative adverse impacts to biological resources. The No Action Alternative does not provide the restoration benefits to WCNH/Birds that would occur through the proposed alternatives.		
Socioeconom	ic Environment		
Alternative A (Preferred):	Land acquisition could have a short-term, minor to moderate impact on socioeconomic resources due to changes in visitor spending and tax impacts. There would be short-term, minor adverse impacts to tourism and recreation during prescribed burns. There would be a long-term, minor to moderate, adverse effect to land and marine management as acquired properties would not be available for development.		
	For site-specific restoration activities and management measures, environmental reviews and surveys would be conducted if cultural resources are suspected in the area. Resources that are eligible for the National Register of Historic Places would be avoided in the design of the restoration activity and management measure. There would be no adverse impact to cultural resources.		
	There would be a minor short-term, adverse impacts to public health and safety. Exposure to smoke during prescribed burns would adversely impact public health, but these impacts are expected to be minor since prescribed burns are typical in this region and short term. The proposed alternative would have a beneficial effect to the surrounding communities. It would promote healthy lifestyles by allowing recreational use on previously private parcels of land.		
Alternative B:	Land acquisition could have a minor to moderate impact on socioeconomic resources due to changes in visitor spending and tax impacts.		
	There would be long-term benefit to tourism and recreation opportunities by expanding the area recreational activities including wildlife observation, hunting, boating, and hiking.		
	There would be long term-benefits to land and marine management by expanding the current public ownership.		
	There would be a beneficial effect to the surrounding communities by promoting healthy lifestyles by allowing recreational use on previously private parcels of land and by preventing development in the floodplain, thereby reducing flood risk.		
Alternative C:	Management activities could have short-term, minor impact to tourism and recreation.		
	For site-specific restoration activities, environmental reviews and surveys would be conducted if cultural resources are suspected in the area. Resources that are eligible for the National Register of Historic Places would be avoided in the design of the restoration activities and management measures. There would be no adverse impact to cultural resources.		
	There would be long-term benefit to land and marine management by habitat restoration benefits to up to 17,500 acres of currently owned lands.		
	There would be minor, short-term adverse impacts to public health and safety. Exposure to smoke during prescribed burns would adversely impact public health. There would be a beneficial effect to the surrounding communities by promoting healthy lifestyles by allowing recreational use on previously private parcels of land and by preventing development in the floodplain, thereby reducing flood risk.		

	Comparison of WCNH/Bird Restoration Type Alternatives
Alternative D:	Alternative D would combine the adverse and beneficial impacts of Alternative B and C.
No Action	This alternative is not expected to contribute to short-term or long term, cumulative adverse impacts to socioeconomics. The The No Action Alternative does not provide the restoration benefits to WCNH/Birds that would occur through the proposed alternatives.
Cumulative I	Effects
Alternatives A (Preferred), B, C and D (Preferred)	There could be minor to moderate, long-term adverse impact to socioeconomic resources (A, B and D due to acquired properties being removed from local tax base and from development. There could be long-term beneficial impacts from increased visitor spending resulting from added recreational access (A-D). Carried out with other past, present, and reasonably foreseeable future actions, long-term cumulative adverse impacts to socioeconomics would not contribute substantially to cumulative adverse impacts to socioeconomic resources. There could be increased visitor use as a result of the acquisition and preservation of lands in perpetuity. Implementation of the proposed WCNH/Bird alternatives (A, C and D) would have short-term, minor to moderate adverse impacts on air equality and greenhouse gas emissions due to smoke generated during prescribed fire that is anticipated for habitat management. Long term beneficial impacts to air and greenhouse gas emissions would also be anticipated due to re-vegetation and carbon sequestration that would occur during habitat management (Alternatives A, C, and D) and as a result of acquisition (Alternatives A and D) that would prevent development and provide for preservation in perpetuity. When the proposed WCHN/Birds alternatives A-D are analyzed in combination with these past present, and reasonably foreseeable future actions, implementation of the alternatives would not contribute substantially to cumulative adverse impacts to air quality and greenhouse gas emissions. The alternatives, carried out in conjunction with other restoration efforts, would also have the potential to result in some long-term beneficial cumulative impacts to air quality.
No Action	There would be no beneficial impacts or short or long-term cumulative adverse impacts to resources.

The MS TIG is proposing to select Alternative A (Preferred): Graveline Bay Land Acquisition and Management and Alternative D (Preferred): Grand Bay Land Acquisition and Habitat Management. Table 3.6.27 above summarizes the environmental consequences for the proposed alternatives in the Draft RP/EA. Subsequent environmental review will occur in addition to this programmatic review to determine whether planned site-specific restoration activities and management measures are within the maximum expected impacts described in this Draft RP/EA. As described in section 3.1.2, an Environmental Evaluation Worksheet would be used to document the results of the environmental evaluation is in Appendix A. If the planned site-specific restoration activities and management measures are likely to exceed the maximum expected impacts described in this Draft RP/EA, the MS TIG will undertake additional environmental review consistent with NEPA requirements and other requirements for protection of the environment or will abandon the planned project. The MS TIG does not propose to take actions that would result in any significant adverse impacts on the environment.

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific restoration activities and management measures in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to resources:

Geology and Substrates

• Allow revegetation of fire breaks or actively revegetation with native species or annual grasses, if prolonged period of greening up is anticipated.

- Develop and implement an oil spill prevention and response plan, including conducting daily inspections during chemical treatment, mechanical treatment, and prescribed fire operations to ensure there are no leaks of antifreeze, hydraulic fluid, pesticide, or other substances.
- To the extent practicable, for equipment use in wet areas, soft track or wide track equipment should be used to distribute the equipment weight and minimize impacts to soils. Alternatively, crews may remove vegetative materials with chainsaws.

Hydrology and Water Quality

- In the execution of land acquisition and the design of habitat management measures the MS TIG would consider resiliency measures to facilitate habitat migration due to sea level rise (CEQ, 2016).
- Develop and implement an erosion control plan to minimize erosion during and after construction and where possible use vegetative buffers (100 feet or greater), revegetate with native species or annual grasses, and conduct work during dry seasons.
- For chemical treatment, personnel applying chemicals would follow all warning labels on chemical containers and proper permits would be secured prior to treatment activities.
 Personnel will apply herbicide in accordance with the direction and guidance provided on the appropriate U.S. Environmental Protection Agency (EPA) labels and state statutes during land-based activities.
- Soft track or wide track equipment would be used in wet areas to the extent practicable. Alternatively, crews may remove vegetative material with chainsaws.
- Avoid and minimize, to the maximum extent practicable, placement of dredged or dill
 material in wetlands and other aquatic resources. Design construction equipment corridors to
 avoid and minimize impacts to wetlands and other aquatic resources to the maximum extent
 practicable. If required, a USACE permit would be obtained; likely a Nationwide 27
 (Aquatic Habitat Restoration, Establishment, and Enhancement Activities) as well as MDMR
 Coastal Wetlands Permit (if required). USACE permit and/or MDME Coastal Wetlands
 permit conditions (if required) would be adhered to in all operations.
- Designate a vehicle staging area removed from any natural surface water resource or wetland
 to perform fueling, maintenance, and storage of construction vehicles and equipment. Inspect
 vehicles and equipment daily prior to leaving the storage area to ensure that no petroleum or
 oil products are leaking.
- Develop and implement a spill prevention and response plan, including conducting daily inspections of all construction and related equipment to ensure there are no leaks of antifreeze, hydraulic fluid, or either substances and cleaning and sealing all equipment that would be used in the water to rid it of chemical residue.
- Controlling dust related to construction site activities through a Soil Erosion Sediment Control Plan that includes spraying of a suppressing agent on dust piles (non-hazardous, biodegradable).
- Covering trucks hauling loose materials.

Air Quality and Greenhouse Gas

- Shut down idling construction equipment, if feasible.
- Encourage the use of the proper size of equipment for the job to maximize energy efficiency.
- Use of ultra-low sulfur diesel fuel in off-road construction equipment with engine horsepower (HP) rating of 60 HP and above.

- Controlling dust related to construction site activities through a soil erosion sediment control plan that includes spraying of a suppressing agent on dust piles (non-hazardous, biodegradable).
- Covering trucks hauling loose materials.

Habitat and Wildlife

Prior to bringing any equipment (including personal gear, machinery, vehicles, or vessels) to
the work site, inspect each item for mud or soil, seeds, and vegetation. If present, clean the
equipment, vehicles, or personal gear until they are free from mud, soil, seeds, and
vegetation. Inspect the equipment, vehicles, and personal gear each time they are being
prepared to go to a site or prior to transferring between sites to avoid spreading exotic,
nuisance species.

Protected Species

• Provide all individuals working on a project with information in support of general awareness of and means to avoid impacts to protected species and their habitats present at the specific project site.

Piping Plover and Red Knot

- Provide all individuals working on a restoration activities associated with the proposed alternative with information in support of general awareness of piping plover or red knot presence and means to avoid birds and their Critical or otherwise important habitats.
- Minimize vegetation planting in preferred habitats and avoid removal of wrack year-around along the shoreline.
- During recreational use, enforce leash or "no pet" policies in Critical or important habitats.

Gopher Tortoise

• If suitable habitat is present, coordinate with the local USFWS Ecological Services Field Office to discuss the need for surveys to identify any gopher tortoise burrows and to develop conservation measures to avoid or minimize impacts. Measures could include establishing a protective buffer (size determined by USFWS and the state trust resource agency) if burrows are within the project area and cannot be avoided, implementing standard procedures to relocate the tortoise within the project site but away from the areas of construction or restoration or considering conservation banks. A Candidate Conservation Agreement with Assurances may be appropriate for project sites within the non-listed range of the species.

Protected Plants

- If suitable habitat is present, coordinate with the local USFWS Ecological Services Field Office to discuss the need for surveys to identify protected plants and to develop conservation measures to avoid or minimize impacts.
- Enhance and protect plants on site and in adjacent habitats to the maximum extent possible.

Migratory Birds

• Use care to avoid birds when operating machinery or vehicles near birds.

153

- Avoid working in migratory bird nesting habitats during breeding, nesting, and fledging
 (approximately mid-February through late August). If project activities must occur during
 this timeframe and breeding, nesting, or fledging birds are present, contact the state trust
 resource agency to obtain the most recent guidance to protect nesting birds or rookeries, and
 their recommendations will be implemented.
- Conservation areas may already be marked to protect bird nesting areas. Stay out of existing marked areas.
- If vegetation clearing is necessary, clear vegetation outside the migratory bird nesting season (approximately mid-February through late August) or have a qualified biologist inspect for active nests. If no active nests are found, vegetation may be removed. If active nests are found, vegetation may be removed after the nest successfully fledges.

Bald Eagles

- If bald eagle breeding or nesting behaviors are observed or a nest is discovered or known, have all activities avoid the nest by a minimum of 660 feet. If the nest is protected by a vegetated buffer where there is no line of sight to the nest, then the minimum avoidance distance is 330 feet. Maintain this avoidance distance from the onset of breeding/courtship behaviors until any eggs have hatched and eaglets have fledged (approximately 6 months).
- If a similar activity (such as driving on a roadway) is closer than 660 feet to a nest, maintain a distance buffer as close to the nest as the existing tolerated activity. If a vegetated buffer is present and there is no line of sight to the nest and a similar activity is closer than 330 feet to a nest, then maintain a distance buffer as close to the nest as the existing tolerated activity.
- In some instances, activities conducted within 660 feet of a nest may result in disturbance, particularly for the eagles occupying the Mississippi barrier islands. If an activity appears to cause initial disturbance, stop the activity and move all individuals and equipment away until the eagles are no longer displaying disturbance behaviors. Contact the USFWS's Migratory Bird Permit Office to determine how to avoid impacts or if a permit may be needed.

3.7 NR (Nonpoint Source) Restoration Type

Section 3.7.1 provides the OPA evaluation for the No Action Alternative and Nutrient Reduction Alternatives A and B. The implementation of conservation practices under these alternatives would be dependent on willing landowners and successful conservation planning to implement those actions. Section 3.7.2 describes the programmatic approach to this NEPA analysis and for NEPA review after site-specific conservation practices have been identified. In addition to incorporating by reference the analysis the USDA-NRCS has conducted on the effects of its conservation practices, the discussion in this Draft RP/EA includes examples of the conservation practices the MS TIG expects would be implemented in the proposed project area and how those practices are expected to impact the environment. Appendix B includes the full list of conservation practices that would be eligible for funding under the alternatives.

3.7.1 OPA Evaluation for NR (Nonpoint Source)

The Nutrient Reduction proposed project alternatives are consistent with the Restore Water Quality Programmatic Goal and the NR Restoration Type in the PDARP/PEIS. Table 3.7-28 provides an OPA

evaluation of each NR alternative and the No Action alternative using the standard OPA evaluation criteria described in OPA implementing regulations at 15 CFR 990.54. These OPA evaluation criteria are listed below:

- The cost to carry out the alternative (The Cost).
- The extent to which each alternative is expected to meet the Trustees' goals and objectives in returning the injured natural resources and services to baseline and/or compensating for interim losses (Restoration Goals and Objectives).
- The likelihood of success of each alternative (Likelihood of Success).
- The extent to which each alternative will prevent future injury as a result of the incident and avoid collateral injury as a result of implementing the alternative (Avoidance of Further Injury/Collateral Injury).
- The extent to which each alternative benefits more than one natural resource and/or service (Multiple Resource Benefits).
- The effect of each alternative on public health and safety (Public Health and Safety).

Table 3.7-1: NR (Nonpoint Source) -OPA Evaluation of Alternatives

Alternative	OPA Evaluation Criteria		
	Cost		
Natural Recovery/No Action Alternative	There is no financial cost associated with the No Action Alternative.		
Alternative A: Upper Pascagoula River Water Quality Enhancement	Alternative A: The cost of \$4 M for development and implementation of conservation plans and practices in the Chunky and Okatibbee watersheds is reasonable for the proposed alternative. USDA-NRCSwould implement this proposed alternative by helping landowners voluntarily implement conservation practices that reduce nutrient and sediment runoff. Through their experience with the Environmental Quality Incentives Program (EQIP), USDA- NRCS is knowledgeable about the cost of successful implementation of the proposed conservation practices. For Alternative A, there would be an opportunity to implement Ecological/NR conservation practices and soil and water conservation practices with willing participants, allowing for a wide array of benefits to cropland, pasture/grassland, associated agriculture lands and riparian areas. This alternative would be more cost-effective as it incorporates soil and water conservation practices on agricultural land, near the source nutrient and sediment runoff. Addressing nutrient and sediment runoff near the source coupled with conservation practices that improve the filtering ability of riparian areas would be more efficient and effective at nutrient reduction than restricting practices only to riparian areas. The MIS TIG anticipates that the proposed alternative would result in improved water quality by reducing nutrient and sediment runoff into coastal waters.		

Alternative	OPA Evaluation Criteria	
Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan	Alternative B: The cost of \$4 M for the enhancement or establishment of riparian buffers in the Chunky and Okatibbee watersheds is reasonable for the proposed project. USDA-NRCS would implement this project by helping landowners voluntarily implement practices that reduce nutrient and sediment runoff by implementing Ecological/NR conservation practices in riparian areas in the proposed project area in the Chunky and Okatibbee watersheds. Through their experience with EQIP, USDA-NRCS is knowledgeable about the cost of successful implementation of the proposed conservations practices. For Alternative B, there would be an opportunity to implement Ecological/NR conservation practices with willing participants, allowing for a wide array of benefits to riparian areas the within farmsteads. The MIS TIG anticipates that the project would result in improved water quality by reducing nutrient and sediment runoff into coastal waters.	
	Restoration Goals and Objectives	
Natural Recovery/No Action Alternative	The No Action Alternative is not consistent with the MS TIG's goal to pursue restoration projects that would provide water quality benefits by NR (nonpoint source).	
Alternative A: Upper Pascagoula River Water Quality Enhancement	Alternative A has a clear nexus to the NR injuries described in the PDARP/PEIS because implementation of conservation practices on privately owned lands would reduce nutrient enrichment and sedimentation and restore water quality in Gulf of Mexico coastal watersheds. The health of the Gulf of Mexico depends upon the health of its estuaries, and the health of those coastal waters is influenced by land use upstream along tributary rivers. The primary goal for this proposed alternative is water quality improvement through the NR restoration type. This watershed-scale proposed alternative restores water quality impacted by the <i>DWH</i> oil spill by reducing the levels of nutrients and sediments entering the Gulf of Mexico. Runoff from cropland, pasture/grassland, and forest contributes nutrients and sediment that adversely impact the health of coastal waters. The proposed conservation practices would reduce nutrient and sediment losses from the landscape, reduce nutrient and sediment loads to streams and downstream receiving waters, and reduce water quality degradation in watersheds that would provide benefits to marine resources and benefits to coastal watersheds.	
	Further, Alternative A is consistent with existing MS TIG goals and objectives that focus on opportunities for leveraged funding, Trustee expertise from state and federal programs and resource management expertise, and projects that are consistent with existing management plans and initiatives. This alternative meets these goals by utilizing Ecological/NR and soil and water conservation/NR practices. It includes an additional \$1 M of leveraged funding from USDA-NRCS for developing conservation plans and implementing conservation practices in the proposed alternative project area. This alternative also utilizes expertise from USDA-NRCS, and is consistent with the PDARP/PEIS and the MGCRP.	
Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan	Alternative B has a clear nexus to the NR injuries described in the PDARP/PEIS because implementation of conservation practices on privately owned lands would reduce nutrient enrichment and sedimentation and restore water quality in Gulf of Mexico coastal watersheds. The health of the Gulf of Mexico depends upon the health of its estuaries, and the health of those coastal waters is influenced by land use upstream along tributaries and rivers. The primary goal for this proposed alternative is water quality improvement through the NR restoration type. by establishing or enhancing riparian buffers within the project area. This watershed-scale proposed alternative restores water quality impacted	

⁵⁵ In general the efficiency of nutrient and sediment removal would depend on the width of riparian buffers,types of plant materials used and storm events.

Alternative	OPA Evaluation Criteria
	by the <i>DWH</i> oil spill by reducing the levels of nutrients and sediments entering the Gulf of Mexico by applying conservation practices in riparian areas. Conservation practices in the riparian area can treat runoff from cropland, pasture/grassland, and forestland that contributes nutrients and sediment that adversely impact the health of coastal waters. The proposed conservation practices would reduce nutrient and sediment losses from the landscape, reduce nutrient and sediment loads to streams and downstream receiving waters, and reduce water quality degradation in watersheds that could provide benefits to marine resources and benefits to coastal watersheds.
	Alternative B would focus on riparian areas within agricultural associated land and forested land in the proposed alternative project area. This alternative would seek to identify opportunities to implement Ecological/NR conservation practices in riparian buffers along the Pascagoula River and its tributaries in the proposed alternative project area. Conservation practice opportunities within one mile of tributaries that already have riparian buffers and areas where new riparian buffers could be successfully established would be a priority. Riparian buffers act to partially protect streams from the impact of adjacent land uses. Buffers would increase water quality in associated streams as sediment is intercepted, serve to provide habitat, and reduce bank erosion by providing bank stabilization. With planning and monitoring, riparian buffers and other related conservation practices would help control channel instability, head-cutting, mass slumping, and wetland degradation. Riparian buffers that exist currently and proper planning of new buffers would help mitigate future water quality degradation. Further, Alternative B is consistent with existing MS TIG goals and objectives that focus on opportunities for leveraged funding, Trustee expertise from state and federal programs and resource management expertise, and projects that arewere consistent with existing management plans and initiatives. This alternative meets these goals by utilizing Ecological/NR conservation practices. It includes an additional \$1 M of leveraged funding from USDA-NRCS for establishing or enhancing riparian areas within the proposed alternative project area. This alternative also utilizes expertise from USDA-NRCS, and is consistent with the PDARP/PEIS and the MGCRP.
	Likelihood of Success
Natural Recovery/No Action Alternative	The No Action Alternative would not contribute to restoring, replacing, or enhancing injured natural resources and would not provide for compensation of interim natural resource losses that occurred as result of the <i>DWH</i> oil spill.
Alternative A: Upper Pascagoula River Water Quality Enhancement; Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan	Alternatives A & B: The MS TIG and its implementing agency Trustee, USDA-NRCS, has demonstrated success in developing and implementing the same types of conservation practices in the proposed alternative project area and other similar watersheds. Given their extensive experience and expertise in conservation practices, the success and legacy of the USDA-NRCS Farm Bill programs, and their established level of trust and cooperation with private landowners, there is a significant opportunity to implement conservation practices on private lands that would reduce the levels of nutrients and sediments entering watersheds that could provide benefits to marine resources and benefits to coastal watersheds.
	Avoidance of Further Injury/Collateral Injury
Natural Recovery/No Action Alternative	The No Action Alternative would not cause further injury, but would also not provide benefit to offset interim losses.
Alternative A: Upper Pascagoula River Water Quality Enhancement; Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan	Alternatives A & B: USDA-NRCS applies conservation practices according to standards that require use of associated and mitigating practices in a "systems approach" to ensure new injuries do not occur and those practice standards would be followed under either Alternative A or B. In addition, the MS TIG would

Alternative	OPA Evaluation Criteria
	ensure compliance with all applicable federal laws, regulations and executive orders prior to implementation of the selected alternative by using a site-specific environmental evaluation process carried out during the conservation planning effort. This process would include conducting any necessary agency consultations and obtaining any required permits. Among other things, the environmental evaluation will identify mitigation measures needed and determine whether there is potential for significant adverse effects to be created. If such potential exists, that particular project will be abandoned or redesigned to minimize the impacts. The proposed alternative would meet all the OPA and NEPA requirements as discussed in Sections 3.0 and 4.0 of this Draft RP/EA. In addition to addressing unique resources site-specifically, the MS TIG is also undertaking programmatic consultations under ESA and other laws for protection of the environment. For example, USDA-NRCS has completed a programmatic ESA consultation with the U.S. Fish and Wildlife Service, confirming USDA-NRCS experience that the conservation practices likely to be implemented under the proposed action alternative may affect but are not likely to adversely affect protected species. The MS TIG has initiated coordination to confirm that the Services agree the NR projects proposed in this RP/EA similarly may affect but would not be likely to adversely affect protected species in the project area. The MS TIG has similar efforts underway to ensure no further injury to other resources, as well.
Natural Recovery/No Action Alterantive	Multiple Resource Benefits The No Action Alternative could provide for multiple resource benefits; however, recovery rates of multiple resources would be less than if the MS TIG pursued active restoration activities included in the Proposed Actions.
Alternative A: Upper Pascagoula River Water Quality Enhancement; Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan	Under both proposed Alternatives A and B, various conservation practices would be conducted on private lands to address nutrient reduction. Through a coordinated and integrated watershed approach to proposed alternative implementation, benefits to multiple resources are anticipated and would include reductions in nutrient and sediment losses from the landscape; reductions in nutrient and sediment loads to streams and downstream receiving waters; that could provide benefits to marine resources and benefits to coastal watersheds. For example, either of these alternatives would reduce nutrient and sediment loading contribution in watersheds in the Upper Pascagoula River basin that contain Gulf sturgeon Critical Habitat. The Gulf sturgeon is anadromous, spending much of its life in marine environments, but spawning occurs in the Upper Pascagoula River and tributaries and in other river systems in the Gulf. Decreasing sediment and other pollutants as proposed under these alternatives may improve Gulf sturgeon spawning success.

Project Alternatives A and B would meet the evaluation criteria established by OPA because:

- Cost estimates are based on comparable projects previously implemented and those costs were considered reasonable;
- The project alternatives have a clear nexus to the NR injuries described in the PDARP/PEIS, and the MS TIG's restoration goals and objectives that would be met include opportunities for leveraged funding, Trustee expertise from state and federal programs and implementing agency resource management expertise, and consistency with existing management plans and initiatives;
- There is a high likelihood of success because these alternatives propose implementing proven conservation practices and tested restoration techniques used by the MS TIG Trustees and project partners on similar types of projects in the region;
- These watershed-scale proposed alternatives improve the quality of coastal waters impacted by the *DWH* oil spill by reducing the runoff of nutrients, and sediment into coastal waters;

- Future and collateral injury would be avoided by employing best practices during project implementation;
- Both alternatives are likely to benefit more than one resource; and
- There would be a long-term benefit to public safety from improved water quality.

Proposed Alternatives A and B are also consistent with the MGCRP and other regional planning initiatives. The nexus between these alternatives and the injury and the programmatic restoration goal is clear because implementation of conservation practices on privately owned lands would reduce nutrient enrichment and sedimentation and restore water quality in Gulf of Mexico coastal watersheds Future conservation planning and implementation of USDA-NRCS conservation practices would not require additional OPA evaluation.

3.7.2 NEPA Analytical Approach for NR (Nonpoint Source) Restoration Type

This section provides the NEPA analytical approach for the NR (Nonpoint Source) Restoration Type in the following order:

- 1. USDA NEPA Analyses for conservation practices incorporated by reference;
- 2. a description of the general NEPA analytical approach for the NR (Nonpoint Source) project alternatives:
- 3. the MS TIG plan for site-specific NEPA review for the selected alternative; and
- 4. the organization of the affected environment and environmental consequences for the proposed alternatives under the NR (Nonpoint Source) Restoration Type.

1) USDA NEPA Analyses for Conservation Practices Incorporated by Reference: The USDA-NRCS has a long-standing structured, interdisciplinary, science-based, and public process for developing conservation practice standards and analyzing the effects of those practices. ⁵⁶ Implementing these conservation practices has been proven to successfully address natural resource concerns related to agricultural and forested lands, and many of these practices can be used to achieve a number of the restoration types identified in the *DWH* PDARP/PEIS. Because of this, both of the proposed action alternatives contemplate using USDA-NRCS conservation practices to achieve certain PDARP restoration goals in Mississippi. This analysis hereby incorporates by reference the standards and specifications for the conservation practices in Appendix B found in the USDA-NRCS National Handbook of Conservation Practices and the analysis of the effects of those practices contained in the USDA-NRCS Conservation Practice Physical Effects (CPPE) matrices, the Network Effects Diagrams, ⁵⁷ and in the USDA-NRCS Conservation Effects Assessment Project reports. ⁵⁸

⁵⁶ See, for example, the Environmental Quality Incentives Program Programmatic EA, March 2016 at https://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/technical/nra/ec/?cid=nrcseprd387616 and research associated with the NRCS Conservation Effects Assessment Project at https://www.nrcs.usda.gov/wps/portal/nrcs/main/national/technical/nra/ceap/. See also the national NRCS conservation practice standards and associated CPPE and Network Effects Diagrams at https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/ ?cid=nrcs143 026849.

⁵⁷ Both the CPPE matrices and network effects diagrams are available from the NRCS National Handbook of Conservation Practices website at https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs143_026849.

⁵⁸ https://www.nrcs.usda.gov/wps/portal/nrcs/main/national/technical/nra/ceap/.

Each of those assessments is based on a review of the best available scientific studies and methodological approaches, as well as professional judgment. ⁵⁹ In addition, this document incorporates by reference the analyses from the USDA-NRCS EQIP Programmatic EA, March 2016, and in particular its discussions of the water quality impacts of NRCS conservation practices.

2) The NEPA Analytical Approach for the Development of NR (Nonpoint Source) Project **Alternatives**: This Draft RP/EA analyzes potential environmental impacts at a broad program scale, identifying the qualitative effects that are a reasonably foreseeable result of each alternative. Under both action alternatives there would be a landowner outreach and a conservation planning phase in which USDA-NRCS would work with private landowners to develop site-specific conservation plans outlining a combination of conservation practices⁶⁰. Conservation planning for proposed Alternative A (Preferred) would be conducted for the purpose of achieving nutrient and sediment reduction from agricultural and forested land, including riparian areas, whereas conservation planning for Alternative B would focus on establishing and maintaining riparian buffers that effectively filter nutrients and sediment from upland runoff, and would not address nutrient and sediment runoff at the source. Conservation practices would be planned and implemented on a site-specific basis, and would vary depending on the physical conditions, characteristics, and environmental constraints (e.g. endangered species, cultural resources, etc.) associated with each site. Because the specific sites are not yet known, this analysis identifies the environmental impacts that normally occur from implementing USDA-NRCS conservation practices to achieve nutrient and sediment reductions. In addition to incorporating by reference the analysis USDA-NRCS has conducted on the effects of its conservation practices, the discussion in this Draft RP/EA includes examples of the conservation practices the MS TIG expects will be implemented in the project area for the proposed alternatives and how those practices are expected to impact the environment.

3) The MS TIG Approach to Site-Specific Environmental Review for the Selected Alternative: Subsequent environmental review will occur in addition to this NEPA analytical approach to determine whether a planned site-specific action is below the maximum impacts described in this Draft RP/EA. An example of the Environmental Evaluation Worksheet used to document this review is attached as Appendix A. If the site-specific action is below the maximum impacts described in this Draft RP/EA, the analysis of the effects will be documented on the Environmental Evaluation Worksheet and the action will proceed. The Environmental Evaluation Worksheet will be routed through the MS TIG to the administrative record, where it will be publicly available. If the evaluation of the planned site-specific action indicates effects are likely to exceed the maximum impacts described in this EA, the MS TIG will undertake additional site-specific environmental review consistent with NEPA requirements and other requirements for protection of the environment. The MS TIG does not propose to take actions that would result in any significant adverse impacts on the environment.

⁵⁹ The majority of conservation practices likely to be implemented under the proposed action have been determined to fall within established NRCS categorical exclusions and therefore would not normally require preparation of an EA or EIS if implemented under NRCS program authorities. However, because this action is proposed for funding under the *DWH* NRDA Consent Decree and not all *DWH* NRDA Trustees have such categorical exclusions, the MS TIG decided to prepare this EA to aid their planning, decision-making and compliance with NEPA.

⁶⁰ The landowner outreach program, conservation planning activities and creation of conservation plans would not require project-specific environmental compliance measures described in this section.

- **4) Organization of the Affected Environment and Environmental Consequences for NR** (Nonpoint Source) Restoration Type: Guidelines for NEPA impact determinations for the PDARP/PEIS are described in Section 6.3.2 of the PDARP/PEIS and are hereby incorporated by reference. NR Alternatives A and B include development and implementation of conservation plans to reduce nutrient and sediment runoff, which would improve water quality in downstream coastal waters. Alternative A (Preferred) would include conservation practices on agricultural and forested land including riparian areas; Alternative B would include practices such as conservation buffers only in riparian areas associated with agricultural and forested land. Section 3.8 below addresses the environmental consequences of the No Action Alternative, which would allow natural recovery to proceed, followed by an overview of the proposed NR (Nonpoint Source) alternatives in section 3.9. The NEPA affected environment and environmental consequences for the NR (Nonpoint Source) Restoration Type alternatives are structured as follows:
 - Section 3.9 NR (Nonpoint Source) Alternatives Description of Common Features and Analytical Approach
 - Section 3.9.1 NR (Nonpoint Source) Alternatives A and B Affected Environment and Environmental Consequences
 - Section 3.9.1.1 Overview of Affected Environment and Environmental Consequences
 - Section 3.9.1.2 Physical Environment
 - Section 3.9.1.3 Biological Environment
 - Section 3.9.1.4 Socioeconomic Environment
 - Section 3.10 Cumulative Impacts for NR (Nonpoint Source)
 - Section 3.11 Summary of the Comparison of the Alternatives

3.8 Natural Recovery/No Action Alternative

In addition to the proposed alternatives listed above for the NR (Nonpoint Source) Restoration Type, the MS TIG evaluated the Natural Recovery/No Action Alternative (No Action). NEPA (§ 1502.14(d) requires consideration of a No Action alternative as a basis for comparison of potential environmental consequences of the action alternatives. The No Action alternative evaluation under NEPA parallels a natural recovery alternative under OPA. OPA regulations also require that "trustees must consider a natural recovery alternative in which no human intervention would be taken to directly restore injured natural resources and services to baseline" (40 CFR § 990.53(b)(2)). The OPA alternatives analysis (which includes the Natural Recovery/No Action Alternative) was presented above in Table 3.7.1-28.

Under the No Action alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes to occur, which could result in one of four outcomes for injured resources: 1) gradual recovery, 2) partial recovery, 3) no recovery, or 4) further deterioration. Although injured resources could presumably recover to or near baseline conditions under this scenario, recovery would take much longer compared to a scenario in which restoration actions were undertaken.

The No Action alternative would have no beneficial impacts to water quality via nutrient reduction because this alternative would largely result in a continuation of the conditions described in the PDARP/PEIS Chapters 3, Ecosystem Setting and Chapter 4, Injury to Natural Resources, and there would be no associated benefits to water quality by the reduction of sediments and nutrient loading.

Under the No Action alternative, some NR (Nonpoint Source) benefits could result from USDA-NRCS programs in the proposed project area, but not from the federal action being evaluated in this Draft RP/EA. The full suite of restoration benefits would not be realized solely with natural processes and without the benefit of leveraged funding opportunities and opportunity for robust monitoring and adaptive management. The No Action alternative does not meet the MS TIG's goals and objectives and clearly does not provide the significant restoration benefit to water quality via nutrient reduction that would occur through the action alternatives.

When analyzed in combination with other past, present and reasonably foreseeable future actions, the No Action alternative would provide no beneficial impacts, because existing conditions would not change in a predictable way. This alternative is not expected to contribute to short-term or long term, cumulative adverse impacts to physical resources, biological resources, or socioeconomics.

3.9 NR (Nonpoint Source) Alternatives -Description of Common Features and Analytical Approach

Both proposed NR (Nonpoint Source) alternatives would be implemented by USDA-NRCS in the Chunky-Okatibbee watershed in Mississippi for the pur

pose of improving water quality by implementing conservation practices to reduce nutrient and sediment runoff. USDA-NRCS and its conservation partners would help voluntarily participating landowners by developing conservation plans that identify natural resource concerns and conservation practices the landowner can implement to reduce nutrient and sediment runoff. The MS TIG proposes providing \$4.0 M for either of these proposed alternatives. USDA-NRCS would invest an additional \$1 M in program funds in the proposed alternative project area to implement similar conservation practices through EQIP. For proposed Alternative A and B, conservation planning would be completed with landowners in a 20,000-acre screening area shown in Figure 3.9-1.

Both alternatives would be implemented over a 5-year period with the first year consisting primarily of landowner outreach and planning. Implementation of the Ecological/NR and Soil and Water Conservation/NR conservation practices would begin in year two and continue through year five. The estimated cost for each of the alternatives is \$4.0 M.

The proposed NR (Nonpoint Source) alternatives would be implemented in portions of Newton, Lauderdale, Clarke, Neshoba, and Kemper counties, Mississippi. Lauderdale and Kemper counties contain the largest percentage of the project area. The project boundary is the Chunky-Okatibbee watershed boundary. That portion of the watershed upstream of the Okatibbee Lake Reservoir in northwestern Lauderdale County is not a part of the project area. The project location for the proposed alternatives would include conservation plans in a 20,000-acre area within the Chunky-Okatibbee watershed as shown on Figure 3.9-1




Figure 3.9-1: Upper Pascagoula River Water Quality Enhancement Project Area

The primary goal for the NR (Nonpoint Source) alternatives is water quality improvement through nutrient and sediment reduction. The health of the Gulf of Mexico depends upon the health of its estuaries, and the health of those coastal waters is influenced by land uses in the watersheds of its tributaries. In the five Gulf States, over 80 percent of the acreage is in private ownership (USDA-NRCS 2014) and is used for forestry and agriculture. These watershed-scale NR (Nonpoint Source) alternatives restore water quality impacted by the *DWH* oil spill by reducing excessive nutrients and the sediment carrying them into coastal waters. Runoff from cropland, pasture/grassland, and forests contributes excess nutrients and sediment that adversely impact the health of coastal waters of the Gulf. While agricultural and forested lands are not the sole contributors (and in many instances, not the leading contributors) of nutrients to coastal waters, there are opportunities to address this resource concern at these sources in the Pascagoula watershed. Given the success of USDA-NRCS Farm Bill programs such as EQIP and their strong acceptance by private landowners, there is a significant opportunity to implement conservation practices on private lands that would reduce the levels of nutrients and sediments entering the Gulf of Mexico from the Pascagoula watershed.

Land Use

The following Land Use categories are located in the Chunky-Okatibbee watershed:

- *Cropland* Land used primarily for the production and harvest of annual or perennial field, forage, food, fiber, horticultural, orchards, vineyards and/or energy crops (e.g.).
- Associated Agriculture Lands Land associated with farms and ranches that are not purposefully managed for food, forage (e.g.) or fiber (e.g.) and are typically associated with nearby production and/or conservation lands. This could include incidental areas such as: idle center pivot corners, odd areas, ditches and watercourses, riparian areas, field edges, seasonal and permanent wetlands, and other similar areas.
- Pasture/Grassland
 - Pasture Lands composed of introduced or domesticated native forage species that are used primarily for the production of livestock. They receive periodic renovation and/or cultural treatments, such as tillage, fertilization, mowing, weed control, and may be irrigated. They are not in rotation with crops.
 - o Grassland Land used primarily for the production of grazing animals. Includes native plant communities and those seeded to native or introduced species, or naturalized by introduced species, that are ecologically managed using range management principles.
- *Forestland* Land on which the primary vegetation is tree cover (climax, natural or introduced plant community) and use is primarily for production of wood products and/or non-timber forest products.
- **Developed Land (Urban)** Land occupied by buildings and related facilities used for residences, commercial sites, public highways, airports, and open space associated with towns and cities.
- *Water* Geographic area whose dominant characteristic is open water/permanent ice or snow. May include intermingled land, including tidal influenced coastal marsh lands.

Table 3.9-1 lists the acreages of the Land Use categories located in the Chunky-Okatibbee watershed:

Table 3.9-1: Land Use Category Acreage

National Resource Inventory⁶¹Land Use in the Chunky-Okatibbee Watershed Acres Associated Agriculture Lands 40,322 Cropland 3,580 Forestland 248,874 Pasture/Grassland 135.078 Developed Land (Urban) 45,689 Water 6,263 479,806 Total

⁶¹ https://www.nrcs.usda.gov/wps/portal/nrcs/rca/national/technical/nra/rca/ida/

Conservation Practices and Analytical Approach

Conservation Practices ⁶²are technical methods designed to help conserve soil, water, air, energy, and related plant and animal resources. Appendix B provides a complete list of conservation practices that will be available for implementation under proposed NR (Nonpoint Source) alternatives A and B. Site-specific planning would be conducted to determine which particular practice is appropriate to use given the conditions at that site.

Certain conservation practices are highlighted for the purposes of this Draft RP/EA, to provide examples of the types of effects that may result from the application of different types of conservation practices with a focus on ground-disturbing practices that have potential for adverse impacts. These practices have been grouped into two categories which are discussed below: 1-Conservation practices that provide Ecological and NR benefits (Ecological/NR conservation practices) and; 2) Conservation practices that provide soil and water conservation and NR benefits (soil and water conservation/NR conservation practices). Some conservation practices, such as Conservation Practice Standard (CPS) 342, Critical Area Planting, can fall into both categories depending on the purpose for which the practice is used.

Table 3.9-2 provides a limited number of examples of conservation practices that provide Ecological/NR Benefits. These practices will apply to both Alternatives A and B. Table 3.9-3 provides a limited number of soil and water conservation/NR Benefits which will apply primarily to Alternative A. The conservation practice standards and their associated purposes and effects analysis, which have been incorporated by reference into this RP/EA, are available on the USDA-NRCS National Handbook of Conservation Practices website at https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs143_02684_9.

Ecological/NR Conservation Practices: Examples of conservation practices that support Ecological/NR benefits (Table 3.9-2) include conservation practices implemented primarily on lands associated with agricultural operations, such as streams, riparian areas and forested lands, because these lands also can help to improve water quality by nutrient reduction via removal of sediment, nitrogen, and phosphorous. Eight conservation practices that include vegetative management, restoration of streambanks and shorelines, and structural measures to accomplish work in streams, wetlands and riparian areas are highlighted in this RP/EA as examples of conservation practices likely to be implemented under the proposed alternatives that also have potential for adverse impacts. The Streambank and Shoreline Protection practice (Conservation Practice Standard (CPS 580), Grade Stabilization Structures (CPS 410) and the Forest Stand Improvement practice (CPS 666⁶³) are ground disturbing practices and are representative of conservation practices with potential for adverse impacts and are discussed further in Section 3.9.1. Critical area planting (CPS 342) is considered to be both an Ecological/NR and Soil and Water Conservation/NR conservation practice. Any of a number of the conservation practices in Appendix B could be implemented under either of the proposed NR (Nonpoint Source) alternatives; the conservation practices funded would not be limited

⁶² https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs143 026849

⁶³ Not all applications of CPS 666 require ground disturbance, but when ground disturbance is required, these are the types of short-term adverse effects that normally occur.

to those discussed here and the actual practices selected for each project site and their anticipated impacts would be documented on the Environmental Evaluation Worksheet, described above.

Table 3.9-2: Exemplar -Ecological/NR Conservation Practices

Table 3.7	-2. Exemplai -Eco	logical/NR Conservation Fractices		1
Conservation Practice Standard Code	Conservation Practice Name	Purpose	Sediment Reduction	Nutrients (Nitrogen and Phosphorous)
314	Brush Management	Create the desired plant community consistent with the ecological site. Restore or release desired vegetative cover to protect soils, control erosion, reduce sediment, improve water quality or enhance stream flow. Maintain, modify, or enhance fish and wildlife habitat. Improve forage accessibility, quality and quantity for livestock and wildlife. Manage fuel loads to achieve desired conditions.	X	X
390	Riparian Herbaceous Cover	Provide or improve food and cover for fish, wildlife and livestock; Improve and maintain water quality. Establish and maintain habitat corridors. Increase water storage on floodplains. Reduce erosion and improve stability to stream banks and shorelines. Increase net carbon storage in the biomass and soil. Enhance pollen, nectar, and nesting habitat for pollinators. Restore, improve or maintain the desired plant communities. Dissipate stream energy and trap sediment. Enhance stream bank protection as part of stream bank soil bioengineering practices.	X	X
644	Wetland Wildlife Habitat Management	To maintain, develop, or improve wetland habitat for waterfowl, shorebirds, fur-bearers, or other wetland dependent or associated flora and fauna.	X	X
391	Riparian Forest Buffer	Create shade to lower or maintain water temperatures to improve habitat for aquatic organisms. Create or improve riparian habitat and provide a source of detritus and large woody debris. Reduce excess amounts of sediment, organic material, nutrients and pesticides in surface runoff and reduce excess nutrients and other chemicals in shallow ground water flow. Reduce pesticide drift entering the water body. Restore riparian plant communities. Increase carbon storage in plant biomass and soils.	X	X
342	Critical Area Planting	Stabilize areas with existing or expected high rates of soil erosion by wind or water. Stabilize stream and channel banks, pond and other shorelines, earthen features of structural conservation practices. Stabilize areas such as sand dunes and riparian areas.	X	
580*	Streambank and Shoreline Protection	Prevent the loss of land or damage to land uses, or facilities adjacent to the banks of streams or constructed channels, shoreline of lakes, or estuaries including the protection of known historical, archeological, and traditional cultural properties. Maintain the flow capacity of streams or channels. Reduce the offsite or downstream effects of sediment resulting from bank erosion. To improve or enhance the stream corridor for fish and wildlife habitat, aesthetics, recreation.	X	
	Grade Stabilization			
410*	Structure	Stabilize grade, reduce erosion, or improve water quality.	X	X
666*	Forest Stand Improvement	Improve and sustain forest health and productivity. Reduce damage from pests and moisture stress. Initiate forest stand regeneration. Reduce fire risk and hazard and facilitate prescribed burning. Restore or maintain natural plant communities. Improve wildlife and pollinator habitat. Alter quantity, quality, and timing of water yield. Increase or maintain carbon storage.	X	
550	-T	i de la compania del compania del compania de la compania del la compania de la compania della c		1

^{*} Practices 580, 410, and 666 are ground disturbing practices and illustrate the types of adverse environmental impacts the MS TIG expects to occur. During implementation of the selected alternative USDA-NRCS would use any of a number of the practices as shown in Appendix B. (https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs143 026849). The Section 3.7.2 describes the environmental review of all site-specific conservation plans that would be developed for the alternative that is selected.

Soil and Water Conservation/NR Practices: Examples of conservation practices that support soil and water conservation/NR benefits (Table 3.9-313) include conservation practices implemented primarily on agricultural lands including cropland and pasture/grassland, and forestland to provide nutrient reduction via removal and management of sediment, nitrogen, phosphorous and animal waste. Twelve conservation practices that include crop management measures, plantings, nutrient management, and construction measures to reduce erosion and control runoff are highlighted in this Draft RP/EA as examples of conservation practices likely to be implemented under the proposed alternatives that also have potential for adverse impacts. The Grassed Waterway practice (CPS 412), Stream Crossing (CPS 578), and Terrace (CPS 600) are ground disturbing practices and are representative of conservation practices with potential for adverse impacts and are discussed further in Section 3.9.1. Because the USDA-NRCS analysis of the effects of the conservation practices listed in Appendix B has been incorporated by reference, any of a number of those practices could be implemented under the proposed action alternative; the conservation practices funded would not be limited to those discussed here and the actual practices selected for each project site and their anticipated impacts would be documented on the Environmental Evaluation Worksheet, described above.

(https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs143 026849).

Table 3.9-3: Exemplar -Soil and Water Conservation/NR Conservation Practices

Conservation Practice Standard Code	Conservation Practice Name	Purpose	Reduction of Sediment	nutrient Reduction (Nitrogen and Phosphorous)	animal waste
412*	Grassed Waterway	Convey runoff from terraces, diversions, or other water concentrations without causing erosion or flooding. To prevent gully formation. To protect/improve water quality.	X	X	
328	Conservation Crop Rotation	Reduce sheet, rill and wind erosion. Maintain or increase soil health and organic matter content. Reduce water quality degradation due to excess nutrients. Improve soil moisture efficiency. Reduce the concentration of salts and other chemicals from saline seeps. Reduce plant pest pressures. Provide feed and forage for domestic livestock. Provide food and cover habitat for wildlife, including pollinator forage, and nesting.	X	X	
342	Critical Area Planting	Stabilize areas with existing or expected high rates of soil erosion by wind or water. Stabilize stream and channel banks, pond and other shorelines, earthen features of structural conservation practices. Stabilize areas such as sand dunes and riparian areas.	X		
329	Residue & Tillage Management	Reduce sheet, rill, and wind erosion and excessive sediment in surface waters. Reduce tillage-induced particulate emissions. Maintain or increase soil health and organic matter content. Reduce energy use.	X	X	
393	Filter Strip	Reduce suspended solids and associated contaminants in runoff and excessive sediment in surface waters. Reduce dissolved contaminant loadings in runoff. Reduce suspended solids and associated contaminants in irrigation tailwater and excessive sediment in surface waters.	X	X	
340	Cover Crop	Reduce erosion from wind and water. Maintain or increase soil health and organic matter content. Reduce water quality degradation by utilizing excessive soil nutrients. Suppress excessive weed pressures and break pest cycles. Improve soil moisture use efficiency. Minimize soil compaction.	X	X	

Conservation Practice Standard Code	Conservation Practice Name	Purpose	Reduction of Sediment	nutrient Reduction (Nitrogen and Phosphorous)	animal waste
	Livestock Shelter	To provide protection for livestock from excessive heat, wind, cold. Protect surface waters from nutrient and pathogen loading. Protect wooded areas from accelerated erosion and excessive nutrient deposition by providing alternative livestock shelter/shade location. Improve the distribution of grazing livestock to enhance wildlife habitat, reduce over-used areas, or correct other resource concerns			
576	Structure	resulting from improper livestock distribution.	X	X	X
578*	Stream Crossing	Provide access to another land unit. Improve water quality by reducing sediment, nutrient, organic, and inorganic loading of the stream. Reduce streambank and streambed erosion.	X	X	X
600*	Terrace	Reduce erosion and trap sediment. Retain runoff for moisture conservation.	X	X	
590	Nutrient Management	Budget, supply, and conserve nutrients for plant production. To minimize agricultural nonpoint source pollution of surface and groundwater resources. To properly utilize manure or organic byproducts as a plant nutrient source. To protect air quality by reducing odors, nitrogen emissions (ammonia, oxides of nitrogen), and the formation of atmospheric particulates. To maintain or improve the physical, chemical, and biological condition of soil.		X	
528	Prescribed Grazing	Improve or maintain desired species composition and vigor of plant communities. Improve or maintain quantity and quality of forage for grazing and browsing animals' health and productivity. Improve or maintain surface and/or subsurface water quality and quantity. Improve or maintain riparian and watershed function. Reduce accelerated soil erosion, and maintain or improve soil condition. Improve or maintain the quantity and quality of food and/or cover available for wildlife. Manage fine fuel loads to achieve desired conditions.	X	X	
317	Composting Facility	Reduce water pollution potential and improve handling characteristics of organic waste solids, reuse organic waste as animal bedding, or use as a soil amendment that provides soil conditioning, slow-release plant-available nutrients and plant disease suppression.		X	X

^{*} Practices 412, 578, and 600 are ground disturbing practices and illustrate the types of adverse the MS TIG expects to occur. During implementation of the selected alternative USDA-NRCS would use any of a number of the practices as shown in Appendix B. (https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs143 026849). The Section 3.7.2 describes the environmental review of all site-specific conservation plans that would be developed for the alternative that is selected.

Table 3.9-4 lists the land use categories, acreages, and the categories of conservation practices that potentially could be prescribed.

Table 3.9-4: Potential Conservation Practice by Land Use Category

Land Use	Acres	Planning Area ⁶⁴	Ecological/NR Conservation Practices	Soil and Water Conservation /NR Practices
Associated Agriculture Lands	40,322		X	X
Cropland	3,580	2,000		X
Pasture/Grassland	135,078	11,000		X
Forestland	248,874	7,000	X	X
Developed Land (Urban)	45,689			
Open Water	6,263			
Total	479,806	20,000		

3.9.1 NR (Nonpoint Source) Alternatives A and B: -Affected Environment and Environmental Consequences

This section describes the affected environment and the environmental consequences for proposed NR (Nonpoint Source) alternatives A and B within the Chunky-Okatibbee watershed. The project area for the proposed alternatives is depicted in Figure 3.9-1 for Alternative A (Preferred): Upper Pascagoula Water Quality Enhancement Project and Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan.

Alternative A (Preferred): Upper Pascagoula River Water Quality Enhancement Project

If selected, Alternative A, Upper Pascagoula Water Quality Enhancement Project (Preferred) would be implemented by USDA-NRCS for the purpose of improving water quality through the development and implementation of conservation plans to reduce nutrient and sediment runoff closest to the source of soil erosion and nutrient application as well as in riparian areas. The Upper Pascagoula River Water Quality Enhancement project (Alternative A-Preferred) would include implementation of conservation practices from both the Ecological/NR and Soil and Water Conservation/NR categories described in Section 3.9 (Table 3.9-2, Table 3.9-3; Appendix B). USDA-NRCS would provide outreach and technical assistance to voluntary participants (landowners) to develop conservation plans and would use all available conservation practices typically planned and funded by USDA-NRCS programs. USDA-NRCS would develop conservation plans within a 20,000-acre area with a priority on opportunities that are within one mile of tributaries (See Table 3.9-4). Conservation practices would be implemented on cropland, pasture/grassland, forestland, and associated agriculture lands within the Okatibbee-Chunky watersheds with emphasis given to properties bordering rivers and streams.. The MS TIG would allocate \$4.0 M from the NR restoration type for this alternative.

169

⁶⁴ Estimated planning area is based on preliminary project development and may be modified (increased or decreased) during project implementation considering factors including but not limited to: landowner participation, proximity of existing conservation practices, costs, and opportunities for implementation of conservation actions and practices.

Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan (Proposed Action)

The Pascagoula River Basin Riparian Buffer Maintenance Plan (Alternative B) would also be implemented by USDA-NRCS for the purpose of improving water quality through the development and implementation of conservation plans to reduce nutrient and sediment runoff by focusing conservation practices such as conservation buffers in riparian areas. This Alternative would include implementation of Ecological/NR conservation practices as described in Section 3.9 (Table 3.9-2; Appendix B). The USDA-NRCS would provide outreach and technical assistance to voluntary participants (landowners) to develop conservation plans in riparian areas and would use all available conservation practices typically planned and funded by USDA programs. The USDA would develop conservation plans within a 20,000-acre area with priority on opportunities that are within one mile of tributaries. Conservation practices would be implemented in riparian areas within forest and associated agriculture lands on farmsteads in the Chunky- Okatibbee watersheds in Mississippi. Similar to Alternative A, conservation planning would be completed within a 20,000-acre area with a priority on opportunities that are within one mile of tributaries (See Table 3.9-4). Alternative B differs from Alternative A only in that the conservation practices would primarily be Ecological/NR practices (Appendix B) that would be implemented in riparian areas within associated agriculture lands and forestland in the Chunky- Okatibbee watersheds in Mississippi. The MS TIG would allocate \$4.0 M from the NR restoration type for this alternative.

Exemplar Conservation Practices Analyzed in this Plan: Table 3.9-5 provides a description of the types of work that would be carried out in order to implement each of the exemplar conservation practices discussed in this Draft RP/EA, including both the Ecological/NR conservation practices and Soil and Water Conservation/NR conservation practices. The affected environment and environmental consequences for these exemplar conservation practices are included in sections 3.9.1.1 through 3.9.1.4. Appendix B provides the list of conservation practices contemplated for proposed NR (Nonpoint Source) Alternatives A and B. Appendix C provides a conservation practice network effects diagram for the example practices analyzed in this Draft RP/EA.

Table 3.9-5: Example Ground-Disturbing Conservation Practices-Description of Work

Practice Code	Conservation Practice Name	Purpose/Description of work
		Exemplar Ecological/NR Conservation Practices (Alternative A and B)
		Purpose/Description of Work: Prevent the loss of land or damage to land uses, or facilities adjacent to the banks of streams or constructed channels, shoreline of lakes, or estuaries including the protection of known historical, archeological, and traditional cultural properties. Maintain the flow capacity of streams or channels. Reduce the offsite or downstream effects of sediment resulting from bank erosion. To improve or enhance the stream corridor for fish and wildlife habitat, aesthetics, recreation. Site-specific work would include
580	Streambank and Shoreline Protection	treatment(s) used to stabilize and protect banks of streams or constructed channels, and shorelines of lakes, reservoirs, or estuaries. Heavy equipment would be used to regrade selected shorelines and streambanks and deposit erosion control materials such as rip rap or green controls. The site will be replanted with native herbaceous/tree species.
410	Grade Stabilization Structure	Purpose/Description of Work: Stabilize grade, reduce erosion, or improve water quality. Site-specific construction would include installation of grade stabilization structure(s) used to control the grade in natural or constructed channels. Heavy equipment would be used to regrade selected streams and install grade control structures such as embankments, drop/chute/box inlet drop spillways, side-inlet, open weir, or pipe-drop drainage structures. The site will be replanted with native herbaceous/tree species.
	Forest Stand	Purpose/Description of work: Improve and sustain forest health and productivity. Reduce damage from pests and moisture stress. Initiate forest stand regeneration. Reduce fire risk and hazard and facilitate prescribed burning. Restore or maintain natural plant communities. Improve wildlife and pollinator habitat. Alter quantity, quality, and timing of water yield. Increase or maintain carbon storage. Site-specific work would include the manipulation of species composition, stand structure, or stand density by cutting or killing
666	Improvement	selected trees or understory vegetation to achieve desired forest conditions or obtain ecosystem services.

Practice Code	Conservation Practice Name	Purpose/Description of work
		Improvement, such as invasive or unwanted species removal, thinning, and planting/seeding would potentially utilize heavy equipment. Treatments could include, but are not limited to, mowing, planting/seeding, felling, tilling, or chemical treatment. Planting could include the use of seed drills or other planting/seeding
		equipment.
	Exe	emplar Soil and Water Conservation/NR Conservation Practices (Alternative A)
		Purpose/Description of work: Convey runoff from terraces, diversions, or other water concentrations without causing erosion or flooding. To prevent gully formation. To protect/improve water quality. Sitespecific work would include the construction of a shaped or graded channel that is established with suitable vegetation to convey surface water at a non-erosive velocity using a broad and shallow cross section to a stable outlet. Selected sites would be prepared for planting by potentially using equipment to remove
	Grassed	vegetation and other debris. Site preparation treatments could include tilling, or chemical treatment. Planting
412	Waterway	could include the use of seed drills or other planting/seeding equipment.
578	Stream Crossing	Purpose/Description of work: Provide access to another land unit. Improve water quality by reducing sediment, nutrient, organic, and inorganic loading of the stream. Reduce streambank and streambed erosion. Site-specific work would include construction of a stabilized area or structure constructed across a stream to provide a travel way for people, livestock, equipment, or vehicles. A ford, bridge, or culvert structure could be installed. Heavy equipment would be used to regrade the stream and construct the structure. The area will be replanted with native vegetation.
378	Sucam Clossing	Purpose/Description of work: Reduce erosion and trap sediment. Retain runoff for moisture conservation.
		Site specific work would include construction of an earth embankment, or a combination ridge and channel, constructed across the field slope. Heavy equipment would be used to regrade the selected area into a terrace
600	Terrace	system.

Exemplar conservation practices are ground disturbing practices and are representative of some of the most impacting practices and are analyzed for environmental impacts in the Draft RP/EA. During implementation of the selected alternative USDA-NRCS would use any of a number of their practices.

(https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs143_026849). The Section 3.7.2 describes the environmental review of all site-specific conservation plans that would be developed for the alternative that is selected.

Best Practices: The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS to avoid and minimize impacts to resources. Best practices listed in the PDARP/PEIS are intended to evolve as an adaptive management component of implementing the PDARP/PEIS; as such, the appendix to the PDARP/PEIS is a living document. As new best practices are established, existing best practices are refined, or new techniques and information are informed by implementation, these measures will be added to or updated in the relevant websites identified in the appendix of the PDARP. In this capacity, new projects will have available the current range of best practices to support project design and implementation. In addition to PDARP/PEIS best practices, the MS TIG could develop best practices for site-specific conservation practices in different locations due to differences in relevant site conditions.

3.9.1.1 Overview of Affected Environment and Environmental Consequences

This analysis incorporates by reference the relevant portions of the affected environment description from Section 3.3.2 for Water Quality from the PDARP/PEIS. Likewise, the PDARP/PEIS provides programmatic evaluation of the environmental consequences from conduct of the restoration approaches "Reduce nutrient loads to coastal watersheds" considered in this plan. Those evaluations are incorporated by reference here, from section 6.4.3 of PDARP/PEIS. Tiering from that analysis, this section presents the Affected Environment of the NR (Nonpoint source) proposed alternatives and environmental consequences of the proposed actions in context of the project-specific affected environment.

The programmatic analysis looked at a series of resources as part of the physical, biological, , and socioeconomic environment. As appropriate in a tiered analysis, the evaluation of each alternative

focuses on the specific resources with a potential to be affected by the proposed project. To avoid redundant or unnecessary information, resources that are not expected to be affected are evaluated summarily in the respective sections. These resources include, air quality and greenhouse gas emissions, noise, invasive species, marine and estuarine fauna, infrastructure, tourism and recreation, fisheries and aquaculture, marine transportation, land and marine management and aesthetics and visual resources which will be discussed in Sections 3.9.1.2, 3.9.1.3, and 3.9.1.4.

3.9.1.2 Physical Environment

<u>Introduction to Affected Environment (Physical Environment)</u>: Geology and Substrates and Hydrology and Water Quality are discussed in this section. PDARP/PEIS sections 3.3.1, 3.3.2, 3.3.3 and 3.5.1 are incorporated by reference here. The affected environment for the proposed alternatives physical environment for the is described in respective sections below.

<u>Programmatic Review of Environmental Consequences (Physical Environment)</u>: Sections 6.4.3.1 of the PDARP/PEIS describe the impacts to Physical Resources for the relevant restoration approaches and are incorporated by reference and briefly described here.

<u>PDARP/PEIS</u> consequences related to geology and substrates and water resources: Some agricultural best practices include small-scale construction projects (e.g., to manage manure and runoff from feedlots). Therefore, during construction, short-term, minor adverse impacts on geology, substrate, hydrology, surface and ground water quality (e.g., nutrients, fertilizers, pesticides, total suspended solids in runoff, and high-conductivity ground water) would be anticipated. Short-term adverse impacts would be minimized by implementing best practices. Long-term benefits are expected to result because these conservation practices would reduce nutrients, slow erosion, stabilize soils, improve water quality, and increase ground water recharge.

As appropriate in a tiered analysis, the evaluation of the proposed alternative focuses on the specific resources with a potential to be affected. Air quality and greenhouse gas emissions and noise impacts for the proposed alternatives would be negligible to minor. To avoid redundant or unnecessary information, these resources are evaluated here.

Air Quality and Greenhouse Gas Emissions: Counties where the proposed alternative project area are located are classified as in attainment, meaning criteria air pollutants do not exceed National Ambient Air Quality Standards (NAAQS)⁶⁵-need citation. The primary sources of emissions during project implementation would include equipment operation such as tractors, dozers, and all-terrain vehicles associated with earth moving, seeding, planting, habitat management and small construction. Implementation of conservation practices would be within the range of normal farmstead operation, which do not impact air quality. Conservation practices would occur seasonally, and would likely not occur simultaneously. Whether activities occurred simultaneously or incrementally, the proposed alternatives would have no long-term adverse impacts on air quality or to emissions of greenhouse gases. Conservation practices on forested areas could result in a long-term beneficial impact on air quality resulting from more vigorous long-standing forested areas, which help to sequester carbon.

_

 $^{^{65}\} https://www.deq.state.ms.us/MDEQ.nsf/pdf/Air_2015AirQualityDataSummary/\$File/2015\%20Air\%20Quality\%20Data\%20Summary.pdf$

In addition, the following best practices would be implemented, to the extent practicable, for the proposed alternatives:

- Shut down idling restoration equipment, if feasible.
- Locate staging areas as close to restoration sites as practicable to minimize driving distances between staging areas and restoration sites.
- Encourage the use of the proper size of equipment for the job to maximize energy efficiency.
- Encourage the use of alternative fuels or power sources for generators at restoration sites, such as propane or solar power, or use electrical power where practicable.

Noise: There would be short-term minor adverse noise impacts from equipment and operations associated with the installation of various conservation practices. Conservation practices would be implemented sporadically and seasonally and on private land, not near densely populated areas. The types of noise produced would be typical of farmstead operations (e.g. plowing, harvesting, small earthmoving activities, land clearing). The operations would be short-term and remote from nearby receptors.

For the physical environment, the following resources are further analyzed in this section:

- Geology and Substrates
- Water Quality and Hydrology

3.9.1.2.1 Geology and Substrates

Affected Environment

The project area for the proposed alternatives is located within the Tombigbee Hills physiographic region. Sediments are generally composed of sands, clays, and gravels of the Tuscaloosa and Eutaw formations (Cretaceous). The soils are highly weathered and acid and include very old ultisols, few alfisols, entisols in stream drainages soil orders (Stewart 2003).

Topography in the area varies from undulating broad plateau areas between major stream systems to rugged dissected uplands, characterized by steep side slopes and narrow ridgetops. All the major streams have fairly broad valleys with floodplains bordered by one or more low terraces. Okatibee Creek and Chunky River flow into the Chickasaway River, which flows into the Pascagoula River (USDA 1983).

According to national land cover database, land use within the proposed alternatives project area is 3,580 acres cropland, 40,322 acres as associated agriculture lands, and 135,078 acres is used for pasture or to grow hay.

Environmental Consequences for NR Proposed Alternatives A (Preferred) and B

Table 3.9-6 provides a summary of the environmental consequences associated with representative exemplar conservation practices proposed for implementation in the project area for Alternative A (Preferred): Upper Pascagoula Water Quality Enhancement and Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan. There would be no adverse impacts to geology as a result of the project; soil impacts are summarized below.

Table 3.9-6: Summary of Soil Impacts

		Alternative A: Upper Pascagoula Water Quality Enhancement- (Preferred)					Alternative B: Pascagoula River B Riparian Buffer Maintenance Pl		
Practice	Conservatio	Adverse	Adverse	Beneficial		Adverse	Adverse	Beneficial	
Code	n Practice Name	Impact Duration	Impact Intensity	Impact Duration		Impact Duration	Impact Intensity	Impact Duration	
	Ty				tha	t Provide NR Bei		Burunon	
	Streambank and Shoreline		minor to	<u> </u>			minor to		
580	Protection	short-term	moderate	long-term		short-term	moderate	long-term	
410	Grade Stabilization Structure	short-term	minor to moderate	long-term		short-term	minor to moderate	long-term	
666	Forest Stand Improvement	short-term	minor			short-term	minor		
	Typical Conserv	vation Practic	es (Soils and	Water Conser	vat	ion/NR) that pr	ovide NR Benefi	ts	
412	Grassed Waterway	short-term	minor to moderate	long-term					
578	Stream Crossing	short-term	minor to moderate	long-term					
600	Terrace	short-term	minor to moderate	long-term					

Conservation Practices (Ecological/NR)

Streambank and Shoreline Protection (580): This practice would be applied to stabilize and protect banks of streams or constructed channels and shorelines of open water bodies and can reduce the offsite effects of sediment resulting from bank erosion. There would be short-term, minor to moderate adverse impacts from grading, reshaping, and planting of stream banks, ponds, lakes, and other aquatic systems. There would be long-term beneficial impacts as stabilization would result in reducing the off-site, downstream effects of sediment, nutrients, and organic material into surface waters. Areas would be replanted with native vegetation and or seeded to prevent erosion after bank regrading. Erosion control plans would be implemented during and after construction. Impacts would be applicable to Alternative A and Alternative B.

Grade Stabilization Structure (410): This practice would be used for grade stabilization and preventing formation of advance gullies and headcuts. There would be short-term minor to moderate adverse impacts from soil excavation, grading, to construct or install grade stabilization structures including berms, rip rap, and hard structures. The majority of these would be installed in agricultural fields, and could be installed in drainageways or tributaries. There would be long term beneficial impacts to geology and soils from prevention of gully formation, reduction of soils, and drainageway stabilization. Areas would be replanted or seeded to prevent erosion after bank regrading. Erosion control plans would be implemented during and after construction. Impacts would be applicable to Alternative A and Alternative B.

<u>Forest Stand Improvement (666)</u>: There would be short-term, minor impacts to soils from use of small equipment to access and complete operations which would include use of chainsaws to cut or kill trees or selected understory vegetation, and dragging of felled materials. Impacts would be applicable to Alternative A and Alternative B.

Conservation Practices (Soil and Water Conservation/NR)

Grassed Waterway (412): There would be short-term, minor to moderate adverse impacts from shaping or grading a channel and grading to form or install a stable outlet. The area would be replanted, where possible with vegetation that would serve to reduce erosion and provide benefit to wildlife. There would be long-term benefit from controlling and managing flow to prevent soil erosion, increases in soil infiltration and increased soil biological activity, and trapping of sediments in the waterways. The grassed waterway practice would be implemented primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

Stream Crossing (578): There would be short-term, minor to moderate impacts to the streambed from stabilizing an area for designated crossing, installation of culverts or small bridges. In some cases, fences would be constructed to direct livestock or people to crossing. There would be long-term beneficial impacts resulting from livestock traversing the stream at one stabilized location versus traversing the stream in various location. Fences would prevent riparian area grazing and resultant animal waste/nutrient contribution in and near waterways. This practice would be implemented primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

<u>Terrace (600)</u>: This practice would be used to create an earth embankment, channel, or a combination of ridge and channel constructed across a slope to intercept runoff. There would be short-term minor to moderate, adverse impacts from soil excavation, grading, to construct or install terraces. The majority of these would be installed in agricultural fields. There would be long-term beneficial impacts to geology and soils from prevention of gully formation and reduction of soils erosion. Areas not in crop production would be replanted or seeded to prevent erosion after bank regrading. Erosion control plans would be implemented during and after construction. This practice would be implemented primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific conservation practices in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to geology and substrates (soils):

- Impacts due to conservation practice implementation would be minimized by limiting operations to favorable conditions when soils are not saturated, and minimizing the disturbance footprint. Any practice that involves disturbance of wetlands in order to complete the intended beneficial long-term goal would need authorization by the U.S. Army Corps of Engineers. A Nationwide Permit 27 Aquatic Habitat Restoration, Establishment, and Enhancement Activities would be obtained with adherence to any permit conditions.
- Develop and implement an erosion control plan to minimize soil erosion during and after construction and where possible use vegetative buffers (100 feet or greater), revegetate with native species or annual grasses, and conduct work during dry seasons.
- Develop and implement a spill prevention and response plan, including conducting daily inspections of all construction and related equipment to ensure there are no leaks of antifreeze, hydraulic fluid, or other substances and cleaning and sealing all equipment that

- would be used in the water to rid it of chemical residue. Develop a contract stipulation to disallow use of any leaking equipment or vehicles.
- Prohibit use of hazardous materials, such as lead paint, creosote, pentachlorophenol, and other wood preservatives during construction in, over or adjacent to, sensitive sites during construction and routine maintenance.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.8). The No Action alternative would not provide benefits to soils or geology when compared to Alternatives A and B. The No Action alternative does not meet the MS TIG's goals and objectives and clearly does not provide the significant restoration benefit to water quality via nutrient reduction that would occur through the action alternatives.

3.9.1.2.2 Hydrology and Water Quality

Section 3.3.2 of the PDARP/PEIS addresses river flows on the Northern Gulf geography and water quality. Section 6.14.2 discusses future sea level rise, storm surge and storm intensity projections and is incorporated by reference here. The affected environment consists of numerous named and unnamed tributaries in the Upper Pascagoula River system as well as various farm ponds, lakes, and wetlands. Mississippi's water quality standards specify the appropriate levels for which various water quality parameters or indicators support a water body's designated use(s). Each use assessed for a water body is determined to be either "Attaining" or "Not Attaining" in accordance with the applicable water quality standards and U.S. Environmental Protection Agency (EPA) guidelines for assessments pursuant to §305(b). A water body's use is said to be impaired when—based on current and reliable site-specific data of sufficient quantity, quality, and frequency of collection—it is not attaining its designated use(s). Where data and information of appropriate quality and quantity indicate non-attainment of a designated use or uses for an assessed water body, the water body would be placed on the Mississippi 2014 Section 303(d) List of Impaired Water Bodies (MDEQ 2014).

The proposed alternatives are located in the Chunky-Okatibbee subbasin. It has a drainage area of approximately 581,002 acres and includes portions of Lauderdale, Newton, Clark, Jasper, and Neshoba counties. Named tributaries within the Chunky-Okatibbee subbasin include (but are not limited to) the Chunky River, Okatibbee Creek, Sowashee Creek, Tallushua Creek, Tallahatta Creek, and Suqualena Creek. All of which are part of the Pascagoula River system. Major rivers carry high sediment loads into the Mississippi Sound. Pollution from agriculture, improperly treated sewage, roadways, accidental spills, industry discharges, and other sources also affect the health of the habitats.

The waters in this area are classified by the State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters (MDEQ 2012) as "public water supply", "recreation", and "fish and wildlife". The following water bodies are listed as impaired on the State of Mississippi 303(d) list (Figure X, MDEQ 2014):

- Sosashee Creek: Total Nitrogen, Total Phosphorus
- Northern Reach of Okatibbe Creek: Biological Impairment, pH, Total Nitrogen
- Southern Reach of Okatibbee Creek: Biological Impairment
- Tallashua Creek: Biological Impairment

• Chunky Creek: Biological Impairment

• Anderson Brand: Biological Impairment

Floodplains

There are three flood zone categories within the proposed alternative(s) project area: A, AE, and X. Zone A is defined as Areas subject to inundation by the 1-percent-annual-chance flood event generally determined using approximate methodologies. Because detailed hydraulic analyses have not been performed, there are no Base Flood Elevations (BFEs). Mandatory flood insurance purchase requirements and floodplain management standards apply. Zone AE is defined as "Base Flood Elevations Determined". Upland areas are mostly Zone X. Zone X are defined as "Areas of 0.2% annual change flood; areas of 1% annual chance flood with average depths of less than 1 foot or with drainage areas less than 1 square mile; and areas protected by levees from 1% annual chance flood".

Wetlands

Wetlands in the proposed alternative(s) project area are a mix of palustrine emergent, palustrine forested, and palustrine scrub-shrub wetlands. They are generally located in shallow depressions at lower elevations or within a floodplain, as fringe wetlands are open water, or adjacent to tributaries or oxbow or lowland features. They can originate from hill seeps, or hold water for long periods of time after rain or flood events.

The National Wetland Inventory identifies over 56,871 acres of land within the total project area for the proposed alternatives (482,662 acres) as wetland or open water.

Environmental Consequences for NR Proposed Alternatives A (Preferred) and B

All of the conservation practices would be implemented voluntarily on privately owned land. Detailed information on the conservation Practices including practice standards, flow charts, and environmental effects can be found at

https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs143_02684_9.

Environmental consequences affecting hydrology, water quality, wetlands, and floodplains are discussed below.

Hydrology

Table 3.9-7 provides a the environmental consequences for representative exemplar conservation practices proposed for implementation in the project area for Alternative A (Preferred): Upper Pascagoula Water Quality Enhancement and Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan.

Table 3.9-7: Summary of Hydrology Impacts

		Alternative A: Upper Pascagoula Water Quality Enhancement- (Preferred)					B: Pascagoula R Buffer Maintena	
Practice Code	Conservation Practice Name	Adverse Impact Duration	Adverse Impact Intensity	Beneficial Impact Duration		Adverse Impact Duration	Adverse Impact Intensity	Beneficial Impact Duration
	Tvi	nical Conservati	on Practices (F	Cological/NR)- th	nat Pro	ovide NR Benefits	S	
	Streambank and	Jieur Conservati	on Tructices (1	leological/1(1t) th		Ovide I (It Delicite)		
	Shoreline						minor to	
580	Protection	short-term	minor	long-term		short-term	moderate	long-term
410	Grade Stabilization	short-term	minor	long-term		short-term	minor to moderate	long-term
	Forest Stand							
666	Improvement	short-term	minor	long-term		short-term	minor	long-term
	Typical Conserv	ation Practice	s (Soils and V	Vater Conserva	tion/l	NR) that provid	le NR Benefits	
412	Grassed Waterway			long-term				
578	Stream Crossing	long-term	minor	long-term				
600	Terrace	short-term	minor to moderate	long-term				

Conservation Practices (Ecological/NR)

Streambank and Shoreline Protection (580): This practice would be applied to stabilize and protect banks of streams or constructed channels and shorelines of open water bodies. There would be short-term, minor, adverse impacts from grading, reshaping, and planting of stream banks, ponds, lakes, and other aquatic systems. These impacts would result from altered hydrologic flow in the stream during construction. There would be long-term beneficial impacts as this practice would result in restoring stream hydrology, and provide the hydrologic benefits of riparian vegetation including staging of stormwater flows. Areas would be replanted with native vegetation and or seeded to restore streambank vegetation. Erosion control plans would be implemented during and after construction. Impacts would be applicable to Alternative A and Alternative B.

Grade Stabilization Structure (410): This practice would be used for grade stabilization, prevent formation of advance gullies and headcuts. There would be short-term, minor, adverse impacts from soil excavation, grading, to construct or install grade stabilization structures including berms, rip rap, and hard structures. The majority of these would be installed in agricultural fields, and could be installed in drainageways or tributaries. There would be long-term, beneficial impacts to hydrology from prevention of gully formation, prevention of headcutting, and drainageway destabilization. Areas would be replanted or seeded to prevent erosion and gully formation after regrading. Erosion control plans would be implemented during and after construction. Impacts would be applicable to Alternative A and Alternative B.

<u>Forest Stand Improvement (666)</u>: There would be short-term, minor, impacts to hydrology from use of small equipment to access and complete operations which would include use of chainsaws to cut or kill trees or selected understory vegetation, and dragging of felled materials. Between the time that any vegetation is cleared to the time that ground cover regrows, runoff and increased hydrology could occur. There would be long-term beneficial impacts from healthier forest stands. Removal of

overstory canopy can increase the amount and vigor of ground cover, slowing runoff and increasing infiltration. Impacts would be applicable to Alternative A and Alternative B.

Conservation Practices (Soil and Water Conservation/NR)

Grassed Waterway (412): There would be no adverse impacts to hydrology from shaping or grading a channel and grading to form or install a stable outlet. The area would be replanted, where possible with vegetation that would serve to reduce erosion and provide benefit to wildlife. There would be long-term, benefits from controlling and managing flow to slow hydrologic flow and prevent soil erosion. The grassed waterway practice would be done primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

Stream Crossing (578): There would be long-term, minor, adverse impacts to the streambed from stabilizing an area for designated crossing, installation of culverts of small bridges. There would be long-term beneficial impacts resulting from livestock traversing the stream at one stabilized location versus traversing the stream in various locations which could result in compromise of stream banks. If fences are installed with the crossing, it would prevent riparian area grazing and ground cover grazing that would result in decreased infiltration. This practice would be done primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

<u>Terrace (600)</u>: This practice would be used to create an earth embankment, channel, or a combination of ridge and channel constructed across a slope to intercept runoff. There would be short-term, minor to moderate, adverse impacts to hydrology as a result of soil excavation and grading to construct or install terraces. The majority of terraces would be installed in agricultural fields. There would be long-term, beneficial impacts to hydrology from the reduction of runoff, increased water storage and prevention of gully formation. Areas not in crop production would be replanted or seeded to prevent erosion after bank regrading. Erosion control plans would be implemented during and after construction. This practice would be implemented primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

Water Quality

Table 3.9-8 provides a summary of the environmental consequences for representative conservation practices proposed for implementation in the project area for Alternative A (Preferred): Upper Pascagoula Water Quality Enhancement and Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan.

Table 3.9-8: Summary of Water Quality Impacts

		Alternative A: Upper Pascagoula Water Quality Enhancement- Preferred Adverse Adverse Beneficial Impact Impact Impact Duration Intensity Duration				e B: Pascagoula R Buffer Maintena	
Practice Code	Conservation Practice Name				Adverse Impact Duration	Adverse Impact Intensity	Beneficial Impact Duration
			WATER	QUALITY			
Typical Conser	vation Practices (Ecolo	ogical/NR) that l	Provide NR Be	nefits			
580	Streambank and Shoreline Protection	short-term	minor	long-term	short-term	minor to moderate	long-term
410	Grade Stabilization	short-term	minor	long-term	short-term	minor to moderate	long-term

		Alternative A: Upper Pascagoula Water Quality Enhancement- Preferred					B: Pascagoula R Buffer Maintena			
Practice Code	Conservation Practice Name	Adverse Impact Duration	Beneficial Impact Duration		Adverse Impact Duration	Adverse Impact Intensity	Beneficial Impact Duration			
666	Forest Stand Improvement			long-term				long-term		
Typical Conso	ervation Practices (S	Soils and Wate	r Conservatio	on/NR) that pr	ovide NR Benefits					
412	Grassed Waterway	short-term	minor to moderate	long-term						
578	Stream Crossing	short-term	minor	long-term						
600	Terrace	short-term	minor- to moderate	long-term						

Conservation Practices (Ecological/NR)

Streambank and Shoreline Protection (580): This practice would be applied to stabilize and protect banks of streams or constructed channels and shorelines of open water bodies. There would be short-term, minor, adverse impacts from the potential for increased erosion during grading, reshaping, and planting of stream banks, ponds, lakes, and other aquatic systems. There would be long-term, beneficial impacts as this practice would result in stabilizing the waterbody and preventing further erosion. Areas would be replanted with native vegetation and or seeded to prevent erosion. Erosion control plans would be implemented during and after construction. Impacts would be applicable to Alternative A and Alternative B.

Grade Stabilization Structure (410): There would be short-term, minor adverse impacts from the potential for increased erosion resulting from soil excavation, grading, to construct or install grade stabilization structures including berms, rip rap, and hard structures. The majority of these would be installed in agricultural fields, and could be installed in drainageways or tributaries. There would be long-term, beneficial impacts from drainageway stabilization. Areas would be replanted or seeded to prevent erosion and gully formation after bank regrading. Erosion control plans would be implemented during and after construction. Impacts would be applicable to Alternative A and Alternative B.

<u>Forest Stand Improvement (666)</u>: There would be no adverse impacts to water quality. There would be long-term benefits as a result of this practice. Reduction of overstory canopy can increase the amount and vigor of ground cover, slowing runoff and increasing infiltration. Managing for desirable plant health and vigor reduces the need for pesticide applications. Reduced stand density can increase infiltration and leaching of salts. Removal of canopy/woody vegetation exposes the site and increases mortality of pathogens that would have otherwise entered surface water. Impacts would be applicable to Alternative A and Alternative B.

Conservation Practices (Soil and Water Conservation/NR):

<u>Grassed Waterway (412)</u>: There would be short-term, minor to moderate, adverse impacts from the potential of increased erosion as a result of shaping or grading a channel and grading to form or install a stable outlet. These impacts would last until vegetation regrows. The area would be replanted, where possible, with vegetation that would serve to reduce erosion and provide benefit to

wildlife. There would be long-term benefits from increased infiltration, filtration of water before it reaches the waterway, and erosion prevention. The grassed waterway practice would be implemented primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

Stream Crossing (578): There would be short-term, minor impacts from the potential of increased erosion as a result of earth moving required to install a stream crossing. There would be long-term, beneficial impacts resulting from livestock traversing the stream at one stabilized location versus traversing the stream in various locations. If fences were installed with the practice, they would prevent riparian area grazing and ground cover grazing that would result in decreased infiltration. This practice would be implemented primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

<u>Terrace (600)</u>: This practice would be used to create an earth embankment, channel, or a combination of ridge and channel constructed across a slope to intercept runoff. There would be short-term, minor to moderate, adverse impacts from the potential of increased erosion during soil excavation and grading to construct or install terraces. The majority of these would be installed in agricultural fields. There would be long-term, beneficial impacts from the reduction of runoff that could contain contaminants, and prevention of erosion. Areas not in crop production would be replanted or seeded to prevent erosion after bank regrading. Erosion control plans would be implemented during and after construction. The grassed waterway practice would be implemented primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

Floodplains

Propose alternative(s) activities would not result in a detectable change to natural and beneficial floodplain values. Stream crossings and grade stabilization installed in streams would be constructed would be designed so as not to cause an appreciable rise in floodwaters.

Wetlands

Various Conservation Practices could have impacts to wetlands. The impacts could be from regrading or clearing areas for streambank stabilization or other similar Conservation Practices. Prior to all Conservation Practices that would impact wetlands, coordination with USACE would be conducted to determine the extent of the wetlands and potential impacts and to secure authorization for proposed wetland fill and in-water activities. Table 3.9-9 provides a summary of environmental consequences for representative conservation practices proposed for implementation in the project area for Alternative A (Preferred): Upper Pascagoula Water Quality Enhancement and Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan.

Table 3.9-9: Summary of Impacts to Wetlands

			ve A: Upper F Quality Enhar Preferred			Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan					
Practice Code	Conservation Practice Name	Adverse Adverse Beneficial Impact Impact Impact Duration Intensity Duration				Adverse Impact Duration	Adverse Impact Intensity	Beneficial Impact Duration			
Typical Conservation Practices (Ecological/NR) that Provide NR Benefits											
	Streambank and			Lites (Ecological) (It) that I			Ī				
7 00	Shoreline		minor to				minor to				
580	Protection	short-term	moderate	long-term		short-term moderate		long-term			
410	Grade Stabilization	short-term	minor to moderate	long-term		short-term	minor to moderate	long-term			
666	Forest Stand Improvement	short-term	minor	long-term		short-term	minor	long-term			
	Typical Conservati	on Practices (S	Soils and Wa	ter Conservati	on/	NR)- that prov	ride NR Benefits				
412	Grassed Waterway	short-term	minor to moderate	long-term							
578	Stream Crossing	short-term	minor to moderate	long-term							
600	Terrace	short-term	minor to moderate	long-term							

There could be short-term, minor to moderate adverse impacts to wetlands depending on the location of the conservation practice. Wetlands would be avoided to the greatest extent possible. Any impacts would be localized to the conservation practice area. All conservation practices are intended to conserve and enhance important resources such as wetlands. The practices would have a long-term, beneficial, impact on wetland water quality, hydrology, species composition and vigor. Wetlands impacts could be located on any land use type and the impacts are applicable to both Alternative A and Alternative B.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific conservation practices in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to wetlands:

- In the design of conservation practices the MS TIG would consider resiliency measures related to increasing storm intensities and changing weather patterns (CEQ, 2016).
- Any practice that involves disturbance of wetlands would require authorization by the U.S. Army Corps of Engineers. A Nationwide Permit 27 Aquatic Habitat Restoration, Establishment, and Enhancement Activities would be obtained, with adherence to any permit conditions.
- Develop and implement an erosion control plan to minimize erosion during and after construction and where possible use vegetative buffers (100 feet or greater), revegetate with native species or annual grasses, and conduct work during dry seasons.
- Develop and implement a spill prevention and response plan, including conducting daily inspections of all construction and related equipment to ensure there are no leaks of antifreeze, hydraulic fluid, or other substances and cleaning and sealing all equipment that

- would be used in the water to rid it of chemical residue. Develop a contract stipulation to disallow use of any leaking equipment or vehicles.
- Prohibit use of hazardous materials, such as lead paint, creosote, pentachlorophenol, and other wood preservatives during construction in, over or adjacent to, sensitive sites during construction and routine maintenance.
- Avoid and minimize, to the maximum extent practicable, placement of dredged or fill material in wetlands and other aquatic resources.
- Design construction equipment corridors to avoid and minimize impacts to wetlands and other aquatic resources to the maximum extent practicable.
- To the maximum extent possible, implement the placement of sediment to minimize impacts to existing vegetation or burrowing organisms.
- Apply herbicide in accordance with the direction and guidance provided on the appropriate U.S. Environmental Protection Agency (EPA) labels and state statutes during land-based activities.
- When local conditions indicate the likely presence of contaminated soils and sediments, test soil samples for contaminant levels and take precautions to avoid disturbance of, or provide for proper disposal of, contaminated soils and sediments. Evaluate methods prior to dredging to reduce the potential for impacts from turbidity or tarballs.
- Designate a vehicle staging area removed from any natural surface water resource or wetland
 to perform fueling, maintenance, and storage of construction vehicles and equipment.
 Inspect vehicles and equipment daily prior to leaving the storage area to ensure that no
 petroleum or oil products are leaking.
- Use silt fencing where appropriate to reduce increased turbidity and siltation in the project vicinity. This would apply to both on land and in-water work.

No Action

Under the No Action alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.8). When compared to Alternatives A and B, the No Action alternative would not provide the benefits to hydrology, water quality, or wetlands that would result from the implementation of conservation practice. The No Action alternative does not meet the MS TIG's goals and objectives and clearly does not provide the significant restoration benefit to water quality via nutrient reduction that would occur through the action alternatives.

3.9.1.3 Biological Environment

<u>Introduction to Affected Environment (Biological Environment)</u>: Habitats, Wildlife, and Protected Species are discussed in this section. PDARP Sections 3.4.3.5, and 3.6 are incorporated by reference here. The affected environment for the biological environment for the proposed alternatives is described in respective sections below.

<u>Programmatic Review of Environmental Consequences (Biological Environment)</u>: Sections 6.4.3.2 of the PDARP/PEIS describe the impacts to biological resources for the relevant restoration approaches and are incorporated by reference and briefly described here.

<u>PDARP/PEIS</u> consequences related to biological resources: Depending on the projects implemented, short-term, minor adverse impacts may be anticipated during construction. For example, if

construction includes earth-moving work, terrestrial vegetation may be disturbed. Benefits to biological resources such as benthic invertebrates, shellfish, finfish, and marine mammals could result from 1) improved water quality in the watershed and associated estuary and 2) reduced contaminant loadings (e.g., pesticides and fuel contaminants such as polyaromatic hydrocarbons and metals).

As appropriate in a tiered analysis, the evaluation the proposed alternative focuses on the specific resources with a potential to be affected. Marine and estuarine fauna impacts for the proposed alternatives would be negligible to minor. To avoid redundant or unnecessary information, these resources are evaluated here.

Marine and Estuarine Fauna (Submerged Aquatic Vegetation, Nearshore Benthic Invertebrates, Marine Mammals, Essential Fish Habitat): There would be no in-water marine work or work adjacent to estuarine habitats associated with these proposed alternatives.

For the biological environment, the following resources are further analyzed in this section:

- Habitats and Wildlife
- Protected Species
- Migratory Birds

3.9.1.3.1 Habitats and Wildlife

Affected Environment

The project area for the proposed alternatives is located in the South Atlantic and Gulf Slope Cash Crops, Forest, and Livestock NRCS Land Resource Region (Land Resource Region P). Abundant moisture and a long growing season favor agricultural production in this region. The climate is hot and humid. It is characterized by long, hot summers and short, mild winters. The mean annual precipitation is 44 to 63 inches (1,120 to 1,600 millimeters). The native vegetation consists of oakpine forests. The diverse array of crops includes cotton (*Gossypium* spp.), soybeans (*Glycine max*), peanuts (*Arachis hypogaea*), corn (*Zea mays*), rice (*Oryza sativa*), sugarcane (*Saccharum officinarum*), and wheat (*Triticum aestivum*). The major management concerns on cropland include maintenance of the productivity of the soils, control of erosion, and prevention of groundwater contamination.

The proposed alternatives project area is located in the Southern Coastal Plain Major Land Resource Area (MLRA 133A-1) within the Land Resource Region P. Timber production, cash-grain crops, and forage production are important in this MLRA. Soybeans, cotton, corn, and wheat are the major crops grown throughout the area. Pastures are grazed mainly by beef cattle (*Bos Taurus*), but some dairy cattle and hogs (*Sus scrofa domesticus*) are raised in the area.

The major resource concerns are erosion, maintenance of the content of organic matter and productivity of the soils, control of surface water, artificial drainage, and management of surface compaction and soil moisture. Conservation practices on cropland generally include systems of crop residue management, cover crops, crop rotations, water disposal, subsoiling or deep tillage, pest management, and nutrient management. The most important conservation practice in pastured areas is prescribed grazing (USDA 2016).

The following land use categories (as previously described in Section 3.9) are located in the project area for the proposed alternatives: Associated Agriculture Lands, Crop, Pasture/Grassland, Forest,

Developed Land (Urban), and Water. Conservation practices would be completed predominantly on cropland, pasture/grassland, forestland, and associated agriculture lands.

There are several conservation practices on forestland and riparian habitats. This area supports mixed oak-pine vegetation. Loblolly pine (*Pinus taeda*), longleaf pine (*Pinus palustris*), slash pine (*Pinus elliottii*), shortleaf pine (*Pinus echinata*), sweetgum (*Liquidambar styraciflua*), yellow-poplar (*Liriodendron tulipifera*), red oak (*Quercus rubra*), and white oak (*Quercus alba*) are the major overstory species. Dogwood (*Cornus* spp.), gallberry (*Ilex coriacea*), and farkleberry (*Vaccinium arboretum*) are the major understory species. Common sweetleaf (*Symplocos tinctoria*), American holly (*Ilex opaca*), greenbrier (*Smilax* spp.), southern bayberry (*Myrica cerifera*), little bluestem (*Schizachyrium scoparium*), Elliott bluestem (*Andropogon gyrans*), threeawn (*Aristida purpurea*), grassleaf goldaster (*Pityopsis oligantha*), native lespedezas (*Lespedeza* spp.), and low panicums (*Panicum* spp.) are other understory species.

Some of the major wildlife species in this area are white-tailed deer (*Odocoileus virginianus*), turkey (*Meleagris gallopavo*), rabbit (*Oryctolagus cuniculus*), squirrel (*Sciurus* spp.), bobwhite quail (*Colinus virginianus*), and mourning dove (*Zenaida macroura*). The species of fish in the area include bass (*Micropterus salmoides*), bluegill (*Lepomis macrochirus*), and channel catfish (*Ictalurus punctatus*) (USDA 2016).

Invasive Species EO 13112 applies to all federal agencies whose actions may affect the status of invasive species, requires agencies to identify such actions, and to the extent practicable and permitted by law, requires agencies to 1) take actions specified in the Order to address the problem consistent with their authorities and budgetary resources and 2) not authorize, fund, or carry out actions that they believe are likely to cause or promote the introduction or spread of invasive species in the United States or elsewhere unless pursuant to guidelines that it has prescribed, the agency has determined and made public its determination that the benefits of such actions clearly outweigh the potential harm caused by invasive species; and the benefits of such actions clearly outweigh the potential harm caused by invasive species; and that all feasible and prudent measures to minimize risk of harm will be taken in conjunction with the actions. Best practices that would be used to control or eliminate invasive species are discussed in the environmental consequences section below.

Environmental Consequences for NR Proposed Alternatives A (Preferred) and B

Table 3.9-10 provides a summary of the environmental consequences to habitats and wildlife for representative conservation practices proposed for implementation in the project area for Alternative A (Preferred): Upper Pascagoula Water Quality Enhancement and Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan.

Table 3.9-10: Summary of Impacts to Habitats and Wildlife

Table 3.5 10. Summary of Impacts to Habitats and Whame											
			e A: Upper P Quality Enhar (Preferred)	0			B: Pascagoula F Buffer Maintena				
Practice Code	Conservation Practice Name	Adverse Impact Duration	Adverse Impact Intensity	Beneficial Impact Duration		Adverse Impact Duration	Adverse Impact Intensity	Beneficial Impact Duration			
	Typi	ical Conservatio	n Actions (Eco	logical/NR) that	Pro	vide NR Benefi	ts				
	Streambank and Shoreline		minor to				minor to				
580	Protection	short-term	moderate	long-term		short-term	moderate	long-term			

			ve A: Upper F Quality Enhar (Preferred)	0			e B: Pascagoula I Buffer Maintena	
Practice Code	Conservation Practice Name	Adverse Impact Duration	Beneficial Impact Duration		Adverse Impact Duration	Adverse Impact Intensity	Beneficial Impact Duration	
410	Grade Stabilization Structure	short-term	minor to moderate	long-term		short-term	minor to moderate	long-term
666	Forest Stand Improvement	short-term	minor	long-term		short-term	minor	
	Conservation 1	Practices (Soils	and Water (Conservation/I	NR)	- that provide	NR Benefits	
412	Grassed Waterway	short-term	minor	long-term				
578	Stream Crossing	short-term minor						
600	Terrace	short-term	minor					

Conservation Practices (Ecological/NR)

Streambank and Shoreline Protection (580): There would be short-term, minor to moderate adverse impacts to habitats resulting from grading, reshaping, and planting of stream banks, ponds, lakes, and other aquatic systems. There would be long-term, benefits by revegetating areas with native species. This practice would improve or enhance the stream corridor for fish and wildlife habitat. Areas would be replanted with native vegetation and or seeded to prevent erosion after bank regrading. Erosion control plans would be implemented during and after construction. Impacts would be applicable to Alternative A and Alternative B.

Grade Stabilization Structure (410): There would be short-term, minor to moderate, adverse impacts to habitats from soil excavation, grading, to construct or install grade stabilization structures including berms, rip rap, and hard structures. Most of these grade stabilization structures would be installed in agricultural fields, and could be installed in drainageways or tributaries. There would be long-term, beneficial impacts to aquatic wildlife by stabilizing stream and waterbody habitat and preventing sediment from entering waterways. Areas would be replanted or seeded to prevent erosion after bank regrading. Erosion control plans would be implemented during and after construction. Impacts would be applicable to Alternative A and Alternative B.

Forest Stand Improvement (666): There would be short-term, minor impacts to wildlife and habitat from use of small equipment to access and complete operations which would include use of chainsaws to cut or kill trees or selected understory vegetation, and dragging of felled materials. The use of equipment could damage vegetation and the noise of and activity in the area would cause wildlife to vacate the area during implementation. Wildlife would return after the practice is completed. As a result of this practice, plant health and productivity would improve; invasive species would be removed; and health and vigor of desirable plants would increase. This conservation practice would be designed to have a long-term benefit to habitat and wildlife. Impacts would be applicable to Alternative A and Alternative B.

Conservation Practices (Soil and Water Conservation/NR)

<u>Grassed Waterway (412)</u>: There would be short-term, minor, adverse impacts to habitats and wildlife from noise and activity disturbance during construction. Wildlife would vacate the area during

construction, but return after construction is finished. This practice would be done primarily on cropland and would not impact wildlife habitat. The area would be replanted, were possible with vegetation that would serve to reduce erosion and provide a long-term benefit to wildlife. The grassed waterway practice would be done primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

Stream Crossing (578): There would be short-term, minor impacts to wildlife and habitat from noise and potential vegetation clearing during stream crossing construction. Wildlife would vacate the area during construction, but return after construction is finished. This practice would be done primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

<u>Terrace (600)</u>: There would be short-term, minor, adverse impacts to wildlife and habitat due to potential vegetation clearing and noise disturbance from the use of equipment. Wildlife would vacate the area during construction, but return after construction is finished. The majority of these would be installed in agricultural fields and would not impact wildlife habitat. Areas not in crop production would be replanted or seeded to prevent erosion after bank regrading. Erosion control plans would be implemented during and after construction. This practice would be done primarily on cropland as part of Alternative A; impacts would not be applicable to Alternative B.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific conservation practices in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to habitats, wildlife, and to reduce the spread of invasive species:

- Conservation practices would use natural material in any conservation practice that advises
 the use of materials and native plantings and seedlings, as well as natural revegetation. The
 footprint of any disturbance would be minimized the extent practicable. Clearing activities
 would be discouraged in forested wetlands.
- All equipment to be used during the project, including personal gear, would be inspected and cleaned such that there is no observable presence of mud, seeds, vegetation, insects and other species.

No Action

Under the No Action alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.8). When compared to Alternatives A and B, the No Action alternative would not provide the benefits to habitats and wildlife that would be provided by the implementation of various conservation practices. The No Action alternative does not meet the MS TIG's goals and objectives and clearly does not provide the significant restoration benefit to water quality via nutrient reduction that would occur through the action alternatives.

3.9.1.3.2 Protected Species

Affected Environment

The U.S. Fish and Wildlife Service (USFWS) and NOAA National Marine Fisheries Services (NMFS) designates (lists) species as threatened or endangered when they meet criteria detailed under the Endangered Species Act (ESA) of 1973, as amended (16 U.S.C. §1531 et seq.). Additionally, Mississippi Wildlife Fisheries and Parks (MWFP) identifies and listsspecies for protection. Section 7(a) (2) of the ESA requires that each federal agency ensure that any action authorized, funded, or carried out by the agency is not likely to jeopardize the continued existence of any endangered or threatened species or result in the destruction or adverse modification of Critical Habitat of those species. When the action of a federal agency, either by activity, permitting, or funding, may affect a protected species or its Critical Habitat, that agency is required to consult with either the NMFS or the USFWS, depending on which agency has jurisdiction over the protected species that may be affected. The USDA-NRCS has already completed a programmatic ESA consultation with the USFWS confirming NRCS experience that the conservation practices likely to be implemented under the proposed action alternative may affect but are not likely to adversely affect protected species. Because USDA-NRCS conservation practices would be used under both these alternatives, the MS TIG is confirming that the USFWS agree the NR projects proposed in this RP/EA similarly may affect but would not be likely to adversely affect protected species in the project area. Appropriate recommendations would be incorporated into the proposed project alternatives to the extent feasible. Compliance with the Migratory Bird Treaty Act and Bald and Golden Eagle Protection Act are also discussed in this section.

Federally protected species that are known to occur or could occur in Newton, Lauderdale, Clarke, Neshoba, and Kemper counties are listed in Table 3.9-39.

Table 3.9-11: Federally threatened, endangered, and proposed species

Common Name	Scientific Name	Federal Status	County	Habitat				
Birds								
Red-cockaded Woodpecker	Picoides borealis	Endangered	Newton	This species excavates nesting and roosting cavities in living pine trees, and is the only species known to do so exclusively. Cavities have been found in most species of southern pines, but longleaf pine appears to be the preferred species. Older, mature trees are selected for cavity excavation.				
Wood Stork	ood Stork Mycteria Americana		All	Freshwater and estuarine wetlands, primarily nesting in cypress or mangrove swamps. They feed in freshwater marshes, narrow tidal creeks, or flooded tidal pools. Particularly attractive feeding sites are depressions in marshes or swamps where fish become concentrated during periods of falling water levels.				
Fish	•							
Gulf Sturgeon	Acipenser oxyrinchus desotoi	Threatened	Clarke	Migrates from large freshwater coastal rivers to brackish and marine coastal bays, estuaries and the Gulf of Mexico.				
Mammals								
Northern long- eared bat	Myotis septentrionalis	Threatened	Kemper, Lauderdale, Neshoba, Newton	During the summer, northern long-eared bats roost singly or in colonies underneath bark, in cavities or in crevices of both live trees and snags (dead trees). Northern long-eared bats spend winter hibernating in caves and mines, called hibernacula.				

Common Name	Scientific Name	Federal Status	County	Habitat			
Reptiles							
Ringed map Turtle, Ringed Sawback	Graptemys oculifera	Threatened	Neshoba	The threatened ringed map turtle is found in the Pearl River. It prefers river stretches with moderate currents, abundant basking sites, and sand bars for nesting.			
Gopher tortoise	Gopherus Polyphemus	Threatened	Clarke	Well-drained, sandy soils, which allow easy burrowing; an abundance of diverse herbaceous ground cover; and an open canopy and sparse shrub cover, which allows sunlight to reach the ground floor (USFWS 2013).			
Yellow- blotched map turtle, Yellow- blotched sawback	Graptemys flavimaculata	Threatened	Clarke	Habitat is streams with strong, consistent current and large sandbars for nesting.			
Plants							
Price's potato bean	Apios priceana	Threatened	Kemper	This species found on slopes or bluffs with open woods that often grade into creek and river bottoms. The species may also be found along forested margins of power-line and road rights-of-ways.			

Birds

Wood Stork (*Mycteria americana Linnaeus*): In Mississippi, wood storks have been observed most frequently along the western edge of the state in those counties bordering the Mississippi River and with increasing frequency in some counties along the eastern edge of the state, although they may occur almost anywhere there are sloughs or swamps to provide feeding habitat. The wood stork occurs primarily in freshwater wetlands, including ponds, bayheads, flooded pastures, oxbow lakes, and ditches. Nesting usually occurs in bald cypress trees in swamps, although breeding has also been observed in mangroves (MS Museum of Natural Science 2014). Therefore, the proposed alternative may affect the wood stork and we will consult with the U.S. Fish and Wildlife Service will be consulted as appropriate. However, strict adherence to the USDA-NRCS conservation practices that have been consulted on previously may lead to a No Effect Determination by USDA-NRCS and would then not require further ESA consultation.

Red-cockaded Woodpecker (*Picoides borealis*): In Mississippi, this species has been recorded primarily from the southern two-thirds of the state. It has not been found in the Delta and only sporadically occurs in the northern counties. The Red-cockaded woodpecker is a species of southern pine forests. The preferred nesting habitat is open, park-like, mature pine woodlands with few or no hardwood trees present. Preferred feeding habitats are pine stands with trees 23 cm (9 in.) and greater in diameter. These may or may not include a significant hardwood component. The Red-cockaded woodpecker excavates nesting and roosting cavities in living pine trees, and is the only species known to do so exclusively. Cavities have been found in most species of southern pines, but longleaf pine (*Pinus palustris*) appears to be the preferred species. Older, mature trees are selected for cavity excavation. (MS Museum of Natural Science 2014). Therefore, the proposed alternative may affect the Red-cockaded Woodpecker and we will consult with the U.S. Fish and Wildlife Service will be consulted as appropriate. However, strict adherence to the USDA-NRCS conservation practices that have been consulted on previously may lead to a No Effect Determination by USDA-NRCS and would then not require further ESA consultation.

Fishes

Gulf Sturgeon (*Acipenser oxyrinchus desotoi*): In the Pascagoula River watershed, the Gulf sturgeon occurs in the Chickasawhay River upstream to at least the town of Waynesboro (MS Museum of Natural Science 2014). Waynesboro is approximately 34 miles south of the southern extent of the proposed alternatives. Therefore, the proposed alternative is expected to have No Effect on Gulf sturgeon and consultation with the U.S. Fish and Wildlife Service will not be requested.

Mammals

Northern Long-eared Bat (*Myotis septentrionalis*): Northern long-eared bats typically hibernate in caves and enter hibernation sometime between September and November. They emerge during the spring between March and May depending on latitude. The species typically does not hibernate as a single species, but with large numbers of other bats of varying species. The species frequents forest interiors and consumes a diet consisting predominantly of moths, beetles, and flies. They forage both under forest canopy and along forest edges primarily during the first two hours after sunset. Mating occurs between July and October, with births taking place between May and July (MSU 2016).

Reptiles

Gopher Tortoise (*Gopherus polyphemus*): The gopher tortoise uses well-drained to excessively well-drained upland soils. Tortoises require soils that are sandy enough to permit construction of burrows and open canopies that allow sufficient herbaceous plant growth and sunny areas in which to nest. In Mississippi, these areas often support a mixture of longleaf pine and scrub oaks.

Ringed Map Turtle (*Graptemys oculifera*): This turtle occurs only in the Pearl River and its tributary, the Bogue Chitto River (MS Museum of Natural Science 2014).

Yellow-blotched Map Turtle (*Graptemys flavimaculata*): A Mississippi endemic, the yellow-blotched map turtle occurs in the Pascagoula, Chickasawhay, Leaf, Bouie, and Escatawpa rivers and in Tallahala, Black, Bluff, Bogue Homa, Bucatunna, Gaines, Okatoma, and Thompson's creeks. This turtle occurs in the Pascagoula River from Jackson County upriver to the confluence of the Leaf and Chickasawhay rivers in George County. It is sporadically distributed up the Leaf River to Covington County and as far upstream as Clarke County in the Chickasawhay River. The largest and most viable population appears to occur in the lower Pascagoula River from the town of Wade downstream to the beginning of the brackish marshes at the mouth of the Pascagoula River. The yellow-blotched map turtle requires streams with strong, consistent current and large sandbars for nesting. It spends much of the day basking, so it needs streams which are wide enough to receive several hours of direct sunlight per day and which have abundant snags and logs on which to bask. This habitat type is most often found in the rivers and larger creeks within its range, but may also be found in bends of medium-sized (15 -30 m wide) creeks. (MS Museum of Natural Science 2014).

Plants

Price's Potato Bean (*Apios priceana*): In Mississippi, populations have been found in Oktibbeha, Lee, and Kemper counties. Historically, this species has been found in Clay County, and new populations may still be found there, as well as in Chickasaw, Pontotoc and Benton counties. Populations occur in open woods and along woodland edges in limestone areas, often where bluffs grade into creek or river bottoms. Several populations extend onto roadside or powerline rights-of-way. The soils are described as well-drained loams on old alluvium or over limestone. Plant associates in Mississippi's populations include chinkapin oak, white ash, basswood, sugar maple, slippery elm, redbud, spicebush, and switchcane. This species is thought to be a native of forest

openings and thrives best in areas with partial canopy. Price's potato bean flowers from late June through July and produces fruit in August.

Environmental Consequences for NR Proposed Alternatives A (Preferred) and B

PDARP programmatic ESA consultations were developed with the National Marine Fisheries Services (NMFS, 2016) and the U.S. Fish and Wildlife Service (USFWS, 2016). Potential impacts to threatened or endangered species and their Critical Habitat are presented in Table 3.9-12. The MS TIG has begun coordination pursuant to ESA under the programmatic ESA consultations. The project area for the proposed alternatives is not within Gulf sturgeon Critical Habitat.

Table 3.9-12: Protected Species Impacts

Species /Critical Habitat	Applicable Habitats	Example Conservation Practices for Applicable Habitats	Potential Impacts to Species/Critical Habitat
Red-cockaded woodpecker (Picoides borealis)	Forest	 Forest Stand Improvement (666) Streambank and Shoreline Protection (580) 	This species may use this habitat for foraging, loafing, and nesting. During project planning, surveys would be done if species occurrence is expected. Activities would be planned so as to avoid disturbing the species or its habitat. Chapter 6 of the PDARP outlines Best Practices for this species. The Best Practices are discussed in this Section.
Wood stork (Mycteria americana)	Forest	 Forest Stand Improvement (666) Streambank and Shoreline Protection (580) 	Wood stork may use this habitat for foraging and loafing. The species does not nest in the project area for proposed alternatives. The species would be able to vacate the area during conservation practice implementation, and return after completion.
Gulf sturgeon (Acipenser oxyrinchus desotoi)	Not located in project area for proposed alternatives	n/a	n/a
Northern long-eared bat (Myotis septentrionalis)	Forest	 Forest Stand Improvement (666) Streambank and Shoreline Protection (580) 	This species may exist in forested areas where there are snags or under exfoliating bark, cracks, or crevices in trees. If habitat exists in the area, either surveys/avoidance or both would be completed during the design of the conservation practice. Best management practices outlined in the 4(d) rule would be implemented.
Ringed map turtle (Graptemys oculifera)	Not located in project area for proposed alternatives	• n/a	n/a
Gopher tortoise (Gopherus polyphemus)	Habitat will not likely be present in Conservation Action areas Forest Grassland	Forest Stand Improvement (666)	If suitable habitat is present at the location of a selected conservation practice, surveys would be conducted. Gopher tortoise burrows would be avoided. Best Practices from Chapter 6 of the PDARP are discussed later in the this Section.
Yellow-blotched map turtle (Graptemys	Water	• Stream Crossing (578)	Conservation practices could result in a noise impact and habitat disturbance

Species /Critical Habitat	Applicable Habitats	Example Conservation Practices for Applicable Habitats	Potential Impacts to Species/Critical Habitat
flavimaculata)		• Streambank and Shoreline Protection (580)	causing the species to temporarily vacate the area. If potential habitat is found, the conservation practice would be designed so as to minimize the exposure to the species.
Price's potato bean (Apios priceana)	ForestGrassland	 Forest Stand Improvement (666) Grassed Waterway (412) 	Conservation practices could result in an impact to vegetation. Prior to conservation practice implementation the USDA-NRCS would coordinate with the USFWS Ecological Services Field Office to determine if a survey is needed. If a survey reveals occurrence of this species in the area, then it would be avoided. If avoidance was not possible, vegetation removal would be done by hand, so as to avoid individuals or colonies.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific conservation practices in different locations due to differences in relevant conditions. The MS TIG would continue to consult with the appropriate regulatory agency to further avoid or minimize take of these species in the planning site-specific conservation practices. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to protected species:

- Northern long-eared bat
 - O The USFWS listed the northern long-eared bat as threatened under the Endangered Species Act on May 4, 2015, and established an interim 4(d) rule to help protect the species. The Endangered Species Act 4(d) rule for the northern long-eared bat best practices are outlined below:
 - No purposeful take of northern long-eared bats within the bat's range with the exception of:
 - removal from human structures,
 - defense of human life (public health monitoring for rabies), and
 - removal of hazardous trees for protection of human
 - life and property.
 - o Additionally, no incidental take of northern long-eared bats are allowed:
 - within a hibernacula,
 - if it results from tree removal activities within 0.25 mile
 - of a known hibernacula, and
 - if it results from tree removal activities that cut or destroy a known, occupied maternity roost tree or other trees within 150 feet of a known, occupied maternity roost tree during June and July (MSU, 2016).

• Red-Cockaded Woodpecker

- Avoid working within active red-cockaded woodpecker clusters (the minimum convex polygon containing the aggregation of cavity trees used by a group of red-cockaded woodpeckers and a 200- foot-wide buffer surrounding the polygon).
- o If avoidance is not possible or management activities in red-cockaded woodpecker suitable habitat are desired, conduct standard surveys to determine if the habitat is supporting any individuals or presence can be assumed. If red-cockaded woodpeckers are present (or assumed to be), avoid cavity trees and use mechanized equipment during the non-nesting season (approximately April 1 through July 31).
- o If tree removal is necessary, survey pine trees approximately 60 or more years old for active cavities within one year of the proposed removal. Extend surveys from the project site out to no less than one- half mile. Replace any cavities affected by the project via drilled cavity construction.
- o If impacts to suitable foraging habitat (pines approximately 30 or more years old and within one-half mile of an active cavity tree) are proposed, conduct a foraging habitat analysis. Foraging habitat may need to be replanted post-project.
- Design projects within red-cockaded woodpecker suitable habitat such that prescribed fire needs are not impeded.

• Gopher Tortoise

o If suitable habitat is present, coordinate with the USFWS Ecological Services Field Office in Jackson, MS to discuss the need for surveys to identify any gopher tortoise burrows and to develop conservation measures to avoid or minimize impacts. Measures could include establishing a protective buffer (size determined by USFWS and the state trust resource agency) if burrows are within the project area and cannot be avoided, implementing standard procedures to relocate the tortoise within the project site but away from the areas of construction or restoration or considering conservation banks. A Candidate Conservation Agreement with Assurances may be appropriate for project sites within the non-listed range of the species.

Protected Plants

- o If suitable habitat is present, coordinate with the USFWS Ecological Services Field Office in Jackson, MS to discuss the need for surveys to identify protected plants and to develop conservation measures to avoid or minimize impacts.
- Enhance and protect plants on site and in adjacent habitats to the maximum extent possible.

Protected Species (Wood stork, Northern long-eared bat)

o Provide all individuals working on a project with information in support of general awareness of and means to avoid impacts to protected species and their habitats present at the specific project site. If suitable habitat is present, coordinate with the USFWS Ecological Services Field Office in Jackson, MS to discuss the need for surveys.

No Action

Under the No Action alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.8). When compared to Alternatives A and B the no action would not enhance habitat that protected species could utilize. The No Action alternative does not meet the MS TIG's goals and objectives and clearly does not provide the significant restoration benefit to water quality via nutrient reduction that would occur through the action alternatives.

3.9.1.3.3 Migratory Birds

Affected Environment

Migratory bird species groups that could occur in the project area for the proposed alternatives include wading birds, raptors, goatsuckers, waterfowl, doves and pigeons, and rails and coots (see Table 3.9-13).

Table 3.9-13: Species Groups Present in Project Area for the Proposed Alternatives

SPECIES	BEHAVIOR	SPECIES/HABITAT IMPACTS
Wading birds (herons, egrets, ibises)	Foraging, feeding, resting, roosting, nesting	Wading birds primarily forage and feed at the water's edge. There would be limited habitat in the project area for the proposed alternatives except for ponds and potential habitat that could occur in streams. As such, they may be impacted locally and temporarily by the project alternatives. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. These birds primarily nest and roost in trees or shrubs (e.g. pines, <i>Bacchurus</i>). Nesting sites if located would be avoided during construction.
Raptors (osprey, hawks, eagles, owls)	Foraging, feeding, resting, roosting, nesting	Raptors forage, feed, rest and nest in the action area. As such, they may be impacted locally and temporarily by the project alternatives. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. Operation of large equipment could disturb birds. Care would be take to identify and avoid raptor nests during the construction/installation of conservation practices.
Goatsuckers	Foraging, feeding, resting, roosting, nesting	Goatsuckers forage, feed, rest, and roost in the action area. However, they are nocturnal/crepuscular and therefore not active during the project work period. They nest in thickets and woodlands. Prior to doing work in woodlands, nesting surveys would be completed or construction would be avoided during nesting season.
Waterfowl (ducks, loons, and grebes)	Foraging, feeding, resting, roosting, nesting	Waterfowl forage, feed, rest, and roost in the action area. As such, they may be impacted locally and temporarily by the project alternatives. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting. These birds primarily roost and nest in low vegetation. There would be limited if any habitat suitable for nesting waterfowl. To the extent nesting waterfowl are encountered in the design of conservation practice, it would be avoided.
Doves and pigeons	Foraging, feeding, resting, roosting	Doves and pigeons could forage, feed, rest, and roost in the action area. As such, they may be impacted locally and temporarily by the project alternatives. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting.

SPECIES	BEHAVIOR	SPECIES/HABITAT IMPACTS
Rails and coots	Foraging, feeding, resting, roosting, nesting	Rails and coots forage, feed, rest, and roost in the action area. As such, they may be impacted locally and temporarily by the project alternatives. It is expected that they would be able to move to another nearby location to continue foraging, feeding and resting if disturbed by the project alternatives. These birds primarily roost and nest in marshes, which are within the action area, and adjacent to project alternative activities which are in-water. There would be limited habitat in the areas where conservation practices are installed. If habitat is present work a survey for presence would be completed and areas would be avoided during nesting season.

Migratory Bird Treaty Act: The Migratory Bird Treaty Act of 1918 (MBTA) implements various treaties and conventions among the United States, Canada, Japan, Mexico, and the former Soviet Union for the protection of migratory birds. Under MBTA, unless permitted by regulations, it is unlawful to pursue; hunt; take; capture or kill; attempt to take, capture, or kill; possess; offer to sell or sell; barter; purchase; deliver; or cause to be shipped, exported, imported, transported, carried, or received any migratory bird, part, nest, egg, or product, manufactured or not. USFWS regulations broadly define "take" under MBTA to mean "pursue, hunt, shoot, wound, kill, trap, capture, or collect, or attempt to pursue, hunt, shoot, wound, kill, trap, capture, or collect".

Bald and Golden Eagle Protection Act: The Bald and Golden Eagle Protection Act (BGEPA) of 1940 (16 U.S.C. 668-668c) prohibits anyone, without a permit issued by the Secretary of the Interior, from "taking" bald eagles, including their parts, nests, or eggs. BGEPA provides criminal penalties for persons who "take, possess, sell, purchase, barter, offer to sell, purchase or barter, transport, export or import, at any time or any manner, any bald eagle ... [or any golden eagle], alive or dead, or any part, nest, or egg thereof." Golden eagles are not present in the project area for the proposed alternatives.

Environmental Consequences NR Proposed Alternatives A (Preferred) and B

Migratory birds could use areas at and around the project alternative(s) project area for foraging, feeding, resting, and nesting. Nesting species include raptors (forest edge near wet areas), wading birds (pine trees/shrubs adjacent to wet areas), and waterfowl (open water); Table 3.9-8. For all planned restoration activities, pre-commencement nesting surveys for migratory birds and raptors within the restoration activity area would be conducted and if evidence of nesting is found, the USDA-NRCS would coordinate with the USFWS to develop and implement appropriate conservation measures, such as those described below due to the implementation of best management practices no "take" of nesting birds is anticipated. There are no golden eagles in the project footprint for the proposed alternatives. Raptor nest surveys would be completed within the restoration activity area where raptor nesting habitat exists. If evidence of nesting is found, USDA-NRCS would coordinate with the USFWS to develop and implement appropriate conservation measures, therefore no impacts to golden or bald eagles are anticipated. Potential adverse effects to birds include elevated noise levels due to the use of mechanical equipment for vegetation clearing, and from noise and smoke during prescribed burning. These species are mobile and would likely exit the area during management activities (no impacts to overall population). Foraging and resting birds may temporarily be displaced during management activities. Bird roosting would not be affected because management activities would occur during daylight hours. Therefore, impacts are expected to be short-term, localized, and minor.

Best Practices

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific conservation practices in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to migratory birds including bald eagles:

Migratory Birds

- Use care to avoid birds when operating machinery or vehicles near birds.
- Avoid working in migratory bird nesting habitats during breeding, nesting, and fledging
 (approximately mid-February through late August). If project activities must occur during
 this timeframe and breeding, nesting, or fledging birds are present, contact the state trust
 resource agency to obtain the most recent guidance to protect nesting birds or rookeries, and
 their recommendations would be implemented.
- Conservation areas may already be marked to protect bird nesting areas. Stay out of existing marked areas.
- If vegetation clearing is necessary, clear vegetation outside the migratory bird nesting season (approximately mid-February through late August) or have a qualified biologist inspect for active nests. If no active nests are found, vegetation may be removed. If active nests are found, vegetation may be removed after the nest successfully fledges.

Bald Eagles

- If bald eagle breeding or nesting behaviors are observed or a nest is discovered or known, have all activities avoid the nest by a minimum of 660 feet. If the nest is protected by a vegetated buffer where there is *no* line of sight to the nest, then the minimum avoidance distance is 330 feet. Maintain this avoidance distance from the onset of breeding/courtship behaviors until any eggs have hatched and eaglets have fledged (approximately 6 months).
- If a similar activity (such as driving on a roadway) is closer than 660 feet to a nest, maintain a distance buffer as close to the nest as the existing tolerated activity. If a vegetated buffer is present and there is no line of sight to the nest and a similar activity is closer than 330 feet to a nest, then maintain a distance buffer as close to the nest as the existing tolerated activity.
- In some instances, activities conducted within 660 feet of a nest may result in disturbance, particularly for the eagles occupying the Mississippi barrier islands. If an activity appears to cause initial disturbance, stop the activity and move all individuals and equipment away until the eagles are no longer displaying disturbance behaviors. Contact the USFWS's Migratory Bird Permit Office to determine how to avoid impacts or if a permit may be needed.

The MS TIG has begun coordination and review of the project alternatives for impacts to bald eagles and migratory birds in accordance with the Bald and Golden Eagle Protection Act (BGEPA) of 1940 (16 U.S.C. 668-668c) and the Migratory Bird Treaty Act (MBTA) of 1918 (16 U.S.C. 703–712) to ensure appropriate conservation practices would be incorporated into the selected project alternative.

No Action

Under the No Action alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes

to occur (outcomes described in Section 3.8). There would be no adverse impacts to migratory birds or bald and golden eagles under the No Action alternative. The No Action alternative does not meet the MS TIG's goals and objectives and clearly does not provide the significant restoration benefit to water quality via nutrient reduction that would occur through the action alternatives.

3.9.1.4 Socioeconomic Resources

<u>Introduction to Affected Environment (Socioeconomic Environment)</u>: Socioeconomic and environmental justice, cultural resources and public health and safety are discussed in this section. PDARP Section 3.2 is incorporated by reference here. The affected environment for the physical environment for the proposed alternatives is described in respective sections below.

<u>Programmatic Review of Environmental Consequences (Socioeconomic Resources)</u>: Sections 6.4.3.1.3 of the PDARP/PEIS describe the impacts to Socioeconomic Resources for the relevant restoration approaches and are incorporated by reference and briefly described here.

<u>PDARP/PEIS</u> consequences related to economic effects: Impacts to socioeconomics resulting from the implementation of this restoration approach are dependent on site-specific conditions associated with a project proposed for implementation. Depending on the techniques employed, short-term benefits to the local economy could accrue through an increase in employment and associated spending in the project area during construction activities.

<u>PDARP/PEIS</u> consequences related to cultural resources: If cultural or historic resources are present, minor adverse impacts to the resource would be anticipated during construction activities.

As appropriate in a tiered analysis, the evaluation the proposed alternative focuses on the specific resources with a potential to be affected. Infrastructure, land and marine management, tourism and recreation, fisheries and aquaculture, fisheries and aquaculture, marine transportation and aesthetic and visual resources impacts for the proposed alternatives would be negligible to minor. To avoid redundant or unnecessary information, these resources are evaluated here.

Infrastructure: No publicly owned or maintained infrastructure would be created or impacted as a result of these proposed alternatives.

Land and Marine Management: The end result of these proposed alternatives would be implementation of conservation practice planning and implementation under the guidance and oversight of USDA-NRCS on cropland, associated agriculture lands, pasture/grassland, forestland and riparian areas. The conservation practices are consistent with current farmstead uses and operation that would otherwise would not have benefit of conservation planning and oversight. This would constitute a benefit to landuse for landowners who voluntarily participate in the program. There would be no adverse impacts to land management.

Tourism and Recreational Use: The proposed alternatives would be carried out by the voluntary application of practices by land owners on their own land. Private land is not subject to tourism and any recreational benefits associated with the implementation of conservation practices (e.g. wildlife habitat, stream stabilization), would primarily benefit participants. Implementation of the selected proposed alternative would have negligible impacts, if any, on tourism and recreational use.

Fisheries and Aquaculture: Implementation of the selected proposed alternative could include streambank stabilization in ephemeral and intermittent tributaries. Monitoring would include in-water work near the site of implemented conservation practices. Implementation of the selected proposed alternative would not affect a commercial fishery or aquaculture operation.

Marine Transportation: There would be no marine in-water work associated with the selected proposed alternative.

Aesthetic and Visual Resources: Conservation practices would be implemented on cropland, associated agricultural lands, pastureland/hayland, and forestland. Conservation practices would not be inconsistent with current farming practices and would have a negligible effect on aesthetic and visual resources.

For the socioeconomic and environmental justice, the following resources are further analyzed in this section:

- Socioeconomics and Environmental Justice
- Cultural resources
- Public Health and Safety

3.9.1.4.1 Socioeconomics and Environmental Justice

Affected Environment

The affected environment for the proposed alternative includes portions of the population of portions of Newton, Lauderdale, Clarke, Neshoba, and Kemper counties (Table 3.9-14). From 2009-2013, median household income in Clarke County was \$31,362, which was 20% lower than the median household income in the State of Mississippi (\$39,464); the median household income in Kemper County was \$29,003, 27% lower than the median household income in the State of Mississippi; the median household income in Lauderdale County was \$36,203, 8% lower than the median household income in the State of Mississippi; the median household income in Neshoba County was \$37,050, 6% lower than the median household income in the State of Mississippi; the median household income in Newton County was \$39,190, <1% lower than the median household income in the State of Mississippi. (U.S. Census Bureau, 2010-2014 American Community Survey 5-Year Estimates)

Table 3.9-14: - Population data (http://www.census.gov/2010census/popmap/)

<u> </u>	oic 515 111 1 opulation data				ittp://www.teensusigov/2010census/poplitup/)									
Topic	Mississippi		ssissippi Clarke County Kemper County		Lauderdale County		Neshoba County		Newton County					
2010 Total Population	2,967,297		16,732		10,456		80,261		29,9676		21,720			
White alone	1,754,684	59%	10,741	64%	3,689	35%	43,957	55%	17,974	61%	13,734	63%		
Black or African American alone	1,098,385	37%	5,759	34%	6,288	60%	34,330	43%	6,207	21%	6,567	30%		
Asian alone	25,742	<1%	29	<1%	10	<1%	580	1%	102	<1%	52	<1%		
American Indian and Alaska Native alone	15,030	<1%	60	<1%	385	4%	178	<1%	4,815	16%	1,092	5%		

Торіс	Mississippi		Clarke County		Kemper County		Lauderdale County		Neshoba County		Newton County	
Native Hawaiian and Other Pacific Islander alone	1,187	<1%	1	<1%	0	0%	30	<1%	9	<1%	0	0%
Some Other Race alone	38,162	1.3%	43	<1%	8	<1%	520	1%	140	<1%	86	<1%
Two or More Races	34,107	1.1%	99	1%	76	1%	666	1%	429	1%	189	1%

Environmental Consequences NR Proposed Alternatives A (Preferred) and B

There would be long-term beneficial socioeconomic benefits to landowners who voluntarily participate in the program from the development of a conservation plan; program investment in the cropland, pasture/grassland, associated agriculture lands, forestland and/or riparian areas; benefits from costs saving practices that result from reduction in erosion/costs for maintaining eroded drainage ways, cost reduction resulting from nutrient management, improved production/yield from crops from the implementation of soil and water conservation practices, and increases in the farmstead value because of the capital investment in farmstead improvements. There would be no adverse impacts to socioeconomics from the implementation of proposed Alternative A or B. There would be no disproportionate impacts to low-income or minority populations as a result of either of the project alternatives.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.8). The No Action alternative would have no widespread impact or benefit to socioeconomic resources. Private landowners would not benefit from additional funds to improve their land. The No Action alternative does not meet the MS TIG's goals and objectives and clearly does not provide the significant restoration benefit to water quality via nutrient reduction that would occur through the action alternatives.

3.9.1.4.2 Cultural Resources

Affected Environment

Cultural resources include historic properties listed in, or eligible for listing in the National Register of Historic Places (36 C.F.R. §60[a-d]). The National Historic Preservation Act of 1966 (NHPA), as amended and recodified (54 U.S.C. § 300308), defines an historic property as "any prehistoric or historic district, site, building, structure, or object included on, or eligible for inclusion on the National Register [of Historic Places]." Under the statute and implementing regulations, historic properties include significant traditional religious and cultural properties important to Indian tribes. Historic properties include built resources (bridges, buildings, piers, etc.), archaeological sites, and Traditional Cultural Properties, which are significant for their association with practices or beliefs of a living community that are both fundamental to that community's history and a piece of the community's cultural identity. Although often associated with Native American traditions, such properties also may be important for their significance to ethnic groups or communities. Historic properties also include submerged resources.

This proposed project alternative(s) are currently being reviewed under Section 106 of the NHPA to identify any historic properties located within the proposed alternative(s) project area and to evaluate whether the alternatives would affect any historic properties. The MS TIG is currently conducting a literature review of the proposed project alternatives. Previously recorded archaeological sites, historical standing structures, historic districts, historic churches, post offices, utilities and other resources that are potential National Register of Historic Places (NRHP) properties, National Register Districts and National Historic Landmarks are being reviewed. The preliminary review of the previously recorded archaeological sites using MDAH records revealed archaeological sites located within the vicinity of the proposed project alternatives.

Environmental Consequences for NR Proposed Alternatives A (Preferred) and B

The NHPA charges the federal government with protecting the cultural heritage and resources of the nation. A complete review of the project alternative(s) under Section 106 of the NHPA would be completed as environmental assessment continues. This selected project alternative would be implemented in accordance with all applicable laws and regulations concerning the protection of cultural and historic resources. For site-specific conservation practices, potential effects to historic properties would be considered when the undertaking is the type of activity that has the potential to cause effects on these resources. Resources that are eligible for the National Register of Historic Places would be avoided in the design of the conservation practices, to the extent practicable.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.8). There would be no adverse impact to cultural resources under the No Action. The No Action alternative does not meet the MS TIG's goals and objectives and clearly does not provide the significant restoration benefit to water quality via nutrient reduction that would occur through the action alternatives.

3.9.1.4.3 Public Health and Safety

Affected Resources

The majority of the conservation practices would occur on associated agriculture lands, cropland, pasture/grassland, forestland or in riparian areas or streams. Conservation practices in floodplains could have FEMA floodplain mapping and engineering requirements for installation of conservation practices.

Environmental Consequences for NR Proposed Alternatives A (Preferred) and B

There would be no adverse impact to public health and safety. The program is voluntary and would be completed on private land under the guidance of the USDA-NRCS. There would be beneficial impacts to water quality in the watershed. The conservation practices for the proposed alternatives would be utilized for NR purposes which would enhance water quality. Improved water quality is beneficial to public health since the waters in the watershed are mostly classified as "recreation" and "fish and wildlife" by the State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters (MDEQ 2012). Conservation practices would be designed so as not result in a detectable change to natural and beneficial floodplain values. There would be limited fill, if any, in the floodplain. Restored hydrology streambank stabilization and grade control structures would enhance floodplain functions.

No Action Alternative

Under the No Action alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes to occur (outcomes described in Section 3.8). The No Action would not result in any additional health and safety benefits or impacts. The No Action alternative does not meet the MS TIG's goals and objectives and clearly does not provide the significant restoration benefit to water quality via nutrient reduction that would occur through the action alternatives.

3.9.2 Site-specific NEPA Review for NR Proposed Alternatives A (Preferred) and B

Section 3.9.1 is a discussion of environmental consequences analysis of proposed Alternative A (Preferred) and B for the NR Restoration Type at a programmatic level. The exact parcels and associated conservation practices on those parcels are not known at this time. The environmental consequences are based on the range of restoration conservation practices contemplated on parcels for proposed alternative project areas. The programmatic analysis provides maximum impacts to each of the resource categories based on the MS TIG's knowledge of the proposed alternative project area. The MS TIG is proposing the selection of Alternative A (Preferred). Section 3.7.2 also presents a process that the MS TIG would follow to complete the requirements of NEPA and other environmental statutes as site-specific conservation practices are planned for Alternative A, if selected.

3.10 Cumulative Impacts for NR

Section 6.6 and Appendix 6B of the PDARP/PEIS are incorporated by reference into the following cumulative impacts analysis including the methodologies for assessing cumulative impacts, identification of affected resources and the cumulative impacts scenario. A development of the analysis in the context of the affected environment of the proposed NR alternatives (X), when added to the impacts from applicable past, present and reasonably foreseeable future actions (Y), to understand the potential cumulative impacts to an affected resource (Z), or where the effects may interact and/or be additive, that is X + Y = Z. This analysis includes the alternatives proposed for the NR Restoration Type in this Draft RP/EA, which include:

Alternative A (Preferred): Upper Pascagoula River Water Quality Enhancement Alternative B: Pascagoula River Basin Riparian Buffer Maintenance Plan

3.10.1 Identification of Resources Affected

Section 3.9 provides an environmental consequences analysis for the following resources that would have minor to negligible effects, and based on their magnitude, with respect to context and intensity, would not contribute to cumulative impacts. These resources are excluded from this cumulative impacts analysis:

- Air Quality and Greenhouse Gas Emissions
- Noise
- Invasive Species
- Marine and Estuarine Fauna

- Infrastructure
- Tourism and Recreation
- Fisheries and Aquaculture
- Marine Transportation
- Land and Marine Management
- Aesthetics and Visual Resources

In the planning of site-specific conservation plans, the Implementing Trustee would avoid and minimize impacts to protected species, cultural resources, and migratory birds; these resources would not contribute to cumulative impacts. Of the resources listed above, most were determined to have impacts that would not contribute to cumulative impacts, based on their magnitude with respect to context and intensity, and are therefore excluded from this cumulative impacts analysis.

The following resources were analyzed in detail for environmental consequences that could result from implementation of the alternative.

- Geology and Substrates
- Hydrology and Water Quality
- Habitats and Wildlife

3.10.2 Cumulative Action Scenario

In order to effectively consider the potential cumulative impacts, the MS TIG identified local and site-specific past, current and reasonably foreseeable future actions which are considered relevant to identifying any cumulative impacts the alternatives may have on a local scale. These actions fall within the established spatial and temporal boundaries. For the purpose of this cumulative impacts analysis the spatial extent will be the same as the project location which includes portions of Newton, Lauderdale, Clarke, Neshoba, and Kemper counties, Mississippi (Figure 3.9.1). The cumulative impacts analysis depends on the availability of information and data about past, present and reasonably foreseeable future actions. For this Draft RP/EA, the MS TIG identified USDA conservation program-funded conservation practices that had been completed in the recent past and are foreseeable and are summarized in Table 3.10-1. The cumulative effects for both Alternatives A and B will be the same with the exception that Alternative A will result in a higher level of treatment on fewer locations than Alternative B, which is more likely to result in more linear miles of riparian buffers but a somewhat lower ability to eliminate nutrient and sediment runoff where it exceeds the buffer's filtering capacity.

Table 3.10-1: Description of past, present, and reasonably foreseeable future actions considered in the cumulative impact analysis

Category/Projects	Project Description	Key Resource Areas with Potential for Cumulative Impacts*
Historic USDA Conservation	USDA conservation programs in portions of Newton, Lauderdale, Clarke, Neshoba, and	Short-term adverse impacts to:
		 Geology and Substrates
Program Practices	Kemper counties, Mississippi from 2010 to 2016	 Hydrology and Water Quality
2010-2016 (Project		 Habitat and Wildlife
Area)		Benefits
		 Geology and Substrates
		 Hydrology and water quality
		 Habitat and Wildlife

Category/Projects	Project Description	Key Resource Areas with Potential for Cumulative Impacts*
Future EQIP-funded Conservation Practices 2017-2021 (USDA \$1 M)	EQIP-funded conservation practices in portions of Newton, Lauderdale, Clarke, Neshoba, and Kemper counties, Mississippi from 2017 to 2021	Short-term adverse impacts to:

The following section describes the cumulative impacts of the alternatives being considered when combined with other past, present and reasonably foreseeable future actions which were identified above. In many situations, implementation of the alternatives would likely help reduce overall long-term adverse impacts by providing a certain level of offsetting benefits, especially when considered in concert with the numerous other present and reasonably foreseeable future actions in the area.

3.10.3 Cumulative impact analysis

Geology and Substrates

For implementation of the proposed NR alternatives (A and B), based on the analysis of representative, ground-disturbing (most impactful) conservation practices, there would be short-term to long-term, minor to moderate, adverse impacts to soil from soil disturbing activities such as streambank and shoreline stabilization, construction of grassed waterways, installation of grade stabilization structures, stream crossings, construction of terraces, and associated activities. There would be long-term benefits to soil because once implemented, conservation practices would reduce nutrient runoff and sedimentation of drainageways and tributaries.

Historic USDA-NRCS conservation program-funded practices and future EQIP-funded conservation practices would result in similar adverse and beneficial effects, but these practices would have small localized adverse impacts normally occurring at different times. The application of conservation practices using a systems approach that includes associated and mitigating practices would also serve to avoid and minimize adverse effects.

When the proposed NR Alternatives A and B are analyzed in combination with other past present, and reasonably foreseeable future actions, short-term cumulative adverse impacts to geology and substrates would likely occur. The alternatives would not contribute substantially to cumulative adverse impacts. The alternatives, carried out in conjunction with other conservation practices, would also have the potential to result in some long-term beneficial cumulative impacts to geology and substrates.

Hydrology and Water Quality

For implementation of the proposed NR alternatives (A and B), based on the analysis of representative, ground-disturbing conservation practices with potential for adverse effects, there would be short-term to long-term, minor to moderate, adverse impacts to hydrology and water quality from soil disturbing activities such as streambank and shoreline stabilization, construction of grassed waterways, installation of grade stabilization structures, stream crossings, construction of terraces, and associated activities. Implementation of conservation practices could result in short-term, minor changes to hydrology and short-term sedimentation resulting from the implementation of practices.

There would be long-term benefits to hydrology and water quality resulting from streambank and shoreline restoration, construction of grassed waterways, installation of grade stabilization and other conservation practices. Conservation practices would result in staged stormwater discharge, reduced nutrient runoff and sedimentation into drainageways and tributaries.

Historic USDA conservation program-funded conservation practices and future EQIP funded conservation practices would result in similar adverse and beneficial effects, but these practices would be small localized impacts that have or would occur at different times.

When the proposed NR Alternatives A and B are analyzed in combination with other past present, and reasonably foreseeable future actions, short-term cumulative adverse impacts to hydrology and water quality would likely occur. The alternatives would not contribute substantially to cumulative adverse impacts. The alternatives, carried out in conjunction with other conservation practices, would also have the potential to result in some long-term beneficial cumulative impacts to hydrology and water quality.

Habitats and Wildlife

For implementation of the proposed NR alternatives (A and B), based on the analysis of representative, ground-disturbing (most impactful) conservation practices, there would be short-term to long-term, minor to moderate, adverse impacts to habitats and wildlife from soil disturbing activities such as streambank and shoreline stabilization, stream crossings, forest stand improvements and associated activities. Implementation of conservation practices would include removal of vegetation, small, localized habitat loss, and short-term disturbance to wildlife. There would be long-term benefits to habitats and wildlife resulting from streambank and shoreline restoration, forest stand improvement, and other conservation practices that would-be habitat enhancements and would result in benefits to wildlife.

Historic USDA-NRCS conservation program-funded practices and future EQIP-funded conservation practices would result in similar adverse and beneficial effects, but these practices would be small localized impacts and likely to occur at different times.

When the proposed NR Alternatives A and B are analyzed in combination with other past present, and reasonably foreseeable future actions, short-term cumulative adverse impacts to habitats and wildlife would likely occur. The alternatives would not contribute substantially to cumulative adverse impacts. The alternatives, carried out in conjunction with other conservation practices, would also have the potential to result in some long-term beneficial cumulative impacts to habitats and wildlife.

3.11 Comparison of the Alternatives-NR (Nonpoint Source) Restoration Type

This section provides a comparison of the NEPA environmental consequences for the reasonable range of alternatives for the NR (Nonpoint Source) Restoration Type. The alternatives include two action alternatives as well as a No Action/Natural Recovery alternative and are described in Table 3.11-1.

Table 3.11-1: Comparison of the NR (Nonpoint Source) Restoration Type Alternatives

Table 5.11-1; Com	Table 3.11-1: Comparison of the NR (Nonpoint Source) Restoration Type Alternatives		
	Comparison of the NR Restoration Type Alternatives		
Alternative A (Preferred):	Upper Pascagoula River Water Quality Enhancement Project Alternative A would provide the opportunity to implement Ecological/NR conservation practices as well as Soil and Water conservation practices with willing participants, allowing for a wide array of benefits on cropland, pasture/grassland, associated agriculture lands and forestland. Under this alternative, fewer farms likely would be treated than under Alternative B because both upland and riparian area resource issues would be addressed on each farm under Alternative B which addresses only riparian areas.		
Alternative B:	Pascagoula River Basin Riparian Buffer Maintenance Plan Alternative B differs from Alternative A only in that the range of conservation practices would be limited to Ecological/NR practices would be applied in riparian areas within associated agriculture lands and forestland. Funds allocated to this alternative likely would be spread across more landowners because only resource concerns within the riparian area would be addressed, resulting in fewer practices being installed per farm. Treatments under this alternative may not prevent runoff of all nutrients and sediments where applied in areas that buffers don't have the capacity to filter it all.		
No Action Alternative:	Under the No Action Alternative, the MS TIG would not implement any projects for the NR (Nonpoint Source) Restoration Type at this time, and would instead allow natural recovery processes to occur, which could result in one of four outcomes for injured resources: 1) gradual recovery, 2) partial recovery, 3) no recovery, or 4) further deterioration. Although injured resources could presumably recover to or near baseline conditions under this scenario, recovery would take much longer compared to a scenario in which restoration actions were undertaken.		
Physical Environ	ment		
Alternative A (Preferred):	Based on the analysis of representative, ground-disturbing (most impactful) conservation practices there would be short-term to long-term, minor to moderate, adverse impacts to soil, hydrology and water quality. There would be short-term, minor to moderate, adverse impacts to wetlands. There would be long-term benefits to soil, hydrology, water quality and wetlands.		
Alternative B:	Adverse and beneficial impacts would be the same for Alternative B as Alternative A but would be restricted to riparian areas within associated agriculture lands and forestland.		
No Action Alternative:	This alternative is not expected to contribute to short-term, long term, indirect or cumulative adverse impacts to physical resources. The No Action alternative would have no beneficial impacts to water quality via NR.		
Biological Enviro	onment		
Alternative A (Preferred):	Based on the analysis of representative, ground-disturbing (most impactful) conservation practices there would be short-term to long-term, minor to moderate, adverse impacts to habitat and wildlife. There would be long-term benefits to habitat and wildlife. The following federally protected species could be present within the proposed alternative		
	project area: red-cockaded woodpecker, wood stork, northern long-eared bat; gopher tortoise, yellow-blotched map turtle, and Price's potato bean. USDA-NRCS would coordinate with the USFWS Ecological Services Field Office in Jackson, Mississippi to identify whether protected species or their habitats could be affected by site-specific conservation plans. If habitat is found to exist in a site-specific plans, either surveys/avoidance or both would be completed during the design of the conservation practice.		
	Migratory bird species groups that could occur in the proposed alternative project area include wading birds, raptors, goatsuckers, waterfowl, doves and pigeons, and rails and coots. For all planned restoration activities, pre-commencement nesting surveys for migratory birds and raptors within the site-specific project area would be conducted and if		

	Comparison of the NR Restoration Type Alternatives	
	evidence of nesting is found, USDA-NRCS would coordinate with the USFWS to develop and implement appropriate conservation measures so that no "take" of nesting birds is anticipated.	
Alternative B:	Adverse and beneficial impacts would be the same for Alternative B as Alternative A but would be restricted to riparian areas within associated agriculture lands and forestland.	
No Action Alternative:	This alternative is not expected to contribute to short-term, long term, indirect or cumulative adverse impacts to biological resources. The No Action alternative would provide no beneficial impacts, because existing conditions would not change in a predictable way.	
Socioeconomic E	nvironment	
Alternative A (Preferred):	There would be no disproportionate impacts to low-income or minority populations that would result from implementation of proposed Alternative A.	
	For site-specific conservation practices, surveys would be conducted if cultural resources are suspected in the area. Resources that are eligible for the National Register of Historic Places would be avoided in the design of the conservation practice. There would be no adverse impact to cultural resources.	
	There would be no adverse impact to public health and safety. There would be beneficial impacts to water quality in the watershed. Improved water quality is beneficial to public health since the waters in the watershed are mostly classified as "recreation" and "fish and wildlife" by the State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters (MDEQ 2012). Conservation Practices would be designed so as not result in a detectable change to natural and beneficial floodplain values. Restored hydrology streambank stabilization and grade control structures would enhance floodplain functions.	
Alternative B:	Adverse and beneficial impacts would be the same for Alternative B as Alternative A but would be restricted to riparian areas within associated agriculture lands and forestland.	
No Action Alternative	This alternative is not expected to contribute to short-term, long term, indirect, or cumulative adverse impacts to socioeconomics. The No Action alternative would provide no beneficial impacts, because existing conditions would not change in a predictable way.	
Cumulative Effec	ets	
Alternative A (Preferred) and B:	Alternative A would result in short-term adverse impacts to geology and substrates but because they are temporary would not contribute substantially to cumulative adverse impacts. Alternative A has potential to result in long-term beneficial impacts to geology and substrates. There would be minor short-term adverse impacts to water quality but there also would be long-term beneficial effects to hydrology and water quality. There would be short-term to long-term, minor to moderate, adverse impacts to habitats and wildlife from soil disturbing activities. Carried out with other past, present, and reasonably foreseeable future actions, long-term cumulative adverse impacts to geology and substrates, water quality, habitats and wildlife would not contribute substantially to cumulative adverse impacts. The alternatives, carried out in conjunction with other restoration efforts, would also have the potential to result in some long-term beneficial effects to geology and substrates, hydrology, and water quality.	
	The cumulative effects for both Alternatives A and B will be the same with the exception that Alternative A is likely to result in a higher level of treatment on fewer farms than Alternative B, and Alternative B is more likely to result in more linear miles of riparian buffers but a somewhat lower ability to eliminate nutrient and sediment runoff where it exceeds the buffer's filtering capacity.	
No Action Alternative	There would be no beneficial impacts or short or long-term cumulative adverse impacts to resources.	

The MS TIG is proposing to select Alternative A (Preferred): Upper Pascagoula River Water Quality Enhancement Project. Table 3-11- 1 above summarizes the environmental consequences for the proposed alternatives in this Draft RP/EA. Subsequent environmental review will occur in addition to this programmatic review to determine whether planned actions are at or below the maximum impacts described in this Draft RP/EA. A copy of the Environmental Evaluation Worksheet to be used to document the results of the environmental evaluation is in Appendix A. If the planned action is likely to exceed the maximum impacts described in this Draft RP/EA, USDA-NRCS will undertake additional environmental review consistent with NEPA requirements and other requirements for protection of the environment or will abandon the planned project. The MS TIG does not propose to take actions that would result in any significant adverse impacts on the environment.

The MS TIG would consider best practices referenced in Section 6.15 and Appendix 6A of the PDARP/PEIS. Additional best practices may be recommended for site-specific conservation practices in different locations due to differences in relevant conditions. The following best practices are contemplated and would be implemented to the extent practicable in order to avoid and minimize impacts to various resources:

Geology and Substrates (Soils)

- Develop and implement an erosion control plan to minimize erosion during and after construction and where possible use vegetative buffers (100 feet or greater), revegetate with native species or annual grasses, and conduct work during dry seasons.
- Develop and implement a spill prevention and response plan, including conducting daily inspections of all construction and related equipment to ensure there are no leaks of antifreeze, hydraulic fluid, or other substances and cleaning and sealing all equipment that would be used in the water to rid it of chemical residue. Develop a contract stipulation to disallow use of any leaking equipment or vehicles.
- Prohibit use of hazardous materials, such as lead paint, creosote, pentachlorophenol, and other wood preservatives during construction in, over or adjacent to, sensitive sites during construction and routine maintenance.

Hydrology and Water Quality

- In the design of conservation practices the MS TIG would consider resiliency measures related to increasing storm intensities and changing weather patterns (CEQ, 2016).
- Develop and implement an erosion control plan to minimize erosion during and after construction and where possible use vegetative buffers (100 feet or greater), revegetate with native species or annual grasses, and conduct work during dry seasons.
- Develop and implement a spill prevention and response plan, including conducting daily
 inspections of all construction and related equipment to ensure there are no leaks of
 antifreeze, hydraulic fluid, or other substances and cleaning and sealing all equipment that
 would be used in the water to rid it of chemical residue. Develop a contract stipulation to
 disallow use of any leaking equipment or vehicles.
- Prohibit use of hazardous materials, such as lead paint, creosote, pentachlorophenol, and other wood preservatives during construction in, over or adjacent to, sensitive sites during construction and routine maintenance.
- Avoid and minimize, to the maximum extent practicable, placement of dredged or fill material in wetlands and other aquatic resources.

- Design construction equipment corridors to avoid and minimize impacts to wetlands and other aquatic resources to the maximum extent practicable.
- To the maximum extent possible, implement the placement of sediment to minimize impacts to existing vegetation or burrowing organisms.
- Apply herbicide in accordance with the direction and guidance provided on the appropriate U.S. Environmental Protection Agency (EPA) labels and state statutes during land-based activities.
- When local conditions indicate the likely presence of contaminated soils and sediments, test soil samples for contaminant levels and take precautions to avoid disturbance of, or provide for proper disposal of, contaminated soils and sediments. Evaluate methods prior to dredging to reduce the potential for impacts from turbidity or tarballs.
- Designate a vehicle staging area removed from any natural surface water resource or wetland to perform fueling, maintenance, and storage of construction vehicles and equipment.
 Inspect vehicles and equipment daily prior to leaving the storage area to ensure that no petroleum or oil products are leaking.
- Use silt fencing where appropriate to reduce increased turbidity and siltation in the project vicinity. This would apply to both on land and in water work.

Habitat and Wildlife

 All equipment to be used during the project, including personal gear, would be inspected and cleaned such that there is no observable presence of mud, seeds, vegetation, insects and other species.

Protected Species

Northern long-eared bat

- The USFWS listed the northern long-eared bat as threatened under the Endangered Species Act on May 4, 2015, and established an interim 4(d) rule to help protect the species. The Endangered Species Act 4(d) rule for the northern long-eared bat best practices are outlined below:
- No purposeful take of northern long-eared bats within the bat's range with the exception of removal from human structures, defense of human life (public health monitoring for rabies, and removal of hazardous trees for protection of human life and property.
- Additionally, no incidental take of northern long-eared bats is allowed within a
 hibernacula, if it results from tree removal activities within 0.25 mile of a known
 hibernacula, and if it results from tree removal activities that cut or destroy a known,
 occupied maternity roost tree or other trees within 150 feet of a known, occupied maternity
 roost tree during June and July (MSU, 2016).

Red-Cockaded Woodpecker

- Avoid working within active red-cockaded woodpecker clusters (the minimum convex polygon containing the aggregation of cavity trees used by a group of red-cockaded woodpeckers and a 200- foot-wide buffer surrounding the polygon).
- If avoidance is not possible or management activities in red-cockaded woodpecker suitable habitat are desired, conduct standard surveys to determine if the habitat is supporting any individuals or presence can be assumed. If red-cockaded woodpeckers are present (or

- assumed to be), avoid cavity trees and use mechanized equipment during the non-nesting season (approximately April 1 through July 31).
- If tree removal is necessary, survey pine trees approximately 60 or more years old for active cavities within one year of the proposed removal. Extend surveys from the project site out to no less than one- half mile. Replace any cavities affected by the project via drilled cavity construction.
- If impacts to suitable foraging habitat (pines approximately 30 or more years old and within one-half mile of an active cavity tree) are proposed, conduct a foraging habitat analysis. Foraging habitat may need to be replanted post-project.
- Design projects within red-cockaded woodpecker suitable habitat such that prescribed fire needs are not impeded.

Gopher Tortoise

• If suitable habitat is present, coordinate with the USFWS Ecological Services Field Office in Jackson, MS to discuss the need for surveys to identify any gopher tortoise burrows and to develop conservation measures to avoid or minimize impacts. Measures could include establishing a protective buffer (size determined by USFWS and the state trust resource agency) if burrows are within the project area and cannot be avoided, implementing standard procedures to relocate the tortoise within the project site but away from the areas of construction or restoration or considering conservation banks. A Candidate Conservation Agreement with Assurances may be appropriate for project sites within the nonlisted range of the species.

Protected Plants

- If suitable habitat is present, coordinate with the USFWS Ecological Services Field Office in Jackson, MS to discuss the need for surveys to identify protected plants and to develop conservation measures to avoid or minimize impacts.
- Enhance and protect plants on site and in adjacent habitats to the maximum extent possible.
- Provide all individuals working on a project with information in support of general awareness of and means to avoid impacts to protected species and their habitats present at the specific project site.

Migratory Birds

- Use care to avoid birds when operating machinery or vehicles near birds.
- Avoid working in migratory bird nesting habitats during breeding, nesting, and fledging
 (approximately mid-February through late August). If project activities must occur during
 this timeframe and breeding, nesting, or fledging birds are present, contact the state trust
 resource agency to obtain the most recent guidance to protect nesting birds or rookeries, and
 their recommendations will be implemented.
- Conservation areas may already be marked to protect bird nesting areas. Stay out of existing marked areas.
- If vegetation clearing is necessary, clear vegetation outside the migratory bird nesting season (approximately mid-February through late August) or have a qualified biologist inspect for

active nests. If no active nests are found, vegetation may be removed. If active nests are found, vegetation may be removed after the nest successfully fledges.

Bald Eagles

- If bald eagle breeding or nesting behaviors are observed or a nest is discovered or known, have all activities avoid the nest by a minimum of 660 feet. If the nest is protected by a vegetated buffer where there is *no* line of sight to the nest, then the minimum avoidance distance is 330 feet. Maintain this avoidance distance from the onset of breeding/courtship behaviors until any eggs have hatched and eaglets have fledged (approximately 6 months).
- If a similar activity (such as driving on a roadway) is closer than 660 feet to a nest, maintain a distance buffer as close to the nest as the existing tolerated activity. If a vegetated buffer is present and there is no line of sight to the nest and a similar activity is closer than 330 feet to a nest, then maintain a distance buffer as close to the nest as the existing tolerated activity.
- In some instances, activities conducted within 660 feet of a nest may result in disturbance, particularly for the eagles occupying the Mississippi barrier islands. If an activity appears to cause initial disturbance, stop the activity and move all individuals and equipment away until the eagles are no longer displaying disturbance behaviors. Contact the USFWS's Migratory Bird Permit Office to determine how to avoid impacts or if a permit may be needed.

4.0 Compliance with Other Laws and Regulations

Additional federal and state laws may apply to the proposed projects considered in this Draft RP/EA. Legal authority applicable to restoration project development were fully described in the context of the *DWH* restoration planning in the PDARP/PEIS, Section 6.9 Compliance with Other Applicable Authorities and Appendix 6.D, Other Laws and Executive Orders. That material is incorporated by reference here.

Federal environmental compliance responsibilities and procedures will follow the Trustee Council Standard Operating Procedures for Implementation of the Natural Resource Restoration for the Deepwater Horizon (*DWH*) Oil Spill, which are laid out in Section 9.4.6 of that document. Following these standard operating procedures, the implementing Trustee for each project will ensure that the status of environmental compliance (e.g., completed versus in progress) is tracked through the Restoration Portal. Implementing Trustees will keep a record of compliance documents (e.g., ESA biological opinions, USACE permits) and ensure that they are submitted for inclusion to the Administrative Record. The MS TIG will ensure compliance with all applicable laws and regulations.

4.1 Additional Federal Laws

Additional federal laws, regulations, and executive orders that may be applicable include but are not limited to:

- Endangered Species Act
- Magnuson-Stevens Fishery Conservation and Management Act
- Marine Mammal Protection Act
- Coastal Zone Management Act
- National Historic Preservation Act
- Coastal Barrier Resources Act
- Migratory Bird Treaty Act
- Bald and Golden Eagle Protection Act
- Clean Air Act
- Clean Water Act
- Rivers and Harbors Act
- Marine Protection, Research and Sanctuaries Act
- Estuary Protection Act
- Archaeological Resource Protection Act
- National Marine Sanctuaries Act
- Farmland Protection Policy Act
- Private Aids to Navigation (C.F.R. Title 33, Chapter 1, Part 66)
- Federal Water Pollution Control Act
- Additional Executive Orders
 - o EO 11988: Floodplain Management
 - o EO 11990: Protection of Wetlands
 - o EO 12898: Environmental Justice

- o EO 12962: Recreational Fisheries
- o EO 13112: Invasive Species
- o EO 13175: Consultation and Coordination with Indian Tribal Governments
- o EO 13186: Responsibilities of Federal Agencies to Protect Migratory Birds
- o EO 13653: Preparing the United States for the Impacts of Climate Change
- o EO 13693: Planning for Federal Sustainability in the Next Decade

4.2 Additional State Laws

Potentially applicable state laws may include but are not limited to:

- Public Trust Tidelands, Miss. Code Ann. §29-1-1 et seq.
- Antiquities Law of Mississippi, Miss. Code Ann. §39-7-1 et seq.
- Mississippi Air and Water Pollution Control Law, Miss Code Ann. § 49-17-1 et seq.
- Coastal Wetlands Protection Act, Miss. Code Ann. § 49-27-1 et seq.
- Marine Resources, Miss. Code Ann. 57-15-1 et seq.

5.0 Draft Monitoring and Adaptive Management Plan

5.1 Introduction

Monitoring, Adaptive Management, and Administrative Oversight was identified as one of the Programmatic Trustee Goals in the PDARP/PEIS. As described in Chapter 5, Appendix E of the PDARP/PEIS, the Trustee Council has committed to a Monitoring and Adaptive Management (MAM) Framework to support restoration activities by infusing best available science into project planning and design, identifying and reducing key uncertainties, tracking and evaluating progress toward restoration goals, determining the need for corrective actions, and supporting compliance monitoring.

The *DWH* NRDA MAM Framework provides a flexible, science-based approach to effectively and efficiently implement restoration over several decades that provides long-term benefits to the resources and services injured by the *DWH* oil spill. MAM plans identify the monitoring needed to evaluate progress toward meeting site-specific objectives and to support adaptive management of the restoration project.

The MAM plans for the three preferred project alternatives are attached as Appendix D, E and F of this Draft RP/EA. MAM Plans are living documents and they will be updated as needed to reflect changing conditions and/or to incorporate new information. For example, the plan may need to be revised if the project design changes, if initial data analysis indicates that the sampling design is inadequate, or if any uncertainties are resolved or new uncertainties are identified during project implementation and monitoring. Any future revisions to these documents will be made publicly available through the Restoration Portal via web links provided here. Full monitoring plans for each project can also be accessed through the web link.

5.2 Summary of Restoration Goals, Objectives, and Performance Criteria

Proposed Project Alternative: Graveline Bay Land Acquisition and Management

Restoration activities include acquisition of up to 1,410 acres of habitat in the vicinity of the Graveline Bay CP and restoration and management activities in the existing and expanded Graveline Bay CP. Management activities would include access restriction, chemical treatment, mechanical treatment, prescribed fire, debris removal and road repair/culvert replacement. This project is intended to restore habitats and resources injured from the *DWH* oil spill, including foraging habitat for multiple bird species. Additional ecosystem services that are provided include preservation of buffer habitat for coastal marsh to promote long-term health of coastal habitats and the species that inhabit and utilize the habitat for reproduction, foraging, and shelter. The MDEQ will act as the Implementing Trustee on behalf of the MS TIG working with the MDMR CP Program. The Draft MAM Plan for the Graveline Bay Land Acquisition and Management proposed alternative is included as Appendix D of this Draft RP/EA. A summary of goals and objectives are provided here.

Goal 1: Restore and Conserve Habitat

Objectives

- 4) Protect estuarine marsh, shoreline (beach) and other coastal riparian habitats from development and increase habitat connectivity to other large conservation parcels, by acquiring priority lands in the Graveline Bay CP for conservation.
- 5) Increase and maintain native vegetation species composition in restored habitats within Graveline Bay CP.
 - a. Performance Criteria: Acquisition
 - i. Fee-simple acquisition of priority habitats in the project area of 1,410 acres
 - b. Performance Criteria: Management
 - i. Vegetation structure for fire-suppressed pine savanna (by year 5)

20-65% canopy cover of longleaf or slash pine
40-100% herbaceous cover
Invasive nonnative plant species in any stratum present but sporadic (1-5 % cover)

ii. Vegetation Composition

95% native flora

Goal 2: Replenish and Protect Living Coastal and Marine Resources

Objectives

- 6) Increase and maintain shorebird (species injured by the *DWH* oil spill) use of beach habitat
- 7) Increase and maintain wading bird habitat (species injured by the *DWH* oil spill) use in acquired habitats
 - a. Performance Criteria: Increase shorebird habitat use by year 5
 - b. Performance Criteria: Maintain wading bird habitat use by year 5

Adaptive Management: The adaptive management approach to the Graveline Bay Land Acquisition and Management proposed alternative is detailed in the Draft MAM Plan (Appendix D). It includes interim performance criteria for helping determine whether adjustments to the project are needed to better ensure the project meets the final performance criteria used to determine project success, as well as the potential adaptive management actions (e.g., mid-course corrections or corrective actions) that may be considered for individual parameters. The Draft MAM Plan includes a list of potential adaptive management actions for each parameter to be considered. Parameters include acres acquired, invasive species, vegetation structure and composition, shorebird and wading bird diversity and abundance. The Draft MAM Plan does not include all possible options; rather, it includes a list of potential adaptive management actions for each individual parameter to be considered. The decision to implement a corrective action should holistically consider the overall outcomes of the restoration project by assessing the results of all monitoring parameters.

Proposed Project Alternative: Grand Bay Land Acquisition and Habitat Management

This restoration project is being implemented within the proposed alternative boundary which includes of the Grand Bay NWR, Grand Bay NERR and Grand Bay Savanna CP. Restoration activities involve the acquisition of private parcel inholdings and restoration of habitats, where applicable. This project is intended to help restore habitats and resources injured from the *DWH* oil spill, including coastal, estuarine, and riparian habitats; and birds. The Implementing Trustees include would MDEQ and DOI working with USFWS and MDMR Mississippi CP Program) and NOAA (as the joint managers of the Grand Bay NERR). The Draft MAM Plan for the Grand Bay Land Acquisition and Habitat Management proposed alternative is included as Appendix E of this Draft RP/EA. A summary of goals and objectives are provided here.

Goal 1: Restore and Conserve Habitat Objectives

- 1. Acquire lands and implement management techniques to increase and maintain native vegetation species composition in restored habitats;
 - a. Performance Criteria: Acquisition
 - i. Fee-simple acquisition of priority habitats in the project area up to 8,000 acres
 - b. Performance Criteria: Management
 - ii. Vegetation structure for coastal pine savanna habitat (by year 5)
 - <20% canopy cover of longleaf or slash pine
 - 40-100% herbaceous cover
 - Invasive nonnative plant species in any stratum present but sporadic (1-5 % cover)
 - iii. Vegetation Composition for coastal pine savanna habitat
 - 95% native flora
 - iv. Base-line habitat characteristics of high quality open pine savanna habitat
 - Use of habitat by wintering Henslow's sparrow (*Ammodramus henslowii*) (presence/absence)

Goal 2: Replenish and Protect Living Coastal and Marine Resources Objectives

- 1. Acquire lands and implement management techniques to increase bird diversity, abundance, and habitat utilization
 - a. Performance Criteria: Use of habitat by injured wading bird species

Adaptive Management: The adaptive management approach to the Grand Bay Land Acquisition and Habitat Management proposed alternatives is detailed in the Draft MAM Plan (Appendix E). It includes interim performance criteria for helping determine whether adjustments to the project are needed to better ensure the project meets the final performance criteria used to determine project success, as well as the potential adaptive management actions (e.g., mid-course corrections or corrective actions) that may be considered for individual parameters. The Draft MAM plan includes a list of potential adaptive management actions for each parameter to be considered. Parameters include acres acquired, invasive species, vegetation structure and composition, and bird species monitoring. The Draft MAM Plan does not include all possible options; rather, it includes a list of potential adaptive management actions for each individual parameter to be considered. The decision to implement a corrective action should holistically consider the overall outcomes of the restoration project by assessing the results of all monitoring parameters.

Proposed Project Alternative: Upper Pascagoula River Water Quality Enhancement

Restoration involves the implementation of agricultural conservation practices to reduce sediment, phosphorus, and nitrogen loadings in target watersheds, and to downstream coastal receiving waters. The proposed conservation practices would reduce nutrient losses from the landscape, reduce nutrient loads to streams and downstream receiving waters, and reduce water quality degradation in watersheds thus providing benefits to marine resources and benefits to coastal watersheds. This project is intended to reduce nutrient and sediment load contribution in watersheds that contain Gulf sturgeon [*Acipenser oxyrinchus desotoi*]) Critical Habitat. The Gulf sturgeon is anadromous, spending much of its life in marine environments, but spawning in the Upper Pascagoula River and tributaries. Sediment and other pollutants may reduce Gulf sturgeon spawning success. The Implementing Trustees are MDEQ, USDA and EPA working with USDA-NRCS. The Draft MAM Plan for the Upper Pascagoula Water Quality Enhancement proposed alternative is included as Appendix F of this Draft RP/EA. A summary of goals and objectives are provided here.

Goal 1: Restore Water Quality Objectives

- 1. Reduce sediment, phosphorus, and nitrogen loads leaving private lands in prioritized watersheds in the Pascagoula Basin
 - a. Performance Criteria: x kg of suspended sediments trapped from treatment site; x kg of phosphorous trapped from treatment site; x kg of nitrogen trapped from treatment site Goal
- 2. Identify instream habitat features that are influenced by upstream sediment and nutrient loads for future instream resource benefits.
 - a. Performance Criteria: N/A

Adaptive Management: Adaptive management on specific conservation practices being implemented beyond inspection and maintenance is not anticipated for this project. Monitoring information from this restoration project would be critical to refine targeting of conservation practice implementation, refining in-stream habitat use by Gulf sturgeon if found, as well as identifying instream habitat that could be enhanced by conservation practices for Gulf sturgeon use as needed.

5.3 MAM Plan Administration

MAM Plans are living documents and will be updated as needed to reflect changing conditions and/or to incorporate new information as projects progress and are implemented. For example, the plan may need to be revised if the project design changes, if initial data analysis indicates that the sampling design is inadequate, or if any uncertainties are resolved or new uncertainties are identified during project implementation and monitoring. Any future revisions to this document and individual project MAM plans will be made publicly available through the Restoration Portal via web links provided here. Full monitoring plans include descriptive information regarding monitoring goals, objectives, parameter details (e.g. methodology, sample size, timing/frequency), project-level decisions, and monitoring schedules and budgets.

6.0 List of Preparers and Reviewers

AGENCY/FIRM	NAME	POSITION
Mississippi DEQ		
MDEQ	Tabatha Baum	Attorney
Balch & Bingham LLP	Bradley A. Ennis	Attorney
Covington Civil & Environmental, LLC	Stephen Parker	Senior Scientist
Covington Civil & Environmental, LLC	Alane C. Young	Senior Geologist
Covington Civil & Environmental, LLC	Thomas Strange	Senior Scientist
Covington Civil & Environmental, LLC	Morgan Boudreaux	Biologist
Covington Civil & Environmental, LLC	Christopher W. Thomas	Project Scientist
NATIONAL OCEANIC AND ATMOSPHERIC ADM	INISTRATION	
National Oceanic and Atmospheric Administration	Corinna Mc Mackin	Attorney
National Oceanic and Atmospheric Administration	Dan Van Nostrand	Marine Habitat Resource Specialist
Earth Resources Technology	Ramona Schreiber	Marine Habitat Resource Specialist
U.S. DEPARTMENT OF AGRICULTURE		
United States Department of Agriculture Natural Resource Conservation Service, Gulf Coast Ecosystem Restoration Team	Ronald Howard	Program Specialist
United States Department of Agriculture Natural Resource Conservation Service	Andrée DuVarney	National Environmental Coordinator
U.S. ENVIRONMENTAL PROTECTION AGENCY		
Gulf of Mexico Program	Troy Pierce	Chief Scientist
Office of Wetlands, Oceans, and Watersheds	Erika Larsen	Physical Scientist
U.S. DEPARTMENT OF THE INTERIOR		
U.S. Department of Interior	Jon Hemming	DWH NRDAR Field Office Supervisor
U.S. Fish and Wildlife	Brian Spears	Restoration Program Manager
U.S. Department of Interior	Robin Renn	NEPA Coordinator

7.0 Literature Cited

- Council on Environmental Quality (CEQ). 2005. Guidance on Consideration of Past Actions in Cumulative Effects Analysis. Memorandum from James L. Connaughton, Chairman. June 24.
- CEQ. August 1, 2016. Memorandum for Heads of Federal Agencies. Final Guidance for Federal Departments and Agencies on Consideration of Greenhouse Gas Emissions and the Effects of Climate Change in National Environmental Policy Act Reviews.
- Ernst, C. H., J. E. Lovich, and R. W. Barbour. 1994. Turtles of the United States and Canada. Smithsonian Institution Press, Washington, D.C. 578 pp.
- Fertl, D., A.J. Schiro, G.T. Regan, C.A. Beck, N. Adimey, L. Price-May, A. Amos, G.A.J. Worthy, and R. Crossland. 2005. Manatee Occurrence in the Northern Gulf of Mexico, West of Florida. Gulf and Caribbean Research 17:69-94.
- Frey, J. 2014. 2014. Letter to James Davis, MDMR. Mississippi Department of Wildlife, Fisheries and Parks review of Construction of Recreational Park In Biloxi, Harrison County, Mississippi. R#10035, DMR-03400. Mississippi Natural Heritage Program. March 31.
- Grand Bay National Estuarine Research Reserve (GBNERR). 2013. Grand Bay National Estuarine Research Reserve Management Plan 2013-2018: Grand Bay National Estuarine Research Reserve, Mississippi Department of Marine Resources, Moss Point, Mississippi. 104pp.
- GBNERR. 2016. Invasive Species Management Plan for Grand Bay National Estuarine Research Reserve. Available at: http://grandbaynerr.org/wp-content/uploads/2010/08/InvasiveSpeciesManagementPlan.pdf.
- Mississippi Department of Environmental Quality (MDEQ). 2012. State Of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters. Adopted by Mississippi Commission on Environmental Quality: June 28. Office of Pollution Control. Jackson, Mississippi.
- MDEQ. 2014. Title 11: Mississippi Department of Environmental Quality Part 6: Wastewater Pollution Control Regulations Part 6, Chapter 9: Mississippi Commission on Environmental Quality, Mississippi 2014 Section 303(d) List of Impaired Water Bodies FINAL LIST Version 5.0 For Commission Approval.
- MDEQ. 2015. Air Quality Data Summary. Available at: http://www.deq.state.ms.us/mdeq.nsf/pdf/Air_2015AirQualityDataSummary/\$File/2015%20Air%20Quality%20Data%20Summary.pdf?OpenElement.
- MDEQ and National Fish and Wildlife Foundation. 2015. The Mississippi Gulf Coast Restoration Plan A Path Toward Sustainable Ecosystem Restoration. Available at http://msrestoreteam.com/NFWFPlan2015/#p=1.
- Mississippi Department of Marine Resources (MDMR). 1998. Grand Bay National Estuarine Research Reserve Final Environmental Impact Statement/Reserve Management Plan. November.

- MDMR. 2016. Graveline Bay Preserve. Mississippi Gulf Ecological Management Site. Available at: http://www.dmr.ms.gov/index.php/mississippi-gems/215-graveline-bay.
- Mississippi Department of Wildlife, Fisheries, and Parks (MDWFP). 2001. Endangered Species of Mississippi. Mississippi Museum of Natural Science. Jackson Mississippi.
- Mississippi Museum of Natural Science (MMNS). 2015. Mississippi State Wildlife Action Plan. Mississippi Department of Wildlife, Fisheries, and Parks, Mississippi Museum of Natural Science, Jackson, Mississippi.
- Mississippi Natural Heritage Program. 2015. Special Animals Tracking List. Museum of Natural Science, Mississippi Dept. of Wildlife, Fisheries, and Parks, Jackson, MS. 11 pp.
- Mississippi State University (MSU) Extension. 2016. Northern Long-Eared Bat and Mississippi Forestry: Impacts of the Endangered Species Act Publication 2935 (POD-02-16).
- The National Coalition of Prescribed Fire Councils. 2007. Guide to Smoke Management. September Version. Available at:

 http://www.garxfire.com/pdf%20files/The_National_Coalition_of_Prescribed_Fire_Councils_Guide_to_Smoke_Management.pdf.
- National Marine Fisheries Service (NMFS). 2016. Memorandum from NMFS to NOAA Re: Framework Biological Opinion of Deepwater Horizon Oil Spill Final Programmatic Damage Assessment and Restoration Plan and Final Programmatic Environmental Impact Statement (ESA Section 7 Consultation Response). February 10, 2016.
- Natural Resources Conservation Service (NRCS) and Forest Service. 2011. Basic Smoke Management Practices. U.S. Department of Agriculture. October. Available at: https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/16/stelprdb1046311.pdf.
- NRCS. 2016. Web Soil Survey for Jackson County. U.S. Department of Agriculture. Available at: http://websoilsurvey.sc.egov.usda.gov/App/HomePage.htmhyr.
- NRCS. 2016a. List of Hydric Soils. National List; all states (December 2015). U.S. Department of Agriculture. Available at: https://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/use/hydric/.
- National Oceanic and Atmospheric Administration (NOAA) Fisheries. 2014a. Hawksbill Turtle (*Eretmochelys imbricata*). NOAA Fisheries Office of Protected Resources. Available at: http://www.nmfs.noaa.gov/pr/species/turtles/hawksbill.htm.
- NOAA Fisheries. 2014b. Kemp's Ridley Turtle (*Lepidochelys kempii*). NOAA Fisheries Office of Protected Resources. Available at: http://www.nmfs.noaa.gov/pr/species/turtles/kempsridley.htm.
- NOAA Fisheries. 2014c. Loggerhead Turtle (*Caretta caretta*). NOAA Fisheries Office of Protected Resources. Available at: http://www.nmfs.noaa.gov/pr/species/turtles/loggerhead.htm.
- NOAA Fisheries. 2015. Green Turtle (*Chelonia mydas*). NOAA Fisheries Office of Protected Resources. Available at: http://www.nmfs.noaa.gov/pr/species/turtles/green.htm.

- Neel-Schaffer. 2009. Gautier Comprehensive Plan. City of Gautier, Mississippi. June 15.
- Niles, L.J., H.P. Sitters, A.D. Dey, P.W. Atkinson, A.J. Baker, K.A. Bennett, K.E. Clark, N.A. Clark, C. Espoz, P.M. Gonzalez, B.A. Harrington, E.E. Hernandez, K.S. Kalasz, R. Matus, C.D.T. Minton, R.I.G. Morrison, M.K. Peck, and I.L. Serrano. 2007. Status of the red knot (*Calidris canutus rufa*) in the Western Hemisphere. Report to the U.S. Fish and Wildlife Service. New Jersey Department of Environmental Protection, Division of Fish and Wildlife, Endangered and Nongame Species Program, Trenton, New Jersey. 236p.
- Planning Works, LLC. 2010. City of Ocean Springs Comprehensive Plan. Public Review Draft. March.
- Priddy, R.R., Crisler, R.M., Jr., Seliren, C.P., Powell, J.D., Burford, H., 1955. Sediments of Mississippi Sound and inshore waters: Mississippi Geological Survey Bulletin 82, 54 pp.
- Rogillio, H.E., Ruth, R.T., Behrens, E.H., Doolittle, C.N., Granger, W.J., and Kirk, J.P. 2007. Gulf sturgeon movements in the Pearl River drainage and the Mississippi Sound. North American Journal of Fisheries Management 27: 89–95.
- Ross, S. T., Slack, W.Y., Heise, R.J., Dugo, M.A., Rogillio, H., Bowen B.R., Mickle, P. and Heard, R. 2009. Estuarine and coastal habitat use of Gulf sturgeon (*Acipenser oxyrinchus desotoi*) in the North-Central Gulf of Mexico. Estuar. Coast 32. 360-364. 360–374.
- Schmid, K. 2015. Geology and Geomorphology of the Coastal Counties in Mississippi Alabama. Available at: http://geology.deq.state.ms.us/coastal/NOAA_DATA/Publications/Publications/Coastwide/Geology%20and%20Geomorphology%20of%20the%20Coastal%20Counties.pdf.
- Shaver D.J. and Rubio C. 2008. Post-nesting movement of wild and head-started Kemp's ridley sea turtles (*Lepidochelys kempii*) in the Gulf of Mexico. Endanger. Species Res.; 4:43–55.
- Stewart, R.A. 2003. Physiographic regions of Mississippi. Handout, Department of Biological Sciences, Delta State University, 6 pp. (with addenda by S.P. Faulkner, 2005).
- Trustee Council for the Deepwater Horizon. 2016. Plan for Deepwater Horizon Oil Spill Natural Resource Injury Restoration: An Overview. April.
- U.S. Army Corps of Engineers (USACE). 2009. Mississippi Coastal Improvements Program (MsCIP) Hancock, Harrison, and Jackson Counties, Mississippi Comprehensive Plan and Integrated Programmatic Environmental Impact Statement.
- USACE. 2009. Mississippi Coastal Improvements Program (MsCIP) Hancock, Harrison, and Jackson Counties, Mississippi Comprehensive Plan and Integrated Programmatic Environmental Impact Statement.
- USFWS. 2005. Draft Mississippi Sandhill Crane National Wildlife Refuge Biological Report. May.
- USFWS. 2007. Mississippi Sandhill Crane National Wildlife Refuge Comprehensive Conservation Plan; U.S. Department of Interior Fish and Wildlife Service; Southeast Region; September.USFWS. 2008. Comprehensive Conservation Plan and Environmental

- Assessment; Grand Bay National Wildlife Refuge; Jackson County, Mississippi and Mobile County, Alabama. U.S. Department of Interior Fish and Wildlife Service; Southeast Region; March.
- USFWS. 2010. Habitat Descriptions Federally Endangered and Threatened and Candidate Species of Mississippi. Available at: http://www.fws.gov/mississippiES/pdf/T&E%20Species%20Habitat%20info%202010%20up dated.pdf.
- USFWS. 2012. Land Protection Plan and Final Environmental Assessment for the Expansion of Grand Bay National Wildlife Refuge. Southeast Region. October. Available online at https://restorethegulf.gov/sites/default/files/documents/images/Grand%20Bay%20Final%20L and%20Protection%20Plan%20and%20Environmental%20Assessment.pdf.
- USFWS. 2012a. Louisiana quillwort (*Isoetes louisianensis*) 5-year Review: Summary and Evaluation. June.
- USFWS. 2013. Habitat Descriptions Federally Endangered and Threatened and Candidate Species of Mississippi. Mississippi Field Office. February.
- USFWS. 2016. Memorandum from FWS to DOI Re: Biological and Conference Opinion for the Deepwater Horizon Final Programmatic Damage Assessment and Restoration Plan (ESA Section 7 Consultation Response). March 18.
- USFWS. 2016a. Resource Management. Grand Bay. National Wildlife Refuge. Available at: https://www.fws.gov/refuge/Grand_Bay/what_we_do/resource_management.htm.
- USFWS. 2016b. Prescribed Fire Beneficial to the Land, the Wildlife and the People. Mississippi Sandhill Crane National Wildlife Refuge. Available at: https://www.fws.gov/mississippisandhillcrane/fire.html.
- Vittor, B. A. and Associates, Inc. 1982. Benthic Macroinfauna Community Characterizations in Mississippi Sound and Adjacent Waters. Contract Report to U. S. Army Corps of Engineers, Mobile District, Mobile, Alabama. 442 pp.

Appendix A. Draft MS TIG Post NRDA Restoration Plan and EA – Example Environmental Evaluation Worksheet

U.S. Department of Agriculture	NRCS	S-CPA-52					
Natural Resources Conservation Se		4/2013	IA Client Name:				
ENVIRONMENTAL E	EVALUATION WORKSHE	EET					
D. Client's Objective(s) (purpose):		C. Identification # (farm, trac		#, etc. as required):			
E. Need for Action:	H. Alternatives						
	No Action √ if RMS	s 🔲	Alternative 1 √ if RM	S 📗	Alternative 2 √ if RM	s 🗌	
			rce Concerns				
	ze, record, and address cond source Planning Criteria for g			ces Inv	entory process.		
F. Resource Concerns	I. Effects of Alternatives	uluanc	.e.j.				
and Existing/ Benchmark	No Action		Alternative 1		Alternative 2		
Conditions (Analyze and record the	Amount, Status,	√ if	Amount, Status,	√ if	Amount, Status,	√ if	
existing/benchmark	Description	does NOT	Description	does NOT	Description	does	
conditions for each	(Document both short and	meet PC	(Document both short and	meet PC	(Document both short and	meet PC	
identified concern) SOIL: EROSION	long term impacts)		long term impacts)	10	long term impacts)	10	
SOIL. ERUSION							
		NOT meet		NOT meet		NOT meet	
		PC		PC		PC	
		NOT meet		NOT meet		NOT meet	
		PC		PC		PC	
SOIL: SOIL QUALITY DEGR	RADATION						
		NOT		NOT		NOT	
		meet PC		meet PC		meet PC	
		NOT meet		NOT meet		NOT meet	
		PC		PC		PC	
WATER: EXCESS / INSUFF	ICIENT WATER						
		NOT		NOT		NOT	
		meet PC		meet PC		meet PC	
WATER: WATER QUALITY	DEGRADATION			10		10	
WATER. WITER GONEITT	BEONIBATION					ПП	
	1	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \					
	1	NOT meet		NOT meet		NOT meet	
		PC		PC		PC	
	1	NOT		NOT		NOT	
		meet		meet		meet	
	1	PC		PC		PC	

F. Resource Concerns	I. (continued)					
and Existing/ Benchmark	No Action		Alternative 1		Alternative 2	
Conditions	Amount, Status,	√if	Amount, Status,	√ if	Amount, Status,	√if
(Analyze and record the	Description	does	Description	does	Description	does
existing/benchmark conditions for each		NOT meet		NOT meet		NOT meet
identified concern)	(Document both short and long term impacts)	PC	(Document both short and long term impacts)	PC	(Document both short and long term impacts)	PC
AIR: AIR QUALITY IMPACTS			iong term impacts)		iong term impacts)	
AIN. AIN QUALITTIMI AUTO						
		Ш		ш		Ш
		NOT		NOT		NOT
		meet PC		meet PC		meet PC
						ш
		NOT meet		NOT meet		NOT meet
		PC		PC		PC
PLANTS: DEGRADED PLAN	IT CONDITION					
		NOT		LIOT.		NOT
		NOT meet		NOT meet		NOT meet
		PC		PC		PC
		NOT		NOT		NOT
		meet		meet		meet
		PC		PC		PC
ANIMALS: INADEQUATE H	ABITAT FOR FISH AND WILD	LIFE				
		NOT		NOT		NOT
		meet		meet		meet
		PC		PC		PC
ANIMALS: LIVESTOCK PRO	DUCTION LIMITATION	ī				ı
		NOT		NOT		NOT
		meet		meet		meet
		PC		PC		PC
		NOT		NOT		NOT
		meet		meet		meet
ENERGY: INEFFICIENT EN	FDCV USF	PC		PC		PC
LINERUT. INEFFICIENT ENL	LNOT USE					
		Ш				Ш
		NOT		NOT		NOT
		meet PC		meet PC		meet PC
				. 0		. ,
		Ш				Ш
		NOT		NOT		NOT
		meet PC		meet PC		meet PC
HUMAN: ECONOMIC AND S	SOCIAL CONSIDERATIONS					
	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2					

Special Environmental Concerns: Environmental Laws, Executive Orders, policies, etc.

In Section "G" complete and attach Environmental Procedures Guide Sheets for documentation as applicable. Items with a "•" may require a federal permit or consultation/coordination between the lead agency and another government agency. In these cases, effects may need to be determined in consultation with another agency. Planning and practice implementation may proceed for practices not involved in consultation.

•	J. Impacts to Special Environmental Concerns					
Concerns	No Action		Alternative 1		Alternative 2	
(Document existing/ benchmark conditions)	Document all impacts (Attach Guide Sheets as applicable)	√if needs further action	Document all impacts (Attach Guide Sheets as applicable)	√ if needs further action	Document all impacts (Attach Guide Sheets as applicable)	√ if needs further action
•Clean Air Act Guide Sheet FS1 FS-2						
Clean Water Act / Waters of the						
U.S. Guide Sheet Fact Sheet						
Coastal Zone Management Guide Sheet Fact Sheet						
Coral Reefs Guide Sheet Fact Sheet						
Cultural Resources / Historic Properties Guide Sheet Fact Sheet						
Endangered and Threatened Species Guide Sheet Fact Sheet						
Environmental Justice Guide Sheet Fact Sheet						
Essential Fish Habitat Guide Sheet Fact Sheet						
Floodplain Management Guide Sheet Fact Sheet						
Invasive Species Guide Sheet Fact Sheet						
Migratory Birds/Bald and Golden Eagle Protection Act Guide Sheet Fact Sheet						
Natural Areas Guide Sheet Fact Sheet						
Prime and Unique Farmlands Guide Sheet Fact Sheet						
Riparian Area Guide Sheet Fact Sheet						
Scenic Beauty Guide Sheet Fact Sheet						
						-


Wetlands Guide Sheet	Fact Sheet						
•Wild and Scenic Guide Sheet	Rivers Fact Sheet						
K. Other Agen Broad Public (No Action		Alternative 1		Alternative 2	
Easements, Perm Review, or Permit Agencies Consult	s Required and						
Cumulative Effect (Describe the cum considered, includ present and know regardless of who actions)	nulative impacts ling past, n future actions						
L. Mitigation (Record actions to minimize, and cor	npensate)		_				
M. Preferred Alternative	√ preferred alternative						
	Supporting reason						
N Context (Re	ecord context	of alternatives analysis)					
The significance	e of an action	must be analyzed in several co	ontexts	such as society as a whole (hu	man, n	national), the affected region, the	9
affected interes		cality. cance or Extraordinary Circu	4.714				
Intensity: Refe agency believes down into small If you answer A	ers to the sevents that on balar component pany of the best and signification.	erity of impact. Impacts may be nee the effect will be beneficial. parts. elow questions "yes" then co ance issues to consider and	both b Signif ntact t a site	eneficial and adverse. A signific ficance cannot be avoided by te the State Environmental Liaise specific NEPA analysis may b	rming a	iired.	
	Is the proximit critical a	referred alternative expected to ty to historic or cultural resource areas?	signifi es, par	·	tics of t	the geographic area such as ild and scenic rivers, or ecologic	·
	Does th environi	ne preferred alternative have hig ment?	ghly un	certain effects or involve unique	or unl		
	principle	e about a future consideration?			_	nt impacts or represent a decisi	
	quality o	of the human environment eithe	er indivi	idually or cumulatively over time	?	cant environment impacts to the	
	 Will the preferred alternative likely have a significant adverse effect on ANY of the special environmental concerns? Use the Evaluation Procedure Guide Sheets to assist in this determination. This includes, but is not limited to, concerns such as cultural or historical resources, endangered and threatened species, environmental justice, wetlands, floodplains, coastal zones, coral reefs, essential fish habitat, wild and scenic rivers, clean air, riparian areas, natural areas, and invasive species. 						
	environi	ment?			w or re	equirements for the protection of	the
P. To the best	of my knowl	edge, the data shown on this	form	is accurate and complete:			
		CS person (e.g. another MS TIG block to verify the information's			re to si	gn the first signature block and	then
				T1(1)			
	Signature (TSP if applicable)		Title		Date	
		iture (NRCS)		Title		Date	
-	f preferred alternative is not a federal action where NRCS has control or responsibility and this NRCS-CPA-52 is shared with someone other than the client then indicate to whom this is being provided.						

T	The following sections are to be completed by the Responsible Federal Official (RFO)				
NRCS is the RF	O if the action is lead federal agency for NRDA-funded actions planned by NRCS.				
Q. NEPA Com The preferred a	pliance Finding (check one) Ilternative:	Action required			
	1) is a federal action that has been sufficiently analyzed in an existing NEPA document to which this environmental evaluation is tiered because the expected effects are within the range of those described in the applicable NEPA document and there are no predicted significant adverse environmental effects or extraordinary circumstances.	Document in "R.1" below. No additional analysis is required.			
	2) is a federal action that has NOT been sufficiently analyzed or may involve predicted significant adverse environmental effects or extraordinary circumstances and may require an EA or EIS.	Contact the State Environmental Liaison. Further NEPA analysis required.			
	upporting the Finding				
R.1 Findings Docum	pentation				
	red the effects of the alternatives on the Resource Concerns, Economic and Social Concerns, and Extraordinary Circumstances as defined by Agency regulation and ed above.				
S. Signature o	f Responsible Federal Official:				
	Signature	Date			
	Oignature Title	Date			
	Additional notes				

Appendix B. Conservation Practices List for Nutrient Reduction Alternative A and B


Code	Practice	Alternative A	Alternative B
201	Edge of Field Water Quality Monitoring Data Collection	X	X
202	Edge of Field Water Quality Monitoring System Implementation	X	X
313	Waste Storage Facility	X	
314	Brush Management (Heavy Equipment)	X	X
315	Herbaceous Weed Control	X	X
317	Composting Facility	X	
327	Conservation Cover	X	
328	Conservation Crop Rotation	X	
329	Residue Management, No-Till	X	
338	Prescribed Burning	X	X
340	Cover Crops	X	
342	Critical Area Planting	X	X
345	Residue and Tillage Management, Reduced Till	X	
350	Sediment Basin	X	
356	Dike	X	
362	Diversion	X	
378	Pond	X	
381	Silvopasture Establishment	X	***
382	Fence	X	X
386 390	Field Border	X	X
390	Riparian Herbaceous Cover Riparian Forest Buffer	X	X
391	Filter Strip	X	Λ
393	Firebreak (New construction)	X	X
410	Grade Stabilization Structure	X	X
410	Grassed Waterways	X	Λ
422	Hedgerow Planting	X	
430	Irrigation Pipeline	X	X
441	Irrigation System, Microirrigation	X	Α
442	Irrigation System, Sprinkler	X	
443	Irrigation System, Surface and Subsurface	X	
449	Irrigation Water Management	X	
460	Land Clearing	X	
464	Irrigation Land Leveling	X	
468	Lined Waterway Or Outlet	X	
484	Mulching	X	X
490	Forest Site Preparation (Chemical or Burning)	X	X
490	Forest Site Preparation (Mechanical)	X	X
511	Forage Harvest Management	X	
512	Pasture and Hay Planting	X	
516	Pipeline	X	
528A	Prescribed Grazing	X	X
554	Drainage Water Management	X	
561	Heavy Use Area Protection	X	
576	Livestock Shelter Structure	X	37
578	Stream Crossing	X	X
580 587	Streambank and Shoreline Protection Structure For Water Control	X X	X X
590	Nutrient Management	X	Λ
595	Pest Management Pest Management	X	
600	Terrace	X	
612	Tree/Shrub Establishment (Hand Planting)	X	X
612	Tree/Shrub Establishment (Mechanical Planting)	X	X
614	Watering Facility	X	X
642	Water Well	X	23
644	Wetland Wildlife Habitat Management	X	
666	Forest Stand Improvement (Chemical/Hand Tools)	X	X
666	Forest Stand Improvement (Cutting/removal with heavy equipment)	X	X
500	12 or out a marto remone (Catalag removal with nearly equipment)	1 21	1 23


Appendix C. Exemplar Conservation Practice Network Effect Diagrams for Nutrient Reduction




NRCS CONSERVATION PRACTICE EFFECTS - NETWORK DIAGRAM

September 2014








Notes:

Effects are qualified with a plus (+) or minus (-). These symbols indicate only an increase (+) or a decrease (-) in the effect upon the resource, not whether the effect is beneficial or adverse. Projects involving long lengths of bank or shoreline, structural controls, substantial earth moving and/or fill, or sensitive waters may need to be evaluated in a site-specific EA or EIS.


Additional information about potential protection measures and their impacts is available in the EIS for the Emergency Watershed Protection (EWP) Program.


² Conventional bank armoring (e.g., rip rap, gabions) may result in decreased (-) channel/flood plain dynamics, and associated impacts, while other less intrusive methods (e.g., stream barbs, stone toes with sloped, vegetated banks) may result in increased (+) channel/flood plain dynamics.



NRCS CONSERVATION PRACTICE EFFECTS - NETWORK DIAGRAM

September 2015





APPENDIX D

Draft Monitoring and Adaptive Management Plan for *Deepwater Horizon* NRDA Project:

Graveline Bay Land Acquisition & Management

Page 2 of 23

Table of Contents

1.0	Introduction	3
1.1	Project Overview	3
1.2	Project Goals and Restoration Objectives	5
1.3	Conceptual Model	6
1.4	Sources of Critical Uncertainty	7
2.0	Project Monitoring	8
3.0	Rationale for Adaptive Management	12
4.0	Evaluation	13
5.0	Project-Level Decisions	14
6.0	Monitoring Schedule	15
7.0	Data Management	16
7.1	Data Review and Clearance	16
7.2	Data Storage and Accessibility	16
7.3	Data Sharing	17
8.0	Reporting	17
9.0	Roles and Responsibilities	17
10.0	Monitoring Budget	18
11.0	References	18
AP	PENDIX 1	20
AP	PENDIX 2	22

1.0 Introduction

Monitoring, Adaptive Management, and Administrative (MAM) Oversight was identified as one of the programmatic goals in the *Deepwater Horizon* (*DWH*) oil spill Programmatic Damage Assessment and Restoration Plan and Programmatic Environmental Impact Statement (PDARP/PEIS). The *DWH* NRDA MAM Framework provides a flexible, science-based approach to effectively and efficiently implement restoration over several decades that provides long-term benefits to the resources and services injured by the *DWH* spill. This project MAM plan identifies the monitoring needed to evaluate progress toward meeting project objectives and to support adaptive management of the restoration project. It identifies key sources of uncertainty, incorporates monitoring data and decision points that address these uncertainties, and establishes a decision-making process for making adjustments where needed.

This MAM Plan is a living document and would be updated as needed to reflect changing conditions and/or new information. For example, the plan may need to be revised if the project design changes, if initial data analysis indicates that the sampling design is inadequate, or if any uncertainties are resolved or new uncertainties are identified during project implementation and monitoring. Any significant future revisions to this document would be made publicly available through the Restoration Portal.

1.1 Project Overview

The Graveline Bay Land Acquisition and Management proposed alternative includes acquiring parcels near publicly owned lands in the Graveline Bay Coastal Preserve (CP) in Jackson County, Mississippi. Habitat management measures are also planned including chemical treatment, mechanical treatment, prescribed fire, access restriction, debris removal and road repair/removal and culvert replacement. The proposed alternative would be implemented at proposed locations in Graveline Bay (Figure 1.1-1). The project planning process has been a collaboration between the MDMR and the MS TIG. Potential acquisitions from in the proposed alternative project area include approximately 1,410 acres of habitat that could be acquired from willing sellers. Estuarine marsh, shoreline (beach) and other coastal riparian habitats are in the proposed alternative area, some of which are expected to provide foraging, loafing and nesting for bird species injured by the *DWH* oil spill. The proposed Graveline Bay Land Acquisition and Management (Alternative A-Preferred), if selected, would be implemented and the MAM plan would be implemented. For the purposes of this Draft Monitoring and Adaptive Management Plan (Draft MAM Plan), proposed Alternative A is referred to as the project.




Figure 1.1-1. Graveline Bay Land Acquisition and Management -Parcels and Habitats¹

This project is being implemented to restore for injuries to natural resources and their services injured by *DWH* oil spill. As outlined within the PDARP/PEIS, this restoration project falls under the following programmatic goal, restoration type, restoration approach, restoration technique, TIG, and restoration plan:

- Programmatic goals: Restore and Conserve Habitat; Replenish and Protect Living Coastal and Marine Resources
- Restoration types: Wetlands, Coastal and Nearshore Habitats; Birds
- Restoration approaches: Protect and Conserve Marine, Coastal, Estuarine, and Riparian Habitats; Restore and Conserve Bird Nesting and Foraging Habitat
- Restoration techniques: Acquire lands for conservation; Develop and implement management actions in conservation areas and/or restoration projects; Enhance habitat through vegetation management
- TIG: Mississippi
- Restoration plan: Mississippi Trustee Implementation Group Draft 2016-2017
 Restoration Plan/Environmental Assessment (Draft RP/EA)

¹ Parcels will be purchased at appraised value for appraisable lands.

This restoration project is being implemented in the Graveline Bay estuary (HUC 10 ,0317000907) and more specifically the private parcels adjacent to Graveline Bay and bayou in Jackson County, Mississippi. The parcels are located in Sections 4, 5, 9, 10, 15, and 16 of Township 8 South, Range 7 West.

Management activities would include access restriction, chemical treatment, mechanical treatment, prescribed fire, debris removal and road repair/removal and culvert replacement.

1.2 Project Goals and Restoration Objectives

Under the Restore and Conserve Habitat Programmatic Goal, the MS TIG would focus on the Wetlands, Coastal and Nearshore Habitats Restoration Type. Specific goals of the restoration type include:

- Restore a variety of interspersed and ecologically connected coastal habitats in each of
 the five Gulf states to maintain ecosystem diversity, with particular focus on maximizing
 ecological functions for the range of resources injured by the spill, such as oysters,
 estuarine-dependent fish species, birds, marine mammals, and nearshore benthic
 communities
- 2) Restore for injuries to habitats in the geographic areas where the injuries occurred, while considering approaches that provide resiliency and sustainability.
- 3) While acknowledging the existing distribution of habitats throughout the Gulf of Mexico, restore habitats in appropriate combinations for any given geographic area. Consider design factors, such as connectivity, size, and distance between projects, to address injuries to the associated living coastal and marine resources and restore the ecological functions provided by those habitats.

The specific restoration objectives for this project under the Wetlands, Coastal and Nearshore Restoration Type are:

- 1) Protect estuarine marsh, shoreline (beach) and other coastal riparian habitats from development and increase habitat connectivity to other large conservation parcels, by acquiring priority lands in the Graveline Bay Coastal Preserve for conservation.
- 2) Increase and maintain native vegetation species composition in restored habitats within Graveline Bay Coastal Preserve.

Under the Replenish and Protect Living Coastal and Marine Resources Programmatic Goal, the MS TIG would focus on Birds Restoration Type. Specific goals of the restoration type include:

- 1) Restore or protect habitats on which injured birds rely.
- 2) Restore injured birds by species where actions would provide the greatest benefits within geographic ranges that include the Gulf of Mexico.

The specific objectives for this project relative to the Birds Restoration Type are:

- 3) Increase and maintain shorebird (species injured by the *DWH* oil spill) use of beach habitat
- 4) Increase and maintain wading bird habitat (species injured by the *DWH* oil spill) use in acquired habitats

The following Restoration Objectives, as outlined in this Draft MAM Plan, are:

- 1) Protect estuarine marsh, shoreline (beach) and other coastal riparian habitats from development and increase habitat connectivity to other large conservation parcels, by acquiring priority lands in the Graveline Bay Coastal Preserve for conservation.
- 2) Increase and maintain native vegetation species composition in restored habitats within Graveline Bay Coastal Preserve.
- 3) Increase and maintain shorebird (species injured by the *DWH* oil spill) use of beach habitat.
- 4) Increase and maintain wading bird habitat (species injured by the *DWH* oil spill) use in acquired habitats.

Performance criteria would be used to determine restoration success or the need for corrective action in accordance with (15 CFR 990.55(b)(1)(vii) and are outlined for each objective in Section 2.

1.3 Conceptual Model

The singular purpose of conservation is to ensure the protection of habitat from development or further degradation. By placing lands under agency stewardship, it prevents development and disturbances in priority habitats that buffer protected coastal wetlands, but then allows for the restoration and enhancement of native vegetation assemblages and structure that support life cycle needs of numerous injured shorebirds and wading birds in coastal Mississippi (Table 1.3-1). The habitats in the project area include estuarine marsh, fire-suppressed pine savannas, beach-magnolia forests, coastal plain small stream riparian forest, beach and open water. Protection of these habitats within this key Gulf Coast watershed would protect downstream natural resources by slowing and filtering nutrient laden runoff, maintain resiliency of dynamic habitats by allowing for free movement in response to changing climate conditions, and provide diverse habitat to serve as refuge for wildlife in the densely populated coastal region. Habitat conservation also enhances habitat connectivity and ties into ecological paradigms of hub and corridors for species movement, habitat migration, and population source sink models. Habitat enhancement of conserved lands through various restoration measures of invasive species removal, restoring hydrological functions, returning fire to the systems increases the natural ecosystem functioning of the respective habitats, resulting in a more resilient and sustainable habitat, increased heterogeneity of habitat patches, and thus increases the diversity of the system. Table 1.3-1. Conceptual model for the project.

Activity	Output	Short-term outcome	Long-term outcome
Implement acquisition actions to inhibit development and increase habitat connectivity	Protection and conservation of priority habitats and birds in the Graveline Coastal Preserve boundary	 Increase in habitat connectivity and core areas Increase in injured bird habitat use 	 Protection of key habitats in perpetuity Enhancement of ecosystem services of Gulf coast habitats and living resources
Implement management actions on acquired parcels	 Increase natural ecosystem functioning Enhanced habitat for shorebird use 	Increase in native vegetation species composition and desired vegetation structure Increase in injured bird habitat use	 Increase in management of connected habitats Enhancement of ecosystem services of Gulf coast habitats and living resources

1.4 Sources of Critical Uncertainty

The focus of adaptive management is to learn through targeted monitoring and use information learned to make more informed decisions through time. Learning for adaptive management takes place in the form of reducing critical uncertainty. Critical uncertainties are defined as those that have the potential to impact or impede the decision-making process and the ability to achieve the restoration objective(s). Although many types of scientific and other uncertainties exist, the focus of uncertainty in an adaptive management context is the uncertainty that affects the decisions being made for a project or groups of projects. Monitoring to resolve critical uncertainties affecting these decisions can allow for more effective expenditure of resources (e.g., optimized project selection) into the future as learning takes place. Further, the learning that takes place through monitoring allows corrective actions to be taken to improve project outcomes. If unresolved, the critical uncertainty may delay the time it takes to achieve the restoration objectives, hinder an implemented project's ability to fully achieve restoration objectives, or in the worst-case scenario, it may have the potential to cause a project to fail altogether, regardless of the corrective actions taken. Based on information in the conceptual ecological model, potential critical uncertainties for the project were identified and evaluated. These critical uncertainties are shown in Table 1.4-1.

Table 1.4-1. Critical uncertainties that may affect success of the Graveline Land Acquisition and Management project.

Critical Uncertainty	Summary of Strategy to Resolve
Native vegetation communities do not regenerate after implementation of restoration/management activities.	Conduct targeted monitoring on metrics related to native plant composition and abundance specific to each habitat type (i.e., fire-suppressed pine flatwoods, etc.) and for each restoration/management action (chemical treatment, prescribed fire, mechanical treatment). Monitoring data would be used to refine future management actions.

Critical Uncertainty	Summary of Strategy to Resolve
Bird species injured by the <i>DWH</i> oil spill fail to use designated bird habitat affected by the project.	Conduct targeted monitoring on habitat metrics specific to wading bird habitat requirements. Monitoring data would be used to refine future management actions.
Targeted habitats do not become available for purchase.	Funding allocated for fee-simple acquisition would be used to implement habitat restoration activities within project boundaries.

2.0 Project Monitoring

The proposed monitoring for this restoration project was developed to evaluate project performance. The monitoring parameters, outlined below, are organized by project objective, with one or more monitoring parameters for each objective. Information is provided on the monitoring methods, timing and frequency, duration, sample size, and sites. In addition, example performance criteria for each parameter are identified (if applicable), including example corrective actions that could be taken if the performance criteria are not met. These parameters would be monitored at the restoration project. The parameters listed below may or may not be tied to performance criteria and/or corrective actions. Project monitoring would be applied to the following objectives:

Objective 1: Acquire targeted land parcels to protect and increase connectivity in coastal habitats

Objective 2: Implement management activities to help restore and manage the structure and function of native vegetation in coastal habitats.

Parameter # 1: Acreage of acquired land, by habitat type - the number of acres acquired through purchase of parcels in the project area.

- a) Rationale: Evaluate progress toward meeting objective 1.
- b) Method: This parameter would record the number and location of acres acquired through purchase of targeted parcels within the project boundaries.
- c) Timing, Frequency, and Duration: Land acquisitions would be recorded after each purchase and reported at the end of the project or at MS TIG request. Acquisition would occur over a 10-year period as parcels become available.
- d) Sample Size: N/A
- e) Sites: Parcels within and adjacent to the Graveline Coastal Preserve boundary.
- f) Performance Criteria: Fee-simple acquisition of up to 1,410 acres of target habitats within the project boundaries.
- g) Corrective Action: Purchase of lands from willing sellers would be subject to negotiations with the State of Mississippi as well as due diligence activities. If for

any reason, the State is unable to purchase the parcel, the next parcel that becomes available within the project area would be sought.

Parameter #2: Vegetation Structure

- a) Rationale: Evaluate progress toward meeting objective 2
- b) Methods:
 - 1. The project would adopt the methodologies described in the *Field Manual for Rapid Assessment Metrics for Wildlife and Biodiversity in Southern Open Pine Ecosystems* (see Nordman et al., 2016) for the habitat "Wet Longleaf & Slash Pine Flatwoods & Savannas". Assessment would consist of walking stands along established transects or visits to sets of random points within stands and documenting site characteristics (see Appendix 1). Then, metric assessment scores would be derived to calculate a score for the canopy, ground layer, and invasive species, and an overall score applied using the worksheet provided in Appendix 2.
- c) Timing, Frequency, and Duration: Habitat management would occur after lands are in the State ownership and a management plan is written. Monitoring activities can begin immediately after the parcel title is in place. Monitoring would take place twice per year (growing season and non-growing season) for the first year after treatment and once per year for the next four years in the growing season. Interannual sampling times may differ based on the timing of restoration actions. After the five-year period, the data would be analyzed and the appropriate corrective actions would be implemented to address the performance criteria.
- d) Sample Size: Vegetation structure sampling design would be determined at a later date when a more detailed assessment of the habitat unit can take place.
- e) Sites: All acres acquired
- f) Performance Criteria:
 - 1. Vegetation structure for fire-suppressed pine savanna (by year 5)
 - i. 20-65% canopy cover of longleaf or slash pine
 - ii. 40 to 100% herbaceous cover
 - iii. Invasive nonnative plant species in any stratum present but sporadic (1-5 % cover)
- g) Corrective Action: Based on the adaptive management plan, adjust management techniques as necessary to reach performance criteria goals. This may include increasing or decreasing the prescribed fire frequency, increasing amount of mechanical removal of canopy species, or an increase in herbicidal treatment for invasive species.

Parameter #3: Vegetation Composition

- a) Rationale: Evaluate progress toward meeting objective 2
- b) Methods: The project would adopt protocols outlined in Long-Term Vegetational Monitoring at the Mississippi Sandhill Crane National Wildlife Refuge, 1997 by A.F. Clewell, R.S. Beaman, and M.E. Lasley, 47 pp. For species composition, using a

- point intercept method, all vascular plants rooted within the station would be identified. For community structure, all plants touching a sampling pole to life form (graminoid, forb, woody) would be documented and the tallest plant at each sampling point measured. Vegetation cover would be derived by dividing the number of sampling points at which each life form was intercepted by the total. Species abundance would be measured in terms of species frequency as the number of sampling points along a transect at which each species was recorded.
- c) Timing, Frequency, and Duration: Habitat management would occur after lands are in the State ownership and a management plan is written. Monitoring activities can begin immediately after the parcel title is in place. Monitoring would take place twice per year (growing season and non-growing season) for the first year after treatment and once per year for the next four years in the growing season. Interannual sampling times may differ based on the timing of restoration actions. After the five-year period, the data would be analyzed and the appropriate corrective actions would be implemented to address the performance criteria.
- d) Sample Size: Vegetation composition sampling design would be determined at a later date when a more detailed assessment of the habitat unit can take place.
- e) Sites: All acres acquired
- f) Performance Criteria: 95% native flora²
- g) Corrective Action: Based on the adaptive management plan, adjust management techniques as necessary to reach performance criteria goals. This may include increasing or decreasing the prescribed fire frequency, increasing amount of mechanical removal of canopy species, or an increase in herbicidal treatment for invasive species.

Parameter #4: Invasive Species

- a) Rationale: Evaluate progress toward meeting objective 2
- b) Methods: The project would adopt protocols establish by MDMR Coastal Preserve System for invasive species assessment. Each site would undergo an initial GIS analysis that would analyze recent historical imagery (best available) and habitat areas. Historical land use in that period would be analyzed for high-risk land use changes which could introduce invasive plant species or increase their competiveness with typical native species. Example land uses would include logging, presence of roadways and other artificial edges, presence of hunting food plots and stands, and areas impacted by storm surge or wind events. This analysis would result in prioritized polygons within the subject property that would be considered as 'high risk' for the presence of invasive species. An initial site reconnaissance would be conducted where property is viewable by roads, trails, or waterways. Occurrences of invasive plant species would be noted and compared to the coverage of 'high risk'

² The performance criteria documented here represents a desired condition for the vegetation for a restored site that is well-managed through time. These conditions will be variable across the project area given uncertainties in the timing of management implementation, weather, and other factors.

- polygons. This comparison may result either in polygons being dropped or added to the original 'high risk' list. Invasive species would be comprehensively documented and the extent mapped while engaged in the vegetation structure survey.
- c) Timing, Frequency, and Duration: Habitat management would occur after lands are in the State ownership and a management plan is written. Monitoring activities can begin immediately after the parcel title is in place. Monitoring would take place twice per year (growing season and non-growing season) for the first year after treatment and once per year for the next four years in the growing season. Interannual sampling times may differ based on the timing of restoration actions. After the five-year period, the data would be analyzed and the appropriate corrective actions would be implemented to address the performance criteria.
- d) Sample Size: Areal extent of invasive species in acquired habitat
- e) Sites: All acres acquired
- f) Performance Criteria: 1-5% cover in invasive species³
- g) Corrective Action: Based on the adaptive management plan, adjust management techniques as necessary to reach performance criteria goals. This may include increasing or decreasing the prescribed fire frequency, increasing amount of mechanical removal of canopy species, or an increase in herbicidal treatment for invasive species.

Parameter # 5: Shorebird Diversity and Abundance

- a) Rationale: This parameter would be used to evaluate progress toward objective 3.
- b) Method: Survey routes would consist of established transects along stretches of shoreline/beach. A total of 20 surveys would be conducted annually, in four survey pulses. Species type and abundance would be documented.
- c) Timing, Frequency, and Duration: Four survey pulses would be conducted each year over a five-year period corresponding to fall migration, winter (overwinter), spring migration, and summer nesting as follows:
 - 1. Fall surveys occur between 20 August and 30 October.
 - 2. Winter surveys occur between 10 January and 20 February.
 - 3. Spring surveys occur between 20 March and 30 May.
 - 4. Summer surveys occur between 1 June- 31 July.
- d) Sample Size: One survey transect over 5 acres of beach front
- e) Sites: Graveline Beach (5 acres) before and after management action
- f) Performance Criteria:
 - 1. Increase shorebird habitat use by year 5
- g) Corrective Actions:
 - 1. Identify actions to benefit priority species (e.g., vegetation management, stewardship actions)
 - 2. Continue to monitor

Parameter # 6: Wading Bird Diversity and Abundance

³ The performance criteria documented here represents a desired condition for the vegetation for a restored site that is well-managed through time. These conditions will be variable across the project area given uncertainties in the timing of management implementation, weather, and other factors.

- a) Rationale: This parameter would be used to evaluate progress toward objective 4.
- b) Method: Survey routes would consist of established transects along stretches of forested riparian habitat. A total of 10 surveys would be conducted annually, in two survey pulses. Species type and abundance would be documented as well as visible nests during the nesting season.
- c) Timing, Frequency, and Duration: Two survey pulses would be conducted each year over a five-year period corresponding to spring and summer nesting. Five surveys would occur in each season.
- d) Sample Size: Survey routes would be established in ten riparian drainage locations across the project site. Each riparian area would have one transect route. Routes would differ in length from 200-500 meters
- e) Sites: Survey routes would be conducted as parcels are acquired. All routes would be located in parcels north of Graveline Bay/Bayou
- f) Performance Criteria:
 - 1. Increase wading bird habitat use by year 5
- g) Corrective Action:
 - 1. Continue to monitor
 - 2. Identify actions to benefit priority species (e.g., vegetation management, stewardship actions)

3.0 Rationale for Adaptive Management

As discussed in the PDARP/PEIS, adaptive management is a form of structured decision-making applied to the management of natural resources in the face of uncertainty (Pastorok et al. 1997; Williams 2011). It is an iterative process that integrates monitoring and evaluation of management actions with flexible decision-making, where adjustments are made to management approaches based on observed outcomes (NRC 2004). Within the context of ecological restoration, adaptive management addresses key uncertainties by linking science to restoration decision-making (Steyer & Llewellyn 2000).

Although adaptive management is a critical component of the restoration plan as a whole, the need for adaptive management may vary on a project by project basis. For example, higher uncertainty may be associated with novel approaches, larger restoration scales (e.g., number and area of projects), limited scientific understanding of target resources, increasing influence of socioeconomic factors, and longer time scales of restoration implementation (LoSchiavo et al. 2013; Simenstad et al. 2006; Steyer & Llewellyn 2000; Williams & Brown 2012; see PDARP/PEIS for more information). The OPA NRDA regulations require that all restoration projects clearly identify performance criteria that would be used to determine project success or the need for corrective action. Projects with more uncertainty may require a more active approach to adaptive management.

4.0 Evaluation

Evaluation of monitoring data is needed to assess the performance of the project in meeting its restoration objectives, resolving uncertainties to increase understanding, and determine whether corrective actions are needed.

The results of the analysis would be used to answer the following questions:

- Were the project restoration objectives achieved? If not, is there a good reason why they were not met?
- Did the restoration project produce unanticipated effects?
- Were there unanticipated events unrelated to the restoration project that potentially affected the monitoring results?
- Were any of the uncertainties identified prior to project implementation resolved?
- Were any new uncertainties identified?

The analysis methods would be applied to all monitoring parameters as follows:

Vegetation structure

Recorded metrics would be compared an annual basis using descriptive summaries to track performance across time by analyzing individual metric scores and final scores for each sampling effort. Comparisons would include canopy cover, ground layer cover, basal area, and invasive species cover (Appendix 2).

Vegetation Composition

All data would be analyzed using software capable of calculating general descriptive statistical analyses. Common analyses include:

- Descriptive summaries of cover for grass, forbs, and shrubs. Cover is calculated by dividing the number of intervals at which a life form was measured by the total number of intervals measured.
- Descriptive summaries of mean grass height, mean forb height, mean shrub height, preand post-treatment. The mean height of a life form is calculated by dividing the sum of the heights by the total number of interception points at which the life form occurred.
- Multivariate statistics (PCA/per MANOVA) can be applied to detect the degree of similarity of species abundance across space and time (Clewell, 1997).

Bird Habitat Use

All data would be analyzed using software capable of calculating general descriptive statistical analyses. Common analyses include:

- Descriptive summaries and tabulation of species richness and species abundance across seasons and years.
- Comparative statistics to determine differences in species richness and abundance before and after management action as well as comparisons with legacy data for the site.

5.0 Project-Level Decisions

The decision-making process requires a structured approach for incorporating new information gained from monitoring and evaluation. As specified in the NRDA regulations, performance criteria would be used to determine restoration success or the need for corrective action (15 CFR 990.55(b)(1)(vii)). However, unanticipated consequences, previously unknown conditions or unanticipated environmental drivers uncovered during the evaluation step may also determine the need for corrective actions. Table 5.0-1 provides the interim performance criteria for helping determine whether adjustments to the project are needed to better ensure the project meets the final performance criteria used to determine project success, as well as the potential adaptive management actions (e.g., mid-course corrections or corrective actions) that may be considered for individual parameters. This table does not include all possible options; rather, it includes a list of potential adaptive management actions for each individual parameter to be considered. The decision to implement a corrective action should holistically consider the overall outcomes of the restoration project by assessing the results of all monitoring parameters compiled in the evaluation step.

Table 5.0-1: Corrective Actions for the Graveline Bay Land Acquisition and Management

Monitoring Parameter	Final Performance Criteria used to determine Project Success (Year 10)	Interim Performance Criteria	Potential corrective actions or mid- course corrections
Acres Acquired	Fee-simple acquisition of 1,400 acres of priority habitats in the project area	Performance criteria not met by year 5	Funding allocated for fee- simple acquisition would be used to implement habitat restoration activities within project boundaries.
Acres Managed for Vegetation Structure	20-65% canopy cover of longleaf or slash pine 40 to 100% herbaceous cover Invasive nonnative plant species in any stratum present but sporadic (1-5 % cover)	Performance criteria not met by year 5	 Change burn frequency Modify mechanical removal strategy Alter herbicide treatments Continue to monitor.
Shorebird Diversity and Abundance	Increase in species diversity	Performance criteria not met for year 5	Identify actions to benefit priority species (e.g.,

Monitoring Parameter	Final Performance Criteria used to determine Project Success (Year 10)	Interim Performance Criteria	Potential corrective actions or mid- course corrections
	2) Increase in species		vegetation management,
	abundance		stewardship actions)
			2) Continue to monitor
Wading Bird	1) Increase in species	Performance criteria not met	Identify actions to benefit
Diversity and	diversity	for year 5	priority species (e.g.,
Abundance	2) Increase in species		vegetation management,
	abundance		stewardship actions)
			2) Continue to monitor

6.0 Monitoring Schedule

The schedule for the project monitoring is shown in Table 6.0-1, separated by monitoring activity. Execution monitoring occurs when project has been fully executed as planned (Year 0). The monitoring of project parameters is dependent on the voluntary participation by landowners to sell targeted parcels. Performance monitoring would occur in the years following initial project execution (Years 1-5), and is restrained by the ten-year duration of the overall project. The length of time a parameter is monitored is contingent on when the restoration action is executed within project timeline. Thus, parameters may receive monitoring for 1-5 years. For example, if a parcel is acquired in year 7 of the project, monitoring would occur for three years to coincide with the overall project timeline. The same timeline structure would apply to the monitoring of managed habitats. The monitoring schedule would be updated as acquisitions are finalized and management actions implemented.

Table 6.0-1. Monitoring Schedule.

Monitoring Parameters	Monitoring Times	rame ⁴				
	Execution Monitoring (initial)	Post-Execution Monitoring (ongoing)				
	As-built (Year 0)	Year 1	Year 2	Year 3	Year 4	Year 5
Parameter 1		x	X	X	X	X
Parameter 2		X	X	X	X	X
Parameter 3		X	X	X	X	X
Parameter 4		X	X	X	X	X
Parameter 5		X	X	X	X	X
Parameter 6		X	X	X	X	X

7.0 Data Management

To the extent practicable, all environmental and biological data generated during monitoring activities would be documented using standardized field datasheets. If standardized datasheets are unavailable or not readily amendable to record project-specific data, then project-specific datasheets would be drafted prior to conducting any project monitoring activities. All tangible forms of field data would be reviewed by Implementing Trustee for completeness and accuracy before being finalized. Original hardcopy datasheets and notebooks and photographs would be retained by the Implementing Trustee.

All field datasheets and notebook entries would be scanned to PDF files and would be archived along with the hardcopy datasheets. Electronic data files should be named with the date on which the file was created and should include a ReadMe file that describes when the file was created and by whom, and any explanatory notes on the file contents. If a data file is revised, a new copy should be made and the original preserved.

Relevant Project data that are handwritten on hardcopy datasheets or notebooks would be transcribed (entered) into Excel spreadsheets (or similar digital format). After transcription of the data, a second person not associated with data transcription would perform a verification of the data in the electronic data sheets against the original hardcopy datasheets and/or notebooks, and would make any corrections to transcription errors as appropriate before data are used for any analyses or distributed outside of the agency. Implementing Trustees would verify and validate monitoring data and information and would ensure that all data is entered or converted into agreed upon/commonly used digital format labeled with metadata.

7.1 Data Review and Clearance

Once data is entered electronically it is reviewed and verified for completeness. A quality check is made by verbally comparing the electronic data entered to the original hard copy data sheet. Data are validated and any corrections needed are made. Upon validation, data are approved for analysis, reporting and archiving. All data are kept in one permanent electronic folder as a permanent record.

After any and all identified errors are addressed, data are considered to be QA/QC'd.

The Implementing Trustee would give the other TIG members time to review the data before making such information publicly available. Before submitting the monitoring data and information package, Implementing Trustees shall confirm with one another that the package is approved for submission. No data release can occur if it is contrary to federal or state laws.

7.2 Data Storage and Accessibility

Once all data has been verified by quality assurance/quality control procedures, it would be submitted to the Restoration Project Database that is maintained by MDEQ.

7.3 Data Sharing

Data would be made publicly available, in accordance with the Open Data Policy, through the DIVER Explorer Interface within a year of when the data collection occurred.

8.0 Reporting

All reporting would occur after field reconnaissance is complete for each assessment effort. This report would summarize the findings for the sampling period including all worksheets transferred into digital format and presented in tabular and graphical formats. The data should be summarized in such a way that it is meaningful to the reader. Additionally, an annual report would be completed that includes:

- Summary data –synthesized data for all efforts during the year
- Graphs vegetation characteristics, acres managed, bird species diversity and abundance, etc.
- Interpretation of graphical data
- Discuss comparison of data if pretreatment and post treatment data are available
- Explanation of results
- Uncertainties with management actions
- Potential data collection issues
- Issues to be resolved
 - Issues to improve data collection or cooperation in getting quality data
 - Issues associated with data loss or inability to collect data for a time period (due

9.0 Roles and Responsibilities

The Trustee Council would encourage and facilitate consistency in monitoring and data procedures to evaluate and report on progress toward meeting the ecosystem goals that are the stated basis for the Restoration Agreement and are foundational to the PDARP/PEIS. The Mississippi TIG is responsible for addressing MAM objectives that pertain to their restoration activities and for communicating information to the Trustee Council or Cross-TIG MAM work group.

At the project level, MDEQ would be the implementing agency with co-lead from the Mississippi Department of Marine Resources (MDMR). MDEQ's primary roles includes coordination with project partners and the MS TIG to track project progress, program management and oversight, lead acquisition of parcels and co-lead for management operations. MDMR would be the resource management agency and support all land management aspects of the project

10.0 Monitoring Budget

The overall budget for project monitoring and adaptive management is anticipated to be approximately 7-12% of the total project budget. This budget range is considered to be in draft form and is subject to change as project planning and implementation progress.

11.0References

Clewell, A.F., B.S. Reed, M.E. Lasley. (1997). Long-term vegetational monitoring at the Mississippi Sandhill Crane National Wildlife Refuge, 1997. 35pp. Submitted to: Mississippi Sandhill Crane National Wildlife Refuge, Gautier, MS

LoSchiavo, A. J., Best, R. G., Burns, R. E., Gray, S., Harwell, M. C., Hines, E. B., and Vearil, J. W. (2013). Lessons learned from the first decade of adaptive management in comprehensive Everglades restoration. Ecology and Society, 18(4), 70.

Nordman, Carl, Rickie White, Randy Wilson, Clay Ware, Catherine Rideout, Milo Pyne, and Chuck Hunter. (2016). Rapid Assessment Metrics to Enhance Wildlife Habitat and Biodiversity within Southern Open Pine Ecosystems, Version 1.0. U.S. Fish and Wildlife Service and NatureServe, for the Gulf Coastal Plains and Ozarks Landscape Conservation Cooperative.

National Research Council. 2004. Adaptive Management for Water Resources Project Planning. National Academies Press, Washington, DC. 138pp.

Pastorok, R. A., MacDonald, A., Sampson, J. R., Wilber, P., Yozzo, D. J., & Titre, J. P. (1997). An ecological decision framework for environmental restoration projects. Ecological Engineering, 9(1), 89-107.

Simenstad, C., Reed, D., & Ford, M. (2006). When is restoration not?: Incorporating landscape-scale processes to restore self-sustaining ecosystems in coastal wetland restoration. Ecological Engineering, 26(1), 27-39.

Steyer, G. D., & Llewellyn, D. W. (2000). Coastal Wetlands Planning, Protection, and Restoration Act: A programmatic application of adaptive management. Ecological Engineering, 15(3), 385-395.

Wouldiams, B. K. (2011). Adaptive management of natural resources—framework and issues. Journal of Environmental Management, 92(5), 1346-1353.

Page 19 of 23

Wouldiams, B. K., & Brown, E. D. (2012). Adaptive management: the US Department of the Interior applications guide. US Department of the Interior, Adaptive Management Working Group. 136pp.

APPENDIX 1

Field Form for Rapid Assessment Metrics for Wildlife and Biodiversity in Southern Open Pine Ecosystems

Date:	Project:	Site ID:
Field Crow Toom Mombores		
Field Crew Team Members:		
Leader:		
Assistants:		
	Photos of Site:	AA Centrum out: _N _E_ S _W ;Buffer in:
_N _E _S _W; Add'l: Y / N		
Photo filenames:		
_		
Assessment Area Shape: Circle, Rectang	gle, Square, Polygon	Bearing:
Assessment Area Dimensions: radius 18	m, 40m, m/ft. o	r rectangle m/ft wide xm/ft long
(fill in values, units)		
State:County:	Twp:Range:	Section:USGS 7.5'
Quad:		
Landowner/Managed Area Name:		Contact Person:
	<u> </u>	
Stand Name: Permit Re	equired? Locked G	ate? Access Difficulties?
(describe)		
SITE DESCRIPTION:		

Notes:


GENERAL					
DRAWING (Optional):					
Provide a drawing of the assessment area, including					
its boundaries, either aerial					
view or transect view.					
LOCATION: Assessmen	t Area CENTRUM (che	ck one) ORIGINAL _	MOVED (why? I	now far?)	\neg
CDC II 4	GPS Filename: Projection:				
GPS Unit:	GPS Filename:		Projection:		
UTM Zone:	Datum: NAD83 WGS84	GPS Accuracy: m/	Projection: PDOP:	# of Sat's:	
	Datum: NAD83	GPS Accuracy: m/ ft Original (GRTS):		# of Sat's: Post-processed:	
UTM Zone:	Datum: NAD83 WGS84	ft	PDOP:		
UTM Zone: UTM X Easting: UTM Y Northing:	Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree select appropriate Souping:	ft Original (GRTS): uthern Open Pine Metri	PDOP: Field: cs Datasheet for p	Post-processed:	
UTM X Easting: —————— UTM Y Northing: ————— Classification (use to field form) Southern Open Pine Grown Other Community Classi	Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree select appropriate Souping:	ft Original (GRTS): uthern Open Pine Metri	PDOP: Field: cs Datasheet for p	Post-processed:	
UTM X Easting: UTM X Easting: UTM Y Northing: Classification (use to field form) Southern Open Pine Ground Community Classification	Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree select appropriate Souping: fication Reference:	ft Original (GRTS): uthern Open Pine Metri	PDOP: Field: cs Datasheet for p	Post-processed:	

APPENDIX 2

Wet Longleaf & Slash Pine Flatwoods & Savannas Metrics Data Sheet					Recorded Measured Value of Metric	Recorded Metric Score (1.0-4.0)
Canopy Metrics						
	Excellent = 4.0	Good = 3.0	Fair = 2.0	Poor = 1.0		
Canopy Southern Yellow Pine Basal Area	20-80 ft²/acre basal area of longleaf or slash pine	10 to <20 or >80 to <90 ft²/acre basal area of longleaf or slash pine	5 to <10 or 90 to <100 ft²/acre basal area of longleaf or slash pine	<5 or ≥100 ft²/acre basal area of longleaf or slash pine	ft²/acre BA	
						x0.25
Southern Yellow Pine Canopy Cover	20-65% canopy cover of longleaf or slash pine	15 to <20% canopy cover or >65-75% canopy cover of longleaf or slash pine	10 to <15% canopy cover or >75-85% canopy cover of longleaf or slash pine	<10% cover or >85% cover of longleaf or slash pine	% cover	
						x0.25
Southern Yellow Pine Stand Age Structure	BA ≥20 ft²/acre of flat- top longleaf or slash pine of any diameter and/or longleaf or slash pine trees ≥14"	BA ≥10 ft²/acre of longleaf or slash pine trees ≥14" DBH class	Longleaf or slash pine trees ≥14" DBH class present, but at <10 ft²/acre BA	No longleaf or slash pine trees ≥14" DBH or with flat-top slash or longleaf pine	ft²/acre BA	
	DBH class					x0.25
		Ground Layer Me	trics			
	Excellent = 4.0	Good = 3.0	Fair = 2.0	Poor = 1.0		

Page 23 of 23

Overall Native	40-100%	30 to <40%	20 to <30%	<20% herbaceous cover	% cover	
Herbaceous	herbaceous	herbaceous cover	herbaceous cover			
Ground Cover	cover					
						x0.25
Invasive Plant	Invasive	Invasive nonnative	Invasive nonnative	Invasive nonnative	% cover	
Presence /	nonnative plant	plant species in any	plant species in any	plant species in any		
Distribution	species absent	stratum present but	stratum uncommon (5-	stratum common		
	or cover is	sporadic (1-5%	10% cover)	(>10% cover)		
	very low (≤1%	cover)				
	cover)					
						-0.25
						x0.25
	Final Score is :					
Canony Score	Canopy Score x0.33 + Midstory Score x0.33 + Ground Layer Score x0.33 =					e =
canopy score		astory seoreA	nee : Ground Edyer Seo	A0100 =	Score	-
Evaluation	Scale: 4.0 to 3.5 =	Excellent, 3.5 to $2.5 = 0$	Good, 2.5 to 1.5 = Fair, 1.5	5 to 1.0 = Poor		
_ , , , , , , , , , , , , , , , , , , ,						



APPENDIX E

Monitoring and Adaptive Management Plan for Deepwater Horizon NRDA Project:

Grand Bay Land Acquisition and Habitat Management

Table	of Contents Introduction	3
1.1	Project Overview	3
1.2	Project Goals and Restoration Objectives	4
1.3	Conceptual Model	5
1.4	Sources of Critical Uncertainty	6
2.0	Project Monitoring	6
3.0	Rationale for Adaptive Management	10
4.0	Evaluation	10
5.0	Project-Level Decisions	12
6.0	Monitoring Schedule	13
7.0	Data Management	14
7.1	Data Review and Clearance	15
7.2	Data Storage and Accessibility	15
7.3	Data Sharing	15
8.0	Reporting	15
9.0	Roles and Responsibilities	16
10.0	Monitoring Budget	16
11.0	References	16
1.1	APPENDIX 1	18
1.2	APPENDIX 2	20

1.0 Introduction

Monitoring, Adaptive Management, and Administrative Oversight was identified as one of the programmatic goals in the *Deepwater Horizon* oil spill Programmatic Damage Assessment and Restoration Plan and Programmatic Environmental Impact Statement (PDARP/PEIS). The *Deepwater Horizon* NRDA Monitoring and Adaptive Management (MAM) Framework provides a flexible, science-based approach to effectively and efficiently implement restoration over several decades that provides long-term benefits to the resources and services injured by the *DWH* spill. This project MAM plan identifies the monitoring needed to evaluate progress toward meeting project objectives and to support adaptive management of the restoration project. It identifies key sources of uncertainty, incorporates monitoring data and decision points that address these uncertainties, and establishes a decision-making process for making adjustments where needed.

This plan would be implemented if Alternative D is selected. The proposed alternative for the purpose of this MAM Plan is referred to as the project. This MAM plan is a living document and would be updated as needed to reflect changing conditions and/or new information. For example, the plan may need to be revised if the project design changes, if initial data analysis indicates that the sampling design is inadequate, or if any uncertainties are resolved or new uncertainties are identified during project implementation and monitoring. Any future revisions to this document would be made publicly available through the Restoration Portal.

1.1 Project Overview

This project is being implemented as restoration for the *Deepwater Horizon* oil spill Natural Resource Damage Assessment (NRDA). As outlined within the *Deepwater Horizon* oil spill PDARP/PEIS, this restoration project falls under the following programmatic goal, restoration type, restoration approach, restoration technique, TIG, and restoration plan:

- Programmatic goal: Restore and Conserve Habitats; Replenish and Protect Living Coastal and Marine Resources.
- Restoration type: Wetland, coastal, and nearshore habitats; Birds
- Restoration approaches: Protect and conserve marine, coastal, estuarine, and riparian habitats; Restore and conserve bird nesting and foraging habitat
- Restoration technique: Acquire targeted lands to protect, restore, and manage coastal habitats; Implement management/restoration activities to help restore the natural function and vegetative structure of coastal habitats.
- TIG: Mississippi
- Restoration plan: Mississippi Trustee Implementation Group Draft 2016-2017 Restoration Plan/Environmental Assessment (Draft RP/EA)

This restoration project is being implemented within the approved acquisition boundary of Grand Bay National Wildlife Refuge (Refuge), Grand Bay National Estuarine Research Reserve (NERR), and the Grand Bay Savanna Coastal Preserve (Coastal Preserve). The project area is located in coastal southeast Mississippi, bordering Grand Bay, and between the municipalities of Grand Bay and Moss Point. Portions of the boundaries of the refuge, NERR and coastal preserve overlap (Figure 1.1-1). Restoration activities involve the acquisition of private parcel inholdings

and restoration of habitats, where applicable. This project is intended to help restore habitats and resources injured from the DWH oil spill, including coastal, estuarine, and riparian habitats; and birds. Implementing Trustees include the Department of the Interior (DOI) and Mississippi Department of Environmental Quality. Implementing Trustees would be working with U.S. Fish and Wildlife Service, the Mississippi Department of Marine Resources as the state agency managing the Grand Bay NERR and the Mississippi Coastal Preserve Program and NOAA as the joint managers of the Grand Bay NERR.




Figure 1.1-1. The Grand Bay Land Acquisition and Management project area.

1.2 Project Goals and Restoration Objectives

The overall goal of this restoration project is to protect, restore and manage habitat within the project boundaries to maximize native vegetative communities. These actions would help restore, replace, or acquire the equivalent of wetland, coastal, and nearshore habitats in Mississippi injured by the Deepwater Horizon spill, and provide services to bird species injured by the spill. The proposal includes two restoration objectives: habitat acquisition and habitat management.

- (1) Habitat acquisition to prevent the potential for habitat loss caused by conversion for development and to increase connectivity in native coastal habitats. The primary objective of habitat acquisition is to acquire adequate land to provide contiguous lands and waters within the project boundary in an effort to protect desired habitat and to increase connectivity in native coastal habitats. Acquiring target habitats also facilitates more efficient and effective restoration and management by leading to larger blocks of contiguous habitat which can be managed and protected as a whole. The project objective is to acquire up to 8,000 additional acres of target habitats including coastal marsh, savanna and flatwoods, forested freshwater scrub-shrub, and freshwater marsh in Grand Bay within a 15-year period.
- (2) Habitat management to restore the structure and function of target habitats within the project boundary. The primary objective of habitat restoration is to restore the structure and function of native vegetation in up to 17,000 acres of target habitats, including coastal marsh, savannas and flatwoods, forested freshwater scrub-shrub, and freshwater marsh in Grand Bay within a 15-year period.

Restoration activities would follow those currently being implemented by the Refuge, NERR and Coastal Preserve. Habitat management and restoration activities include, but are not limited to, invasive species mapping and treatment (i.e., mechanical treatment, prescribed fire, and chemical treatment), prescribed burning, and mechanical thinning to remove woody vegetation. Fire management and mechanical thinning would serve to replicate the natural ecological processes that historically shaped these coastal ecosystems and would help restore the natural function of

each habitat type, assist in providing habitat interconnectivity, and help support the natural expected processes in these habitats (e.g. inland migration of coastal marsh caused by expected sea level rise).

Performance criteria would be used to determine restoration success or the need for corrective action in accordance with 15 CFR 990.55(b) (1) (vii)). Specific, measurable performance criteria are defined for monitoring parameters associated with each of the restoration objectives (see Section 2.0).

1.3 Conceptual Model

Acquisition and management of lands in the proposed project area would protect and enhance ecosystem services of habitats and living resources. The singular purpose of conservation is to ensure the protection of habitat from development or further degradation. By placing lands under agency stewardship, it prevents development and disturbances in priority habitats but then allows for the restoration and enhancement of native vegetation assemblages and structure that support coastal, wetland and nearshore habitats in Mississippi, and life cycle needs of birds injured by the spill. The habitats within the project boundary include coastal marsh, savannas and flatwoods, freshwater marsh, and forested freshwater scrub-shrub, among others. Protection of these habitats within this key Gulf Coast watershed would protect downstream natural resources by slowing and filtering nutrient laden runoff, maintain resiliency of dynamic habitats by allowing for free movement in response to changing climate conditions, and provide diverse habitat to serve as refuge for wildlife in the densely populated coastal region. Habitat conservation also enhances habitat connectivity and ties into ecological paradigms of hub and corridors for species movement, habitat migration, and population source sink models. Habitat enhancement of conserved lands through various restoration measures of invasive species removal, restoring hydrological functions (though not contemplated in this plan), and returning fire to the system increases the natural ecosystem functioning of the respective habitats, resulting in a more resilient and sustainable habitat, increased heterogeneity of habitat patches, and thus increases the diversity of the system.

Table 1.3-1. Conceptual ecological model.

	into the treatment econogram models								
Activity	Output	Short-term outcome	Long-term outcome						
Implement acquisition actions to inhibit development and increase habitat connectivity	Protection and conservation of priority habitats and birds within the project boundary	 Maintain or increase in habitat connectivity and core areas Maintain or increase in habitat use by injured bird species bird 	 Protection of key habitats in perpetuity Enhancement of ecosystem services of Gulf coast habitats and living resources 						
Implement management actions on acquired and existing publicly- owned parcels	Increase natural ecosystem functions	Increase in native vegetation species composition and desired vegetation structure Increase in habitat use by injured bird species	 Increase in management of connected habitats Enhancement of ecosystem services of Gulf coast habitats and living resources 						

1.4 Sources of Critical Uncertainty

The focus of adaptive management is to learn through targeted monitoring and use information learned to make more informed decisions through time. Learning for adaptive management takes place in the form of reducing critical uncertainty. Critical uncertainties are defined as those that have the potential to impact or impede the decision-making process and the ability to achieve the restoration objective(s). Although many types of scientific and other uncertainties exist, the focus of uncertainty in an adaptive management context is the uncertainty that affects the decisions being made for a project or groups of projects. Monitoring to resolve critical uncertainties affecting these decisions can allow for more effective expenditure of resources (e.g., optimized project selection) into the future as learning takes place. Further, the learning that takes place through monitoring allows corrective actions to be taken to improve project outcomes. If unresolved, the critical uncertainty may delay the time it takes to achieve the restoration objectives, hinder an implemented project's ability to fully achieve restoration objectives, or in the worst-case scenario, it may have the potential to cause a project to fail altogether, regardless of the corrective actions taken.

Based on information in the conceptual ecological model, potential critical uncertainties for the project were identified and evaluated. These critical uncertainties are shown in Table 1.4-2.

Table 1.4-2. Critical uncertainties that may affect success of the Grand Bay Land Acquisition and Habitat Restoration project.

Critical Uncertainty	Summary of Strategy to Resolve
Native vegetation communities do not regenerate after implementation of restoration/management activities.	Conduct targeted monitoring on metrics related to native plant composition and abundance specific to each habitat type (i.e., open pine savanna, forested freshwater scrub-shrub, etc.) and for each restoration/management action (chemical treatment, prescribed fire, mechanical treatment). Monitoring data would be used to refine future management actions.
Injured bird species fail to use designated bird habitat effected by the project.	Consider expanding survey area to document regional presence of survey bird species (are they in the area?). Conduct targeted monitoring on habitat metrics specific to wading bird habitat requirements. Monitoring data would be used to determine the need to implement restoration activities (i.e., prescribed fire, mechanical treatment) and/or if additional wading bird habitat should be acquired.
Targeted habitats do not become available for purchase.	Funding allocated for fee-simple acquisition would be used to implement habitat restoration activities within project boundaries

2.0 Project Monitoring

The proposed monitoring for this restoration project was developed to evaluate project performance and the need for corrective actions. Information is provided on the intended purpose of each monitoring parameter (e.g., monitor progress toward meeting one of the restoration objectives, regulatory compliance, and support adaptive management of the project), monitoring

methods, timing and frequency, duration, sample size, and sites. In addition, performance criteria are defined for each performance monitoring parameter and potential corrective actions that could be taken if the performance criteria are not met.

These parameters would be monitored at the restoration project site and may also be monitored at appropriate reference and/or control sites to demonstrate how the restoration project is trending toward the performance criteria. The parameters listed below may or may not be tied to performance criteria and/or corrective actions.

Objective 1: Acquire targeted land parcels to protect and increase connectivity in coastal habitats

Objective 2: Implement management activities to help restore and manage the structure and function of native vegetation in coastal habitats.

Parameter #1: Acreage of acquired land, by habitat type - the number of acres acquired through purchase of parcels in the project area.

- a) Rationale: Evaluate progress toward meeting objective 1.
- b) Method: This parameter would record the number and location of acres acquired through purchase of targeted parcels within the project boundaries.
- c) Timing, Frequency, and Duration: Land acquisitions would be recorded after each purchase and reported at the end of the project or at MS TIG request. Acquisition would occur over a 15-year period as parcels become available.
- d) Sample Size: N/A
- e) Sites: Acquired parcels
- f) Performance Criteria: Fee-simple acquisition of up to 8,000 acres of target habitats within the project boundaries.
- g) Corrective Action: Purchase of lands from willing sellers would be subject to negotiations. If, for any reasons, the Implementing Trustees are unable to purchase the parcel, the next available parcel within the project area would be sought and/or funding allocated for fee-simple acquisition could be used to implement habitat restoration activities within project boundaries.

Parameter #2: Presence, relative abundance, status, and distribution of invasive species within the 17,000 acres of target habitats.

- a) Rationale: At this time the extent of invasive species within some of the proposed project boundary is unknown. The intent of this activity is to acquire base-line data on the presence, relative abundance, status and distribution of invasive species in order to effectively develop management strategies focused on removing and/or suppressing infestations.
- b) Method: Base-line survey of invasive species within the project boundary would be conducted using aircraft-based digital photography in conjunction with the collection of ancillary field data (i.e., ground-truthing) to identify and map locations of invasive weeds. Areas of closed canopy would be considered for additional ground truthing data collection. Using GPS, polygons would be delineated around patches of invasive weeds and any co-occurring vegetation. Data would be entered into a geographic

- information system (GIS) for weed management planning purposes. All co-occurring vegetation within the delineated polygon would be identified and recorded following protocols outlined in *National Vegetation Classification Standard*, *Version 2* (FGDC Document number FGDC-STD-005-2008).
- c) Timing and Frequency: Surveys can occur for target habitats in the project boundary prior to management activities being initiated and then again at the end of the project.
- d) Sample Size: 2
- e) Sites: Project boundary
- f) Performance Criteria: N/A
- g) Corrective Action: N/A. Data would be used for reporting purposes.

Parameter #3: Vegetation structure

- a) Rationale: These metrics would measure project success toward increasing native species composition and desired vegetation structure of restored open pine savanna habitat.
- b) Method: The project would adopt the methodologies described in the *Field Manual* for Rapid Assessment Metrics for Wildlife and Biodiversity in Southern Open Pine Ecosystems (Nordman et al., 2016) for the habitat "Wet Longleaf & Slash Pine Flatwoods & Savannas". Assessment would consist of walking stands along established transects or visits to sets of random points within stands and documenting site characteristics (see Appendix 1). Metric assessment scores would be derived to calculate a score for the canopy, ground layer, and invasive species, and an overall score applied using the metrics provided in Appendix 2 and compared to performance criteria described below.
- c) Timing, Frequency, and Duration: Monitoring would be conducted twice per year (growing season and non-growing season) for the first year after treatment and then on an annual basis during the growing season. Inter-annual sampling times may differ based on the timing of restoration actions.
- d) Sample Size: Per survey protocols
- e) Sites: Targeted and/or acquired pine savanna habitats
- f) Performance Criteria:
 - i. <20% canopy cover of longleaf or slash pine
 - ii. 40-100% herbaceous cover
 - iii. Invasive non-native plant species in any stratum present but sporadic (1-5% cover)
- g) Corrective Action: Refine or adjust management techniques as necessary to reach performance criteria goals. This may include increasing or decreasing prescribed fire frequency, increasing amount of mechanical removal of canopy species, or an increase in herbicidal treatment.

Parameter #4: Vegetation composition

- a) Rationale: These metrics would measure project success toward increasing native species composition of restored open pine savanna habitat.
- b) Method: The project would adopt protocols outlined in Long-Term Vegetational Monitoring at the Mississippi Sandhill Crane National Wildlife Refuge, 1997 (Clewell et al., 1998) and Initial Survey Instructions: Long-term Vegetation

Monitoring-Life Form (Clewell Plots) (Wilder, 2016). Four long-term monitoring plots would be established within the project boundaries. Plots would consist of two parallel 200-ft transects spaced 100 ft. apart. Both species composition and community structure surveys would use the point intercept method at 2-foot intervals along each transect (n=200). Species abundance surveys would identify and record all vascular plants rooted within the plot. Community structure surveys would document and record the presence and maximum intercept height for each life form encountered (i.e., grasses, forbs, shrubs, and other). Vegetation cover would be derived by dividing the number of sampling points at which each life form was intercepted by the total. Species abundance would be measured in terms of species frequency as the number of sampling points along the transect at which each species was recorded.

- c) Timing, Frequency, and Duration: Long-term monitoring would rely on both annual surveys documenting changes in the abundance of vegetation life forms (grasses, forbs, and shrubs) and periodic surveys (once within one year after prescribed burns) of plant species composition over time.
- d) Sample Size: 4 long-term monitoring plots (two baseline and two treated)
- e) Sites: Baseline and treated habitats
- f) Performance Criteria: 95% native flora
- g) Corrective Action: Refine or adjust management techniques as necessary to reach performance criteria goals. This may include increasing or decreasing prescribed fire frequency, increasing amount of mechanical treatment or removal of canopy species, or an increase in chemical treatment/herbicidal treatment.

Parameter #5: Presence/absence of wintering Henslow's sparrow (Ammodramus henslowii)

- a) Rationale: Henslow's sparrow are an indicator species of high quality open pine savanna habitat. This metric would measure project success towards the restoration of open pine savanna habitat.
- b) Method: The project would adopt protocols outlined in *Project Prairie Birds: A Citizen Science Project for Wintering Grassland Birds* (Shackelford et al., 2001). In short, survey crews of three would include two outside individuals each using bamboo cane poles to beat the vegetation to flush skulking birds. The center person starting at the transect start point and between the pole operators, would aim for the end marker and commence walking while maintaining pole operators' rhythm and position. The center person would monitor the entire transect for birds as the flush in from of the survey line. All three individuals would spot birds and maintain a straight survey line approximately 20 m wide while walking 100 m.
- c) Timing, Frequency, and Duration: Surveys would be conducted a minimum of three times per winter season at specific intervals. Surveys would take approximately 90 to 120 seconds per transect.
- d) Sample Size: The number of transects would be dependent upon the size of the grassland site. Transects would be 100 m long and approximately 20 m wide.
- e) Sites: Pine savanna
- f) Performance Criteria: Presence/absence of wintering Henslow's sparrow.
- g) Corrective Action: Refine or adjust restoration management activities. This may include increasing or decreasing prescribed fire frequency, increasing amount of mechanical removal of canopy species, or an increase in herbicidal treatment.

Parameter #7: Diversity and abundance of injured bird species in targeted forested freshwater scrub-shrub habitats.

- a) Rationale: This metric would measure injured bird species use of bottomland hardwood habitats within the project boundary. Acquisition and protection of this targeted habitat would potentially benefit "wading bird species" with quantified injuries identified in the PDARP.
- b) Method: Walking or boating surveys would be conducted along transects within bottomland hardwood habitats. Data collection would include injured bird species identification, species abundance, and location.
- c) Timing, Frequency, and Duration: Surveys would occur annually in the spring (March 20 through May 30) and fall (August 20 through October 30).
- d) Sample Size: 2 per site/year
- e) Sites: Targeted and/or acquired bottomland hardwood habitat
- f) Performance Criteria: Use of this habitat by injured bird species.
- g) Corrective Action: Consider expanding survey area to document regional presence of survey bird species (are they in the area?). Conduct targeted monitoring on habitat metrics specific to wading bird habitat requirements. Monitoring data would be used to refine future management actions.

3.0 Rationale for Adaptive Management

As discussed in the PDARP/PEIS, adaptive management is a form of structured decision-making applied to the management of natural resources in the face of uncertainty (Pastorok et al. 1997; Williams 2011). It is an iterative process that integrates monitoring and evaluation of management actions with flexible decision-making, where adjustments are made to management approaches based on observed outcomes (NRC 2004). Within the context of ecological restoration, adaptive management addresses key uncertainties by linking science to restoration decision-making (Steyer & Llewellyn 2000).

Although adaptive management is a critical component of the restoration plan as a whole, the need for adaptive management may vary on a project by project basis. For example, higher uncertainty may be associated with novel approaches, larger restoration scales (e.g., number and area of projects), limited scientific understanding of target resources, increasing influence of socioeconomic factors, and longer time scales of restoration implementation (LoSchiavo et al. 2013; Simenstad et al. 2006; Steyer & Llewellyn 2000; Williams & Brown 2012; see PDARP/PEIS for more information). Under OPA NRDA regulations, restoration projects clearly identify performance criteria that would be used to determine project success or the need for corrective action. At a minimum, all project MAM plans should include identification of potential corrective actions. Projects with more uncertainty may require a more active approach to adaptive management.

4.0 Evaluation

Evaluation of monitoring data is needed to assess the performance of the project in meeting its restoration objectives, resolving uncertainties to increase understanding, and determine whether corrective actions are needed.

As part of the larger decision-making context beyond the project scale, the evaluation of monitoring data from the individual projects would be compiled and assessed at the Restoration Type and TIG level, and the results would be used to update the knowledge base to inform decisions such as future TIG project prioritization and selection, implementation techniques, and the identification of critical uncertainties.

The results of the analysis would be used to answer the following questions:

- Were the project restoration objectives achieved? If not, is there a reason why they were not met?
- Did the restoration project produce unanticipated effects?
- Were there unanticipated events unrelated to the restoration project that potentially affected the monitoring results (e.g., hurricanes)?
- Were any of the uncertainties identified prior to project implementation resolved?
- Were any new uncertainties identified?

Analysis Methods:

Vegetation structure

Recorded metrics would be compared an annual basis using descriptive summaries to track performance across time by analyzing individual metric scores and final scores for each sampling effort. Comparisons would include canopy cover, ground layer cover, basal area, and invasive species cover.

Vegetation Composition

Data would be analyzed using software capable of calculating general descriptive statistical analyses. Common analyses include:

- Descriptive summaries of cover for grass, forbs, and shrubs. Cover is calculated by dividing the number of intervals at which a life form was measured by the total number of intervals measured.
- Descriptive summaries of mean grass height, mean forb height, mean shrub height, preand post-treatment. The mean height of a life form is calculated by dividing the sum of the heights by the total number of interception points at which the life form occurred.
- Multivariate statistics (PCA/perMANOVA) can be applied to detect the degree of similarity of species abundance across space and time (Clewell, 1997).

Injured Bird Species Diversity and Abundance

Data would be analyzed using appropriate software capable of calculating general descriptive statistics.

- Descriptive summaries of bird species abundance (total number of individuals per species per survey) and
- Species diversity (total number of species per survey).

5.0 Project-Level Decisions

An adaptive approach to decision making involves exploring different ways (i.e., alternatives) to meet restoration objectives, predicting the outcomes of those alternatives based on the current state of knowledge, implementing one or more of these alternatives, monitoring to learn about the impacts of alternatives, and then using the results to update knowledge and improve future decisions (DOI Tech Guide). In this section, we describe how updated knowledge gained from the evaluation of monitoring data would be used at the project scale to determine whether the project, once implemented, is considered successful or whether the project requires corrective actions. A project may not be achieving its intended objectives because of previously identified critical uncertainties, unanticipated consequences, previously unknown conditions, or unanticipated environmental drivers. The decision to implement (or not implement) corrective actions is one type of decision within the larger adaptive management decision-making framework.

Learning through monitoring allows for informed corrective actions to be made to the project to achieve desired outcomes. This table identifies corrective actions for each performance criteria (as defined in NRDA regulations (15 CFR 990.55(b) (1) (vii)) but may not include all possible options; rather, it includes a list of potential actions for each individual parameter to be considered if the project is not performing as expected once implemented. Other corrective actions may be identified post-implementation, as appropriate. The decision of whether or not a corrective action should be implemented for a project should holistically consider the overall outcomes of the restoration project (i.e. looking at the combined evaluation of multiple performance criteria) in order to understand why project performance deviates from the predicted or anticipated outcome. The decision to implement a corrective action and the knowledge gained from the process could also inform the larger decision making framework, such as whether prioritization of the restoration technique should change or how to implement the restoration technique to improve the likelihood of achieving favorable project outcomes in future applications.

Table 5.0-1. List of project monitoring parameters, performance criteria, and potential corrective actions.

Monitoring Parameter	Final Performance Criteria used to determine Project Success	Interim Performance Criteria	Potential corrective actions or mid-course corrections*
Acquired Acres	Fee-simple acquisition of targeted habitats within project boundary	N/A	Funding allocated for fee-simple acquisition would be used to implement habitat restoration activities within project boundaries.
Vegetation Structure	 < 20% canopy cover of longleaf or slash pine 40 to 100% herbaceous cover Invasive nonnative plant species in any stratum present but sporadic (1-5 % cover) 	Performance criteria not met at year 5 after first treatment	 Change burn frequency Modify mechanical removal strategy Alter herbicide treatments Explore additional restoration alternatives (e.g., plantings) Continue to monitor.

Monitoring Parameter	Final Performance Criteria used to determine Project Success	Interim Performance Criteria	Potential corrective actions or mid-course corrections*
Vegetation composition	90% native flora	Performance criteria not met at year 5 after first treatment	 Change burn frequency Modify mechanical removal strategy Alter herbicide treatments Explore additional restoration alternatives (e.g., plantings) Continue to monitor.
Presence/Absence Henslow's sparrow	Presence	Presence	Adaptive management would follow that for native vegetation performance criteria
Diversity and abundance of injured bird species on targeted bottomland hardwood habitats	Use of target habitat by injured bird species	Performance criteria not met after 3 years	 Consider expanding survey area to document regional presence of survey bird species (are they in the area?). Conduct targeted monitoring on habitat metrics specific to wading bird habitat requirements.

^{*}The table provides the triggers for helping determine whether adjustments to the project are needed based on the performance criteria; potential corrective actions for unknown or unanticipated conditions should they arise would need to be determined.

6.0 Monitoring Schedule

The schedule for the project monitoring is shown in Table 6.0-1, separated by monitoring activity. Execution monitoring relates to baseline surveys (e.g., before habitat acquisition and/or management). Post-execution monitoring occurs in years following treatments (e.g., year 1 = within the first year following a prescribed burn).

Table 6.0-1. Monitoring Schedule.

Monitoring Parameters	Monitoring Timeframe ¹						
	Pre-Execution Monitoring	Execution Monitoring (initial)	Post-Execution Monitoring (years related to those following treatment)			ent)	
		As-built (Year 0)	Year 1	Year 2	Year 3	Year 4	Year 5
Acquired acres							
Presence, relative abundance, status, and distribution of invasive	Base-line surveys	х					х

Monitoring Parameters	Monitoring Timeframe ¹						
	Pre-Execution Monitoring	Execution Monitoring (initial)	Post-Execution Monitoring (years related to those following treatment)				
		As-built (Year 0)	Year 1	Year 2	Year 3	Year 4	Year 5
species within the project boundary							
Vegetation structure	Base-line surveys per protocol	X	X	X	X	X	X
Vegetation composition	Base-line surveys per protocol	Х	X	Х	X	X	Х
Henslow's sparrow presence/absence	Base-line surveys per protocol	Х	X	х	Х	Х	х
Diversity and abundance of injured bird species in targeted bottomland hardwood habitats.	Base-line surveys per protocol	x	X	х	х	х	х

7.0 Data Management

To the extent practicable, all environmental and biological data generated during monitoring activities would be documented using standardized field datasheets. If standardized datasheets are unavailable or not readily amendable to record project-specific data, then project-specific datasheets would be drafted prior to conducting any project monitoring activities. All tangible forms of field data would be reviewed by the Implementing Trustee for completeness and accuracy before being finalized. Original hardcopy datasheets and notebooks and photographs would be retained by the Implementing Trustee.

All field datasheets and notebook entries would be scanned to PDF files and would be archived along with the hardcopy datasheets. Electronic data files should be named with the date on which the file was created and should include a ReadMe file that describes when the file was created and by whom, and any explanatory notes on the file contents. If a data file is revised, a new copy should be made and the original preserved.

Relevant Project data that are handwritten on hardcopy datasheets or notebooks would be transcribed (entered) into Excel spreadsheets (or similar digital format). After transcription of the data, a second person not associated with data transcription would perform a verification of the data in the electronic data sheets against the original hardcopy datasheets and/or notebooks, and would make any corrections to transcription errors as appropriate before data are used for any analyses or distributed outside of the agency. Implementing Trustees would verify and validate monitoring data and information and would ensure that all data is entered or converted into agreed upon/commonly used digital format labeled with metadata.

All data would have properly documented FGDC/ISO metadata², a data dictionary (defines codes and fields used in the dataset), and/or a Readme file as appropriate (e.g., how data was collected, QA/QC procedures, other information about data such as meaning, relationships to other data, origin, usage, and format – can reference different documents).

7.1 Data Review and Clearance

Once data is entered electronically it is reviewed and verified for completeness. A quality check is made by verbally comparing the electronic data entered to the original hard copy data sheet. Data are validated and any corrections needed are made. Upon validation, data are approved for analysis, reporting and archiving. All data are kept in one permanent electronic folder as a permanent record.

After any and all identified errors are addressed, data are considered to be QA/QC'd.

The Implementing Trustee would give the other TIG members time to review the data before making such information publicly available. Before submitting the monitoring data and information package, co-Implementing Trustees shall confirm with one another that the package is approved for submission. No data release can occur if it is contrary to federal or state laws.

After any and all identified errors are addressed, data are considered to be QA/QC'd. The Implementing Trustee would give the other TIG members time to review the data before making such information publicly available. Before submitting the monitoring data and information package, co-Implementing Trustees shall confirm with one another that the package is approved for submission. No data release can occur if it is contrary to federal or state laws.

7.2 Data Storage and Accessibility

Trustees would provide DWH NRDA MAM data and information to the MS TIG and the Restoration Portal as soon as possible and no more than 1 year from when data are collected. Once all data has been QA/QC'd it would be submitted to the TIG and stored in the Restoration Project Database managed by the Trustees.

7.3 Data Sharing

Data would be made publicly available, in accordance with the Open Data Policy, through the DIVER Explorer Interface within a year of when the data collection occurred.

8.0 Reporting

All reporting would occur after field reconnaissance is complete for each assessment effort. This report would summarize the findings for the sampling period including all worksheets transferred into digital format and presented in tabular and graphical formats. The data should be summarized in such a way that it is meaningful to the reader. Additionally, an annual report would be completed that includes:

• Summary data –synthesized data for all efforts during the year

- Graphs vegetation characteristics, acres managed, bird species diversity and abundance, etc.
- Interpretation of graphical data
- Discuss comparison of data if pretreatment and post treatment data are available
- Explanation of results
- Uncertainties with management actions
- Potential data collection issues
- Issues to be resolved
 - Issues to improve data collection or cooperation in getting quality data
 - Issues associated with data loss or inability to collect data for a time period

9.0 Roles and Responsibilities

The Trustee Council would encourage and facilitate consistency in monitoring and data procedures to evaluate and report on progress toward meeting the goals of the project and are foundational to the PDARP/PEIS. The Mississippi TIG is responsible for addressing MAM objectives that pertain to their restoration activities and for communicating information to the Trustee Council or Cross-TIG MAM work group.

At the project level, USFWS, NOAA, and MDMR would be the implementing agencies. The implementing agency roles include coordination with project partners and the MS TIG to track project progress, program management and oversight, lead acquisition of parcels and co-lead for management operations.

10.0 Monitoring Budget

The overall budget for project monitoring and adaptive management is anticipated to be approximately 10-15% of the total project budget. This budget range is considered to be in draft form and is subject to change as project planning and implementation progress.

11.0 References

Clewell, A.F., B.S. Reed, M.E. Lasley. 1998. Long-term vegetational monitoring at the Mississippi Sandhill Crane National Wildlife Refuge, 1997. 35pp. Submitted to: Mississippi Sandhill Crane National Wildlife Refuge, Gautier, MS

Federal Geographic Data Committee. 2008. National vegetation classification standard, version 2. Document number FGDC-STD-005-2008. Pp221.

LoSchiavo, A. J., Best, R. G., Burns, R. E., Gray, S., Harwell, M. C., Hines, E. B., and Vearil, J. W. 2013. Lessons learned from the first decade of adaptive management in comprehensive Everglades restoration. Ecology and Society 18: 70.

Neckles, H. A., M. Dionne, D. M. Burdick, C. T. Roman, R. Buchsbaum and E. Hutchins. 2002. A monitoring protocol to assess tidal restoration of salt marshes on local and regional scales. Restoration Ecology 10: 556-563.

Neckles, H.A., G.R. Guntenspergen, W.G. Shriver, N.P. Danz, W.A. Wiest, J.L. Nagel, and J.H. Olker. 2013. Identification of metrics to monitor salt march integrity on National Wildlife

Refuges in relation to conservation and management objectives. Final Report to U.S. Fish and Wildlife Service, Northeast Region. USGS Patuxent Wildlife Research Center, Laurel, MD. 226 pp.

Nordman, C., R. White, R. Wilson, C. Ware, C. Ridout, M. Pyne, and C. Hunter. 2016. Rapid assessment metrics to enhance wildlife habitat and biodiversity within southern open pine ecosystems, Version 1.0. U.S. Fish and Wildlife Service and NatureServe, for the Gulf Coastal Plains and Ozarks Landscape Conservation Cooperative. March 31, 2016.

National Research Council. 2004. Adaptive Management for Water Resources Project Planning. National Academies Press, Washington, DC. 138pp.

Pastorok, R. A., MacDonald, A., Sampson, J. R., Wilber, P., Yozzo, D. J., & Titre, J. P. (1997). An ecological decision framework for environmental restoration projects. Ecological Engineering, 9(1), 89-107.

Shackelford, C.E., N.R. Carrie, C.M. Riley and D.K. Carrie. 2001. Project prairie birds: a citizen science project for wintering grassland birds. Second Edition. Texas Parks and Wildlife PWD BK W7000-485 (1/01). Booklet; 24pp.

Simenstad, C., Reed, D., & Ford, M. 2006. When is restoration not?: Incorporating landscape-scale processes to restore self-sustaining ecosystems in coastal wetland restoration. Ecological Engineering 26: 27-39.

Steyer, G. D., & Llewellyn, D. W. 2000. Coastal Wetlands Planning, Protection, and Restoration Act: A programmatic application of adaptive management. Ecological Engineering, 15(3), 385-395.

Wilder, S. 2016. Initial survey instructions: long-term vegetation monitoring-life form (Clewell plots). Submitted to: Mississippi Sandhill Crane National Wildlife Refuge, Gautier, MS.

Williams, B. K. 2011. Adaptive management of natural resources—framework and issues. Journal of Environmental Management, 92(5), 1346-1353.

Williams, B. K., & Brown, E. D. 2012. Adaptive management: the US Department of the Interior applications guide. US Department of the Interior, Adaptive Management Working Group. 136pp.

1.1 APPENDIX 1

Field Form for Rapid Assessment Metrics for Wildlife and Biodiversity in Southern Open Pine Ecosystems

Date:	Project:	Site ID:	
Field Crew Team Members:			
Leader:			
Assistants:			
		 AA Centrum out: _N _E_ S _W ;Buffer in	: _N _E
_S _W; Add'l: Y / N			
Photo filenames:			
(fill in values, units)	adius 18m, 40m, m/ft. o	r rectangle m/ft wide xm/ft	long
State:County:		Section:USGS 7.5	
Quad: Landowner/Managed Area Nam		Contact Person:	
Stand Name: Pe	 rmit Required? Locked Gat	e? Access Difficulties?	
(describe)			
SITE DESCRIPTION:			

Notes:


GENERAL DRAWING				
(Optional): Provide a				
drawing of the assessment				
area, including its				
boundaries, either aerial view or transect view.				
view of transect view.				
LOCATION: Assessment A	rea CENTRUM (check one	e) ORIGINAL MOV	ED (why? how far?)	
LOCATION: Assessment A GPS Unit:	rea CENTRUM (check one GPS Filename:	e) ORIGINAL MOV	ED (why? how far?) Projection:	
	<u> </u>	ORIGINAL MOV		# of Sat's:
GPS Unit:	GPS Filename:	•) ORIGINAL MOV	Projection:	
GPS Unit:	GPS Filename: Datum: NAD83		Projection:	
GPS Unit: UTM Zone:	GPS Filename: Datum: NAD83 WGS84	GPS Accuracy: m/ ft	Projection: PDOP:	# of Sat's:
GPS Unit: UTM Zone: UTM X Easting: — — — — —	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree	GPS Accuracy: m/ ft	Projection: PDOP:	# of Sat's:
GPS Unit: UTM Zone:	GPS Filename: Datum: NAD83 WGS84	GPS Accuracy: m/ ft	Projection: PDOP:	# of Sat's:
GPS Unit: UTM Zone: UTM X Easting: — — — — —	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree	GPS Accuracy: m/ ft	Projection: PDOP:	# of Sat's:
GPS Unit: UTM Zone: UTM X Easting: UTM Y Northing:	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft Original (GRTS):	Projection: PDOP: Field:	# of Sat's: Post-processed:
GPS Unit: UTM Zone: UTM X Easting: —————— UTM Y Northing: ———————— Classification (use to se	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft	Projection: PDOP: Field:	# of Sat's: Post-processed:
GPS Unit: UTM Zone: UTM X Easting: UTM Y Northing:	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft Original (GRTS):	Projection: PDOP: Field:	# of Sat's: Post-processed:
GPS Unit: UTM Zone: UTM X Easting: —————— UTM Y Northing: ———————— Classification (use to se	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft Original (GRTS):	Projection: PDOP: Field:	# of Sat's: Post-processed:
GPS Unit: UTM Zone: UTM X Easting: UTM Y Northing: Classification (use to se Southern Open Pine Grou	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft Original (GRTS): rn Open Pine Metrics Dat	Projection: PDOP: Field:	# of Sat's: Post-processed:
GPS Unit: UTM Zone: UTM X Easting: —————— UTM Y Northing: ———————— Classification (use to se	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft Original (GRTS): rn Open Pine Metrics Dat	Projection: PDOP: Field:	# of Sat's: Post-processed:
GPS Unit: UTM Zone: UTM X Easting: UTM Y Northing: Classification (use to se Southern Open Pine Ground Community Classification)	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft Original (GRTS): rn Open Pine Metrics Dat	Projection: PDOP: Field:	# of Sat's: Post-processed:
GPS Unit: UTM Zone: UTM X Easting: UTM Y Northing: Classification (use to se Southern Open Pine Ground Other Community Classification) USNVC Association	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft Original (GRTS): rn Open Pine Metrics Dat	Projection: PDOP: Field:	# of Sat's: Post-processed:
GPS Unit: UTM Zone: UTM X Easting: UTM Y Northing: Classification (use to se Southern Open Pine Ground Community Classification)	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft Original (GRTS): rn Open Pine Metrics Dat	Projection: PDOP: Field:	# of Sat's: Post-processed:
GPS Unit: UTM Zone: UTM X Easting: UTM Y Northing: Classification (use to se Southern Open Pine Ground Other Community Classification USNVC Association	GPS Filename: Datum: NAD83 WGS84 LAT: decimal degree LONG: decimal degree	GPS Accuracy: m/ ft Original (GRTS): rn Open Pine Metrics Dat	Projection: PDOP: Field:	# of Sat's: Post-processed:

1.2 APPENDIX 2

Wet Longleaf & Slash Pine Flatwoods & Savannas Metrics Data Sheet					Recorded Measured Value of Metric	Recorded Metric Score (1.0-4.0)
Canopy Metrics						
	Excellent = 4.0	Good = 3.0	Fair = 2.0	Poor = 1.0		
Canopy Southern	20-80 ft ² /acre	10 to <20 or >80 to	5 to <10 or 90 to <100	<5 or ≥100 ft²/acre	ft²/acre BA	
Yellow Pine Basal Area	basal area of longleaf or slash pine	<90 ft²/acre basal area of longleaf or slash pine	ft²/acre basal area of longleaf or slash pine	basal area of longleaf or slash pine		
						x0.25
Southern Yellow Pine Canopy Cover	20-65% canopy cover of longleaf or slash pine	15 to <20% canopy cover or >65-75% canopy cover of longleaf or slash pine	10 to <15% canopy cover or >75-85% canopy cover of longleaf or slash pine	<10% cover or >85% cover of longleaf or slash pine	% cover	
						x0.25
Southern Yellow Pine Stand Age Structure	BA ≥20 ft²/acre of flat-top longleaf or slash pine of any diameter and/or longleaf or	BA ≥10 ft²/acre of longleaf or slash pine trees ≥14" DBH class	Longleaf or slash pine trees ≥14" DBH class present, but at <10 ft²/acre BA	No longleaf or slash pine trees ≥14" DBH or with flat-top slash or longleaf pine	ft²/acre BA	
	slash pine trees ≥14" DBH class					x0.25
Ground Layer Metrics						
	Excellent = 4.0	Good = 3.0	Fair = 2.0	Poor = 1.0		
Overall Native Herbaceous Ground Cover	40-100% herbaceous cover	30 to <40% herbaceous cover	20 to <30% herbaceous cover	<20% herbaceous cover	% cover	
						x0.25

Page **21** of **21**

Invasive Plant	Invasive	Invasive nonnative	Invasive nonnative	Invasive nonnative	% cover	
Presence /	nonnative	plant species in any	plant species in any	plant species in any		
Distribution	plant species	stratum present but	stratum uncommon	stratum common		
	absent or	sporadic (1-5%	(5-10% cover)	(>10% cover)		
	cover is very	cover)				
	low (<u><</u> 1%					
	cover)					x0.25
Final Score is :					Ground	Layer
Canopy Scorex0.33 + Midstory Scorex0.33 + Ground Layer Scorex0.33 =					Score	e =
Evaluation	Evaluation Scale: 4.0 to 3.5 = Excellent, 3.5 to 2.5 = Good, 2.5 to 1.5 = Fair, 1.5 to 1.0 = Poor					



APPENDIX F Monitoring and Adaptive Management Plan for Deepwater Horizon NRDA Project:

Upper Pascagoula River Water Quality Enhancement

Page 2 of 22

Table	e of Contents	
1.0	Introduction	3
1.1	Project Overview	3
1.2	Project Goals and Restoration Objectives	4
1.3	Conceptual Model	5
1.4	Sources of Critical Uncertainty	6
2.0	Project Monitoring	7
3.0	Rationale for Adaptive Management	13
4.0	Evaluation	14
4.1	Project-Level Decisions	15
5.0	Monitoring Schedule	15
6.0	Data Management	16
6.1	Data Review and Clearance	17
6.2	Data Storage and Accessibility	19
6.3	Data Sharing.	19
7.0	Reporting	19
8.0	Roles and Responsibilities	21
9.0	Monitoring and Adaptive Management Budget	21
10.0	References	21

1.0 Introduction

Monitoring, Adaptive Management, and Administrative Oversight was identified as one of the programmatic goals in the *Deepwater Horizon* (*DWH*)oil spill Programmatic Damage Assessment and Restoration Plan and Programmatic Environmental Impact Statement (PDARP/PEIS). The *DWH* NRDA Monitoring and Adaptive Management Framework provides a flexible, science-based approach to effectively and efficiently implement restoration over several decades that provides long-term benefits to the resources and services injured by the spill. Project monitoring and adaptive management is important to measure the beneficial impacts of restoration and support restoration decision-making. This project Monitoring and Adaptive Management plan identifies the monitoring needed to evaluate progress toward meeting project objectives and to support adaptive management of the restoration project. It identifies key sources of uncertainty, incorporates monitoring data and decision points that address these uncertainties, and establishes a decision-making process for making adjustments where needed. This MAM Plan is a living document and would be updated as needed to reflect changing conditions and/or new information. Any future revisions to this document would be made publicly available through the Restoration Portal.

1.1 Project Overview

The health of the Gulf of Mexico depends upon the health of its estuaries, and the health of those coastal waters is influenced by land use upstream along tributary rivers. The primary goal for this project is water quality improvement through nutrient reduction. This watershed-scale project restores water quality impacted by the DWH oil spill by reducing nutrient load contributions and the sediment carrying them into coastal waters. Runoff from cropland, and pastureland contributes nutrients and sediment that adversely impact the health of coastal waters of the Gulf. While agricultural and pasture lands are not the sole contributors (and in many instances, not the leading contributors) of nutrients to coastal waters, there are tremendous opportunities to address this resource concern at its sources in the Pascagoula basin. Given the success of USDA, NRCS Farm Bill programs and their strong acceptance by private landowners, there is a significant opportunity to implement conservation practices on private lands. The USDA-NRCS would provide outreach and technical assistance to voluntary participants (landowners), especially on the most vulnerable acres in the watersheds, to develop conservation plans and would use all available conservation practices typically planned and funded by USDA-NRCS programs. The project proposes to implement clusters of projects within the smallest watershed (to the extent practicable) with the goal of making a discernable difference in water quality (at the watershed level). While this targeted and concentrated approach is desired, the projects proponents understand the voluntary nature of conservation implementation and will strive to reach the critical sources within the watershed. The proposed conservation practices would reduce nutrient losses from the landscape; reduce nutrient loads to streams and downstream receiving waters; reduce water quality degradation in watersheds that could provide benefits to marine resources and benefits to coastal watersheds.

This project is being implemented as restoration for the *DWH* oil spill Natural Resource Damage Assessment (NRDA). As outlined within the *DWH* oil spill PDARP/PEIS, this restoration project falls under the following programmatic goal, restoration type, restoration approach, restoration technique, TIG, and restoration plan:

- Programmatic goals: Restore Water Quality
- Restoration type: Nutrient Reduction (Non-Point source)
- Restoration approach: Reduce nutrient loads to coastal watersheds
- Restoration techniques: Agricultural conservation practices; Forestry management practices
- TIG: Mississippi Restoration Area
- Restoration plan: Mississippi Trustee Implementation Group Draft 2016-2017 Restoration Plan/Environmental Assessment (Draft RP/EA)

This restoration project is being implemented within the Pascagoula River basin (HUC 6, 031700) and more specifically in the Chunky-Okatibbee subbasin (HUC 8, 03170001). Subwatersheds include: Chunky River watershed (HUC 10, 0317000157); Tallahatta Creek watershed (HUC 10, 0317000158); Upper Okatibbee Creek watershed (HUC 10, 0317000159) (Note: much of this watershed upstream of a dam was excluded from GIS analysis); Okatibbee Creek watershed (HUC 10, 0317000160); Sowashee Creek watershed (HUC 10, 0317000161).

The monitoring of project parameters is dependent on the voluntary participation by landowners to implement conservation practices on their land. Implemented conservation practices may or may not be located in the same subwatershed, therefore sampling efforts may vary by scale at different watershed levels. The proposed conservation practices would reduce nutrient losses from the landscape, reduce nutrient loads to streams and downstream receiving waters, and reduce water quality degradation in watersheds that would provide benefits to marine resources and benefits to coastal watersheds. This project is intended to reduce nutrient and sediment loads contribution in watersheds that contain Gulf sturgeon [Acipenser oxyrinchus desotoi]) critical habitat. The Gulf sturgeon is anadromous, spending much of its life in marine environments, but spawning in the Upper Pascagoula River and tributaries. Sediment and other pollutants may reduce Gulf sturgeon spawning success. The Implementing Trustee is the Mississippi Department of Environmental Quality and the USDA-NRCS in partnership with the USEPA.

1.2 Project Goals and Restoration Objectives

Under the Restore Water Quality Programmatic Goal, the MS TIG would focus on the Nutrient Reduction (Nonpoint Source) Restoration Type, and specific goals of the Restoration Type:

- 1) Reduce nutrient loadings to Gulf Coast estuaries, habitats, and resources that are threatened by chronic eutrophication, hypoxia, or harmful algal blooms or that suffer habitat losses associated with water quality degradation.
- 2) Where appropriate, co-locate nutrient load reduction projects with other restoration projects to enhance ecological services provided by other restoration approaches.
- 3) Enhance ecosystem services of existing and restored Gulf Coast habitats.

The specific objectives for this project are:

Objective 1: Reduce sediment, phosphorus, and nitrogen loads during storm events leaving private lands in prioritized watersheds in the Pascagoula Basin;

Objective 2: *Identify in-stream habitat features that are influenced by upstream sediment and nutrient loads for future in-stream resource benefits.*

Performance criteria would be used to determine restoration success or the need for corrective action in accordance with (15 CFR 990.55(b)(1)(vii)) and are outlined for each objective in Section 2.

1.3 Conceptual Model

The conceptual model here is intended to explain the general relationships among project activities and outcomes derived from the implementation. The implementation of conservation practices in agricultural and forestry landscapes are well known management actions that reduce nonpoint source pollutant loads of nutrients and sediment impacting downstream receiving waters (Kröger et al., 2015). Conservation practices would follow the USDA-NRCS paradigm of avoid, control, and trap. Thus, practices are designed to reduce erosion, slow runoff velocities, and increase hydraulic residence time within the field or tract, and/or edge of field, all which are imperative to the physical, chemical, and biological processes that decrease nutrient and sediment loadings (Barlow and Kröger, 2014). Utilizing model outputs as well as observational data, conservation practices can be targeted into small watershed areas to produce measurable decreases in nutrients and sediments from the field itself, as well as within the downstream receiving water body. Reducing nutrient and sediment loading to the system is imperative for the functionality of in-stream habitats that are used by aquatic organisms to fulfill critical life history cycles. Increased sediment and nutrient loading in streams can result in siltation of in-stream gravel beds, as well as in low-flow clear water conditions, the proliferation of algae and other periphyton on benthic substrates. Siltation and excessive periphyton growth can cover in-stream gravel beds, which are important spawning habitats for Gulf sturgeon. Targeting conservation practices in high sediment and nutrient yielding watersheds will reduce nutrient and sediment loads entering downstream receiving stream reaches. Habitat mapping to identify potential Gulf sturgeon spawning habitat, and eDNA studies to detect presence/absence of Gulf sturgeon, will be completed as part of this MAM plan in order to relate sediment and nutrient reduction practices to potential Gulf sturgeon presence and spawning activities on potential in-stream habitats.

Table 1.3-1 Conceptual Model for the Upper Pascagoula Water Quality Enhancement Project

Activity		Output		Short-term outcome			Long-term outcome		
Implement conservations practices in nutrient a sediment into receiver.	ion to reduce nd	•	Reduced nutrient and sediment loading into the system	•	Decrease in nutrient and sediment loadings in targeted watersheds	•	Enhancement of ecosystem services of Gulf coast habitats and living marine resources		

1.4 Sources of Critical Uncertainty

Critical uncertainties are defined as those that have the potential to impact or impede the decision-making process and the ability to achieve the restoration objective(s). Although many types of scientific and other uncertainties exist, the focus of uncertainty in this context is the uncertainty that affects the decisions being made for the project. Monitoring to resolve critical uncertainties affecting these decisions can allow for more effective expenditure of resources (e.g., optimized project selection) into the future as learning takes place. Further, the learning that takes place through monitoring allows corrective actions to be taken to improve project outcomes. If unresolved, the critical uncertainty may delay the time it takes to achieve the restoration objectives, hinder an implemented project's ability to fully achieve restoration objectives, or in the worst-case scenario, it may have the potential to cause a project to fail altogether, regardless of the corrective actions taken.

Based on information in the conceptual ecological model, potential critical uncertainties for the project were identified and evaluated. These critical uncertainties are shown in Table 1.4-1.

Table 1.4-1. Critical uncertainties that may affect success of the Upper Pascagoula River Water Quality Enhancement Project

Critical Uncertainty	Summary of Strategy to Resolve				
Conservation practices may not result in measurable change in the receiving waters	Conduct targeted in-stream monitoring at locations that are upstream and downstream of the conservation implementation area. Monitoring data would be used to refine future management actions.				
Conservation practices may not result in reduced sediment build-up on in-stream habitat features	Conduct targeted monitoring for gravel beds identified by benthic habitat mapping data. Monitoring data would be used to refine future managemen actions.				
Suitable habitat features for Gulf sturgeon may not exist in the project area	Conduct benthic mapping/sub-bottom profiling activities to locate in-stream gravel beds that may serve as spawning habitat for Gulf sturgeon; Conduct targeted monitoring for Gulf sturgeon presence using eDNA techniques in areas that have the potential to support spawning habitat. Data would be used to refine future management actions.				

2.0 Project Monitoring

The proposed monitoring for this restoration project was developed to evaluate project performance. The monitoring parameters, outlined below, are organized by project objective, with one or more monitoring parameters for each objective. Information is provided on the monitoring methods, timing and frequency, duration, sample size, and sites. In addition, example performance criteria for each parameter are identified (if applicable), including example corrective actions that could be taken if the performance criteria are not met. These parameters would be monitored at the restoration project site, in adjacent streams, and may also be monitored at appropriate reference and/or control sites to demonstrate how the restoration project is trending toward the performance criteria. The parameters listed below may or may not be tied to performance criteria and/or corrective actions. Project monitoring would be applied to the following objectives:

Objective 1: Reduce sediment, phosphorus, and nitrogen loads during storm events leaving private lands in prioritized watersheds in the Pascagoula Basin

Parameter # 1: Total suspended solids (mg/L) and Turbidity (NTU)

- a) Rationale: This parameter would be used to determine whether the conservation practices are successful at meeting Objective 1 of this project and is a required water quality constituent for the NRCS Edge of Field Water Quality Monitoring Data Collection and Evaluation Conservation Activity (Code 201) to measure sediment loads.
- b) Methods:
 - i. Edge of Field: In-situ water sample collection at site drainage locations using automated collection systems. The system scenario outlined in Edge-of-Field Water Quality Monitoring Data Collection and Evaluation (201) is considered the "typical system" designed to meet the stated purposes of edge-of-field water quality monitoring. Event Mean Concentration (EMC) and accurate flow (discharge) measurements are required for each runoff event. All systems must be capable of sampling runoff events throughout the year.
 - ii. In-stream: Fixed station parameter reading using a data sonde, under baseflow conditions when possible, using standard monitoring protocols would occur at appropriately located upstream and downstream stations that bracket portions of watersheds with conservation practices.
- c) Timing, Frequency, and Duration:
 - i. Edge of Field: Data would be collected for storm events using an automated sampler across a hydrograph. Sites would be visit at least once per week or on alternating weeks when sampling events are not anticipated to maintain equipment and ensure proper functioning of the collection system. After collection events, sites would be visited as soon as possible after sampling events to retrieve samples, inspect flow measurement and automated sampler function,

- and make necessary repairs. Excessive delay in retrieving water samples can result in changes to their chemical composition and thus inaccurate representation of actual water quality.
- ii. In-stream: Ten samples per year would be collected at one or more sets of one upstream and two downstream stations that bracket portions of watersheds with conservation practices. Samples when possible, would be taken at baseflow conditions.
- iii. Duration of the project: 5 years.
- d) Sample Size: A paired design would be used at each conservation practice implementation site monitored. The total number of sites is not yet determined. A paired approach provides for a determination of conservation practice effectiveness by comparing a control field and a treatment field that are similar in terms of soil, slope, vegetation, hydrology, initially receive identical management, and receive the same weather (e.g., precipitation events) (Clausen and Spooner 1993). Monitor both fields (watersheds) under identical crop and management conditions without any new practice implementation during the baseline period. Follow this with monitoring of both fields after conservation practice implementation in the treatment field. The monitoring regime (i.e., sample location, method, and frequency) must remain the same through both baseline and post-implementation periods.
- e) Sites: Conservation practice implementation would be dependent on the participation of landowners in the target watersheds described in section 1.1. Locations would be updated in the monitoring plan when landowners sign participation agreements with the NRCS. Site selection criteria would adhere to the guidelines stated in the NRCS Edge of Field Water Quality Monitoring Data Collection and Evaluation Conservation Activity (Code 201). The geographic scope of the in-stream monitoring design would depend on the location of lands enrolled in the conservation program. Where a large number of acres are co-located in a small watershed (e.g. HUC 12), the design would likely include one upstream station (could be optional depending on upstream conditions) and one or more downstream stations (near-field and further downstream) depending on the location of the cluster of conservation practices.

Performance Criteria: x kg of suspended sediments trapped from treatment site

f) Corrective Action: Actions would vary depending on the type of conservation practice that is implemented. Some conservation practices may require inspection and maintenance. Information on the operations and maintenance of conservation practices can be found at http://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs_143_026849.

Parameter #2 Total phosphorus (mg/L)

a) Rationale: This parameter would be used to determine whether the restoration actions are successful at meeting Objective 1. This parameter is a required water quality constituent

for the NRCS Edge of Field Water Quality Monitoring Data Collection and Evaluation Conservation Activity (Code 201).

b) Method(s):

- i. Edge of Field: In-situ water sample collection at site drainage locations using automated collection systems. The system scenario outlined in Edge-of-Field Water Quality Monitoring Data Collection and Evaluation (201) is considered the "typical system" designed to meet the stated purposes of edge-of-field water quality monitoring. Event Mean Concentration (EMC) and accurate flow (discharge) measurements are required for each runoff event. All systems must be capable of sampling runoff events throughout the year.
- ii. In-stream: Sample collection using standard monitoring protocols would occur at appropriately located upstream and downstream stations that bracket portions of watersheds with conservation practice(s).

c) Timing, Frequency, and Duration:

- i. Edge of Field: Data would be collected for storm events using an automated sampler across a hydrograph. Sites would be visit at least once per week or on alternating weeks when sampling events are not anticipated to maintain equipment and ensure proper functioning of the collection system. After collection events, sites would be visited as soon as possible after sampling events to retrieve samples, inspect flow measurement and automated sampler function, and make necessary repairs. Excessive delay in retrieving water samples can result in changes to their chemical composition and thus inaccurate representation of actual water quality.
- ii. In-stream: Ten samples per year would be collected at one or more sets of one upstream and two downstream stations that bracket portions of subwatersheds, especially where conservation practices are densely co-located.
- iii. Duration of the project: 5 years.

d) Sample Size:

i. Edge of Field: A paired design would be used at each conservation practice implementation site. The total number of sites is not yet determined. A paired approach provides for a determination of conservation practice effectiveness by comparing a control field and a treatment field that are similar in terms of soil, slope, vegetation, hydrology, initially receive identical management, and receive the same weather (e.g., precipitation events) (Clausen and Spooner 1993). Monitor both fields (watersheds) under identical crop and management conditions without any new practice implementation during the baseline period. Follow this with monitoring of both fields after conservation practice implementation in the treatment field. The monitoring regime (i.e., sample location, method, and

- frequency) must remain the same through both baseline and post-implementation periods.
- ii. In-stream: Samples for MDEQ's Ambient Fixed Station Monitoring QAPP (MDEQ 2015) would be collected off bridges where possible that cross the water bodies in question or in wadeable streams if needed. The total number of sites is not yet determined and would be dependent on the amount and location of conservation practices in the targeted watersheds. It is anticipated that a total of 10 samples would be collected per year at each station. Samples would be taken at baseflow conditions when possible.
- e) Sites: Conservation practice implementation would be dependent on the participation of landowners in the target watersheds described in section 1.1. Locations would be updated in the monitoring plan when landowners sign participation agreements with the NRCS. Site selection criteria would adhere to the guidelines stated in the NRCS Edge of Field Water Quality Monitoring Data Collection and Evaluation Conservation Activity (Code 201). The geographic scope of the in-stream monitoring design would depend on the location of lands enrolled in the conservation program. Where a large number of acres are co-located in a small watershed (e.g. HUC 12), the design would likely include one upstream station (could be optional depending on upstream conditions) and one or more downstream stations (near-field and further downstream) depending on the location of the cluster of conservation practices.

Performance Criteria: x kg of phosphorus trapped from treatment site

d) Corrective Action: Actions would vary depending on the type of conservation practice that is implemented. Some conservation practices may require inspection and maintenance. Information on the operations and maintenance of conservation practices can be found at http://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs_143_026849

Parameter #3 Total Nitrogen (mg/L)

a. Rationale: This parameter would be used to determine whether the restoration actions are successful at meeting Objective 1. This parameter is a required water quality constituent for the NRCS Edge of Field Water Quality Monitoring Data Collection and Evaluation Conservation Activity (Code 201).

b. Method(s):

i. Edge of Field: In-situ water sample collection at site drainage locations using automated collection systems. The system scenario outlined in Edge-of-Field Water Quality Monitoring – Data Collection and Evaluation (201) is considered

- the "typical system" designed to meet the stated purposes of edge-of-field water quality monitoring. Event Mean Concentration (EMC) and accurate flow (discharge) measurements are required for each runoff event. All systems must be capable of sampling runoff events throughout the year.
- ii. In-stream: Sample collection using standard monitoring protocols would occur at appropriately located upstream and downstream stations that bracket portions of subwatersheds, especially where conservation practices are densely co-located.

c. Timing, Frequency, and Duration:

- i. Edge of Field: Data would be collected for storm events using an automated sampler across a hydrograph. Sites would be visit at least once per week or on alternating weeks when sampling events are not anticipated to maintain equipment and ensure proper functioning of the collection system. After collection events, sites would be visited as soon as possible after sampling events to retrieve samples, inspect flow measurement and automated sampler function, and make necessary repairs. Excessive delay in retrieving water samples can result in changes to their chemical composition and thus inaccurate representation of actual water quality.
- ii. In-stream: Ten samples per year would be collected at one or more sets of one upstream and two downstream stations that bracket portions of subwatersheds, especially where conservation practices are densely co-located. Samples would be taken at baseflow conditions when possible.
- iii. Duration of the project: 5 years.

a) Sample Size:

- i. Edge of Field: A paired design would be used at each conservation practice implementation site. The total number of sites is not yet determined. A paired approach provides for a determination of conservation practice effectiveness by comparing a control field and a treatment field that are similar in terms of soil, slope, vegetation, hydrology, initially receive identical management, and receive the same weather (e.g., precipitation events) (Clausen and Spooner 1993). Monitor both fields (watersheds) under identical crop and management conditions without any new practice implementation during the baseline period. Follow this with monitoring of both fields after conservation practice implementation in the treatment field. The monitoring regime (i.e., sample location, method, and frequency) must remain the same through both baseline and post-implementation periods.
- ii. In-stream: Samples for MDEQ's Ambient Fixed Station Monitoring QAPP (MDEQ 2015) would be collected off bridges where possible that cross the water bodies in question or in wadeable streams if needed. The total number of sites is not yet determined and would be dependent on the amount and location of conservation practices in the targeted watersheds. It is anticipated that a total of

- 10 samples would be collected per year at each station. Samples would be taken at baseflow conditions when possible.
- b) Sites: Conservation practice implementation would be dependent on the participation of landowners in the target watersheds described in section 1.1. Locations would be updated in the monitoring plan when landowners sign participation agreements with the NRCS. Site selection criteria would adhere to the guidelines stated in the NRCS Edge of Field Water Quality Monitoring Data Collection and Evaluation Conservation Activity (Code 201). The geographic scope of the in-stream monitoring design would depend on the location of lands enrolled in the conservation program. Where a large number of acres are co-located in a small watershed (e.g. HUC 12), the design would likely include one upstream station (could be optional depending on upstream conditions) and one or more downstream stations (near-field and further downstream) depending on the location of the cluster of conservation practices.
- c) Performance Criteria: x kg of nitrogen trapped from treatment site
- d) Corrective Action: Actions would vary depending on the type of conservation practice that is implemented. Some conservation practices may require inspection and maintenance. Information on the operations and maintenance of conservation practices can be found at http://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/technical/cp/ncps/?cid=nrcs_143_026849

Parameter # 4: Benthic Substrate

- a) Rationale: This parameter is required to identify potential suitable spawning habitat for Gulf sturgeon in the study area in order to document additional ecosystem benefits of the project. Benthic substrate types would be delineated to illustrate spatial heterogeneity of riverine substrates.
- b) Method: Utilize Swath Bathymetric and Sub-Bottom Profiling Systems that are capable of detecting locations of gravel beds and sub-surface materials. Habitats would be ground- truthed in areas that are wadeable for accuracy assessment.
- c) Timing, Frequency, and Duration: Data would be collected in one survey event to reduce potential variability in water volumes over time. Survey duration has to be determined and would depend on the number and location of conservation practice sites in the study area.
- d) Sample Size: 100% bathymetric coverage of waterways located adjacent to conservation practices identified in objective 1.

- e) Sites: Locations would be dependent on the locations of participating landowners in the target watersheds described in section 1.1. Locations would be updated in the monitoring plan when landowners sign participation agreements with the NRCS.
- f) Corrective Action: not applicable

Parameter # 5: Gulf sturgeon eDNA samples

- a) Rationale: This parameter is required to determine the presence and specific locations of Gulf sturgeon in the project area waterways.
- b) Method: Water samples would be collected at strategic locations using 150 meter transects. One liter water samples would be collected at 0, 75, and 150 meters along the transect. Environmental DNA collection methods would follow the procedure outlined by Pfleger et al. (2016). At each site, three sites replicates would be sampled. Quality control measures, such as sterile technique for collecting and decontamination would be taken at each site to avoid contamination and reduce the possibility of false positives. Collected samples would be immediately placed on ice in a sterilized source cooler storage container to prevent DNA degradation.
- c) Timing, Frequency, and Duration: Data would be collected annually during the spring migration for Gulf sturgeon (April-August). Single event sampling efforts would occur at each site once per month during the migration time period. Subsequent laboratory analysis would take place after samples have been collected and stored. Sampling would occur for five years.
- d) Sample Size: Transect samples would be collected at a broad level to cover the entire waterways of the Okatibbee and Chunky rivers. Sampling would occur every 5 kilometers from the mouth of each waterway, upstream until waters become unnavigable. This includes approximately 60 km of the Okatibbee and approximately 40 km of the Chunky. The sample size equals 12 and 8, respectively. Over a 5-month period (migration), the total number of sample events equals 100. Locations would be refined for sampling after benthic habitat data has been analyzed for potential spawning habitat. eDNA sampling efforts would then concentrate specifically on these areas.
- e) Sites: Locations would be dependent on the locations of participating landowners in the target watersheds described in section 1.1. Locations would be updated in the monitoring plan when landowners sign participation agreements with the NRCS.

3.0 Rationale for Adaptive Management

Implementation of the conservation practices, benthic investigations and eDNA monitoring and evaluation would utilize standardized actions using accepted tools and protocols at specific

locations (NRCS, 2012; Pfleger, 2016). Although adaptive management is a critical component of the restoration plan as a whole, the need for adaptive management on specific conservation practices being implemented is not needed for this project due to the nature of the sampling approaches, the objectives of the project, and the scale of the sites in which the data would be collected (crop field scale; waterway segment scale), and an understanding of the conservation practices that would be applied. Data, analysis and information obtained from this project would be used to help inform future Restoration Plan development, priorities and project selection.

4.0 Evaluation

Evaluation of monitoring data is needed to assess the performance of the project in meeting its restoration objectives, resolving uncertainties to increase understanding, and determine whether corrective actions are needed.

The results of the analysis would be used to answer the following questions:

- Were the project restoration objectives achieved? If not, is there a good reason why they were not met?
- Did the restoration project produce unanticipated effects?
- Were there unanticipated events unrelated to the restoration project that potentially affected the monitoring results?
- Were any of the uncertainties identified prior to project implementation resolved?
- Were any new uncertainties identified?

The analysis methods would be applied to all monitoring parameters as follows:

Water Quality Data

Edge of Field:

Paired field data are often analyzed by Analysis of Covariance (ANCOVA), a procedure that combines linear regression with Analysis of Variance (ANOVA) (Grabow et al. 1998). The USEPA recommends an ANCOVA model for paired watershed data analysis, using matched event loads from control and treatment watersheds to determine effects of conservation practices (USEPA, 1993; USEPA, 1997).

In-stream:

Standard analytical techniques would be used to document water quality improvements between upstream and downstream stations that bracket project areas with conservation systems and follow the guidelines provided in MDEQ's Ambient Fixed Station Monitoring quality assurance project plan (QAPP) (MDEQ 2015). This QAPP has been prepared according to the requirements and guidance provided in the following documents:

USEPA Requirements for Quality Assurance Project Plans, (USEPA QA/R-5 EPA/240/B-01/003)) U.S. Environmental Protection Agency, Office of Environmental Information, Washington, D.C., March 2001 and USEPA Guidance for Quality Assurance Project Plans, (USEPA QA/G5 EPA/240/R-02/009), U.S. Environmental Protection Agency, Office of Environmental Information, Washington, D.C., December 2002.

Benthic Habitat Mapping

Multibeam bathymetric data would be processed in CARIS HIPS or similar software and delivered in Bathymetric Attributed Grid (BAG) format. Grid resolution would be 0.25 meters. Depth, uncertainty, and coverage data would be included with each BAG. The gridded bathymetry data would be used to generate an ESRI shapefile of 1 foot contours. Bathymetric grids would also be used for 3D volumetric analyses.

Multibeam backscatter data would be processed in CARIS SIPS and delivered in georeferenced tagged image file format (GeoTIFF). Sub-bottom profiler data would be processed and interpreted in SonarWiz, or similar software. Along-track seismic reflection profiles would be delivered in TIFF image format. An ESRI shapefile of the sub-bottom profiler trackline would be included. Raw sub-bottom profiler files would also be delivered in native JSF format. Acreages of gravel beds would be delineated in a GIS system using the processed data.

eDNA

DNA extraction methods should follow the best available science. Procedures outlined in Pfleger (2016) resulted in positive DNA hits for both Gulf sturgeon and Alabama sturgeon. Specific numbers of positives and negatives by site per month would be documented and graphed so that a comparative analysis of sites and timing (month) can be analyzed. Additionally, site characteristics would be analyzed with benthic habitat data to better understand species presence and habitat relationships.

4.1 **Project-Level Decisions**

The need for adaptive management on specific conservation practices being implemented beyond inspection and maintenance is not needed for this project. Monitoring information from this restoration project would be critical to refine targeting of conservation practice implementation, refine potential in-stream habitat that could be used by Gulf sturgeon if found, as well as identify in-stream habitat that could be enhanced by conservation practices for Gulf sturgeon as needed.

5.0 Monitoring Schedule

The schedule for the project monitoring is shown in Table 5.0-1, separated by monitoring activity. Execution monitoring occurs when the project has been fully executed as planned (Year 0).

<u>Performance Monitoring for Objective 1:</u> The monitoring of project parameters is dependent on the voluntary participation by landowners to implement conservation practices. Performance monitoring would occur in the years following initial project execution (Years 1-5), but is restrained by the five-year duration of the overall project. The length of time a conservation practice is monitored is contingent on when the treatment is executed within project timeline. Thus, treatments may receive monitoring for 1-5 years. However, it is anticipated that project sites would execute treatments in the second year following project planning and outreach to landowners. The monitoring schedule would be updated as conservation practices are planned and implemented.

Benthic habitat mapping and eDNA sampling: Benthic habitat mapping would occur in year 1 of the project and would be conducted as a single event. eDNA sampling would occur every 5 kilometers from the mouth of each waterway, upstream until waters become unnavigable in year 1 of the project for the broad level analysis of potential Gulf sturgeon presence. Locations would be refined for sampling after benthic habitat data has been analyzed for potential spawning habitat. eDNA sampling efforts would then concentrate specifically on these areas in subsequent monitoring years to account for potential inter-annual migration shifts. eDNA sampling would occur annually for the duration of the project to maximize detection of the potential presence of Gulf sturgeon and the relationship of that potential presence to possible habitat use by Gulf sturgeon, in response to in-stream changes from conservation practice implementation.

Table 5.0-1. Monitoring Schedule.

	Monitoring Timeframe								
Monitoring Parameters	Execution Monitoring (initial)	Post-Execution Monitoring (ongoing)							
	As-built (Year 0)	Year 1	Year 2	Year 3	Year 4	Year 5			
Parameters 1,2,3		X	X	X	X	X			
Parameter 4		X							
Parameter 5		X	X	X	X	X			

6.0 Data Management

To the extent practicable, all environmental and biological data generated during monitoring activities would be documented using standardized field datasheets. If standardized datasheets are unavailable or not readily amendable to record project-specific data, then project-specific datasheets would be drafted prior to conducting any project monitoring activities. All tangible forms of field data would be reviewed by Implementing Trustee for completeness and accuracy before being finalized. Original hardcopy datasheets and notebooks and photographs would be retained by the Implementing Trustee.

All field datasheets and notebook entries would be scanned to PDF files and would be archived along with the hardcopy datasheets. Electronic data files should be named with the date on which

the file was created and should include a ReadMe file that describes when the file was created and by whom, and any explanatory notes on the file contents. If a data file is revised, a new copy should be made and the original preserved.

Relevant project data that are handwritten on hardcopy datasheets or notebooks would be transcribed (entered) into Excel spreadsheets (or similar digital format). After transcription of the data, a second person not associated with data transcription would perform a verification of the data in the electronic data sheets against the original hardcopy datasheets and/or notebooks, and would make any corrections to transcription errors as appropriate before data are used for any analyses or distributed outside of the agency. Implementing Trustees would verify and validate monitoring data and information and would ensure that all data is entered or converted into agreed upon/commonly used digital format labeled with metadata.

6.1 Data Review and Clearance

All components of this project would be subject to the formal Quality Management Program developed by Mississippi Department of Environmental Quality (MDEQ 2014). This program dictates that all data collection and monitoring efforts be performed under a project specific Quality Assurance Project Plan (QAPP). To meet this requirement, MDEQ would provide a Comprehensive Quality Assurance Plan (CompQAP) for all the project activities. Quality Assurance procedures for this monitoring plan, all field methods and associated data collection, recording and storage efforts would be included in the CompQAP. This document would be used to ensure that environmental and related data collected, compiled, and/or generated for this project are of the type, quantity, and quality required for their intended purpose.

Water Quality Data Collection

Edge of Field:

Data would be QA/QC'd in accordance with procedures outlined in the NRCS Edge of Field Water Quality Monitoring Data Collection and Evaluation Conservation Activity (Code 201). A QAPP is required for NRCS-assisted water quality monitoring and must be used as the basis of the QAPP when NRCS is the lead funding agency. Among other items, a QAPP would fully describe the process of sample preservation, handling, and processing. The QAPP documents the results of a project's technical planning process, providing in one place a clear, concise, and complete plan for the environmental data operation and its quality objectives and identifying key project personnel.

In-stream:

Appropriate QA procedures from MDEQ's Ambient Fixed Station Monitoring quality assurance project plan (QAPP) (MDEQ 2015) would be used for in-stream monitoring. This QAPP presents the sampling, analytical, QC requirements for the Ambient Fixed Station Monitoring program conducted under the CWA §106. The QAPP requirements are designed to ensure reproducible and defensible data are generated for use in surface water assessments

Benthic Mapping Data Collection

The quality of hydrographic data depends on precise calibration and maintenance of the accuracy through automatic calibration techniques and periodic verification of the results through data monitoring and statistical analyses of data sets. The quality control system for this project is designed to continuously monitor data quality and query system conditions, which allows for the delivery of high-quality data products. Thus, in addition to the quality control plan described below, there would be near real-time quality control of data in the field as it is acquired. A QA/QC plan for hydrographic data collection would be required before data collection occurs.

eDNA Collection

Protocols should adhere to Mahon et al. (2010) Environmental DNA Monitoring and Surveillance: Standard Operation Procedures. Report to the United States Army Corps of Engineers, Environmental Laboratories, and Cooperative Environmental Studies Unit, Vicksburg, Mississippi or similar protocol that is applicable to the habitat type. Additionally, numerous scientific manuscripts outline protocols for data methods, quality and control.

Data would be QA/QC'd in accordance with procedures outlined in the QA/QC Clearance and Release document approved by the Trustees.

To the extent practicable, all environmental and biological data generated during monitoring activities would be documented using standardized field datasheets. If standardized datasheets are unavailable or not readily amendable to record project-specific data, then project-specific datasheets would be drafted prior to conducting any project monitoring activities. All tangible forms of field data would be reviewed by Implementing Trustee for completeness and accuracy before being finalized. Original hardcopy datasheets and notebooks and photographs would be retained by the Implementing Trustee.

All field datasheets and notebook entries would be scanned to PDF files and would be archived along with the hardcopy datasheets. Electronic data files should be named with the date on which the file was created and should include a ReadMe file that describes when the file was created and by whom, and any explanatory notes on the file contents. If a data file is revised, a new copy should be made and the original preserved.

Relevant Project data that are handwritten on hardcopy datasheets or notebooks would be transcribed (entered) into Excel spreadsheets (or similar digital format). After transcription of the data, a second person not associated with data transcription would perform a verification of the data in the electronic data sheets against the original hardcopy datasheets and/or notebooks, and would make any corrections to transcription errors as appropriate before data are used for any analyses or distributed outside of the agency. Implementing Trustees would verify and validate MAM data and information and would ensure that all data is entered or converted into agreed upon/commonly used digital format labeled with metadata following FGDC/ISO standards to the extent practicable and in accordance with individual agency requirements.

After any and all identified errors are addressed, data are considered to be QA/QC'd. The Implementing Trustee would give the other TIG members time to review the data before making such information publicly available. Before submitting the monitoring data and information package, co-Implementing Trustees shall confirm with one another that the package is approved for submission. No data release can occur if it is contrary to federal or state laws.

6.2 Data Storage and Accessibility

Once all data has been QA/QC'd it would be submitted to the Restoration Project Database that is held at MDEQ.

6.3 Data Sharing

Data would be made publicly available, in accordance with the Open Data Policy, through the DIVER Explorer Interface within a year of when the data collection occurred.

7.0 Reporting

Water Quality

Edge of Field:

For each water quality station, rainfall and flow data would accompany electronic (.pdf) copies of the laboratory analysis for each event. Weekly or bi-weekly checklists and/or a log book should provide information about the performance of the monitoring system, specifically noting any malfunctions, gaps in data collection, or conditions that might be useful in interpreting the results of collected data. The operations form should be completed for the reporting period. Weekly or bi-weekly photos of the field and the system would be provided digitally. An Excel spreadsheet containing all water quality data for all the events of the reporting period would be submitted. All information in this paragraph is required as the documentation for a semi-annual data submittal.

The annual submittal includes all requirements of a semi-annual data submittal for the second half of the monitoring year. In addition, this report would summarize the findings for the year and would include a status review with the participant. The data should be summarized in such a way that it is meaningful to the participant. NRCS must complete a quality assurance check of existing practice management known as the Annual Field Check form. All information in this paragraph is required as the documentation for an annual submittal. The report should include:

- Summary data Tabular
- Graphs Discharge (cfs), Runoff (inches) and Load (lbs/acre)
- Interpretation of graphical data
- Discuss comparison of control and treatment sites
- Explain Results
- Event mean concentration (EMC) vs. discharge

- Unexpected events (data outliers)
- Explain the difference between nutrient inputs and nutrient loads leaving the field (lb/acre)
- Potential data collection issues
- Issues to be resolved
 - Issues to improve data collection or cooperation in getting quality data
 - Issues associated with data loss or inability to collect data for a time period (due diligence)

In-stream:

Field data, field observations and analytical data would be compiled and presented via paper and electronic means. Reporting would include: methodology, including describing field and analytical methods; tabulation of analytical results and field measurements, a QA/QC summary; and a discussion addressing problems, corrective actions, or other characteristics of the data that are required for scientifically sound interpretation of the data. This information would be provided to the project administration by the PM and the Project Data Manager on a quarterly basis.

The monitoring reports would be prepared for each sample collected and include the following items:

- Site identification and location information;
- *in situ* field measurements:
- Analytical results, including analytical methods and dates of analyses; and
- Any additional observations recorded on the sample collection field forms.

Data Quality Reports would include the following:

- Summary of analytical results, including a summary of QA/QC data (i.e., results of field duplicates, analytical duplicates, spikes, and blanks);
- Methods of data analysis; and
- Tabular summaries of all direct and non-direct measurements.

Benthic Habitat Mapping

Progress reports would be submitted to the project lead on a weekly basis to detail progress to date, in addition to current and anticipated survey schedule. A Data Acquisition and Processing Report (DAPR) would be submitted along with mapping deliverables after the conclusion of the field effort.

eDNA

Progress reports would be submitted to the project lead on a weekly basis to detail progress to date, in addition to current and anticipated sampling schedule. A final laboratory report would be submitted that includes all raw data and analysis results.

8.0 Roles and Responsibilities

The Trustee Council would encourage and facilitate consistency in monitoring and data procedures to evaluate and report on progress toward meeting the ecosystem goals that are the stated basis for the Restoration Agreement and are foundational to the PDARP/PEIS. The Mississippi TIG is responsible for addressing MAM objectives that pertain to their restoration activities and for communicating information to the Trustee Council or Cross-TIG MAM work group.

At the project level, Implementing Trustees include MDEQ, USDA, and the USEPA. The Implementing Trustee would work with USDA's NRCS on the project and would perform landowner outreach activities and implementation of conservation practices in targeted watersheds. MDEQ's primary roles includes coordination with project partners and the MS TIG to track project progress, program management and oversight, lead for edge of field sampling, and provide a Comprehensive Quality Assurance Plan (CompQAP) for all of the project activities. USEPA would provide coordination support as well as take the lead on in-stream water quality monitoring in the field under the MDEQ CompQAP; the MDEQ laboratory would process and analyze the total nitrogen and total phosphorus samples taken in-stream.

9.0 Monitoring and Adaptive Management Budget

The overall budget for project monitoring is anticipated to be approximately 10-15% of the total project budget. This budget range is considered to be in draft form and is subject to change as project planning and implementation progress. Adaptive management is not a component of this project and is not included in the budget.

10.0 References

Barlow, J. R., & Kröger, R. (2014). Nitrogen transport within an agricultural landscape: Insights on how hydrology, biogeochemistry, and the landscape intersect to control the fate and transport of nitrogen in the Mississippi Delta. Journal of Soil and Water Conservation, 69(1), 11A-16A.

Clausen, J. C., & Spooner, J. (1993). Paired watershed study design (No. PB-94-154820/XAB; EPA--841/F-93/009). Environmental Protection Agency, Washington, DC (United States). Office of Wetlands, Oceans and Watersheds.

Grabow, G. L., Spooner, J., Lombardo, L. A., Line, D. E., & Tweedy, K. L. (1998). Has water quality improved? Use of SAS for Statistical Analysis of Paired Watershed, Upstream/Downstream, and Before/After Monitoring Designs. Biological and Agricultural Engineering Department, North Carolina State University, Raleigh, NC. USEPA-NCSUCES Grant, (X825012).

Mahon, A., Rohly, A., Budny, M., Elgin, E., Jerde, C., Chadderton, W., Lodge, D. (2010). Environmental DNA Monitoring and Surveillance: Standard Operation Procedures. Report to the United States Army Corps of Engineers, Environmental Laboratories, Cooperative Environmental Studies Unit, and Vicksburg, Mississippi.

MDEQ. (2015). Quality Assurance Project Plan for Ambient Fixed Station Monitoring. This quality assurance project plan (QAPP) has been prepared according to the requirements and guidance provided in the following documents: USEPA Requirements for Quality Assurance Project Plans, (USEPA QA/R-5 EPA/240/B-01/003)) U.S. Environmental Protection Agency, Office of Environmental Information, Washington, D.C., March 2001 and USEPA Guidance for Quality Assurance Project Plans, (USEPA QA/G5 EPA/240/R-02/009), U.S. Environmental Protection Agency, Office of Environmental Information, Washington, D.C., December 2002.

Kröger, R., Prince Czarnecki, J. M., Tank, J. L., Christopher, S. F., & Witter, J. D. (2015). Implementing Innovative Drainage Management Practices in the Mississippi River Basin to Enhance Nutrient Reductions. JAWRA Journal of the American Water Resources Association, 51(4), 1020-1028.

USDA-NRCS (2012). Edge-of-field water quality monitoring data collection and evaluation—Conservation Activity Code 201.

Pfleger, M. O., Rider, S. J., Johnston, C. E., & Janosik, A. M. (2016). Saving the doomed: Using eDNA to aid in detection of rare sturgeon for conservation (Acipenseridae). Global Ecology and Conservation, 8, 99-107.

USEPA. (1993). Paired watershed study design. EPA 841-F-93-009.2001. USEPA Office of Water, Washington, DC

USEPA. (1997). Monitoring guidance for determining the effectiveness of nonpoint-source controls. EPA 841-B-96-004. Washington, D.C.